

GRAPH THEORETICAL REPRESENTATION OF KNOT SYMMETRIC ALGEBRA

*M. KAMARAJ and **R. MANGAYARKARASI

*Government Arts College, Melur-625 106, Maduraidt, Tamilnadu, India

**E. M. G. Yadava Women's college, Madurai 625 014, Tamilnadu, India

(Received on: 25-10-11; Revised & Accepted on: 22-11-12)

ABSTRACT


In this paper we introduce Graph theoretical representation of Knot symmetric Algebra.

INTRODUCTION

In [Br], Brauer algebra was introduced by Richard Brauer (1937) in connection with the finding irreducible representation of the orthogonal group. Generators of Brauer algebra were represented by a graph with $2n$ vertices arranged in two rows such that each row contains n vertices. In [KM], we introduced a new class of algebras which are known as Knot symmetric algebras. Brauerdiagram (graph) motivated as to represent every generator of Knot symmetric algebras as a special type of graph which we call them as Knot graphs.


1. PRELIMINARIES

Brauer algebras 1.1. For $k \in \mathbb{Z}$ and $x \in \mathbb{C}$, the Brauer algebra $B_k(x)$ is the algebra over \mathbb{C} whose basis consists of all diagrams on $2k$ vertices that have any combination of horizontal and vertical edges. An example of Brauer diagram is in below Fig 1



The dimension formula for $B_k(x)$ is $(2k-1)!$

Where $(2k-1)! = (2k-1)(2k-3).....3 \cdot 1$. Multiplying Brauer diagrams introduce a parameter x which comes in to play when a loop forms in the middle rows of two diagram being multiple. A loop can be formed by two or more horizontal edges in the middle rows. When this occurs the loops disappear and we multiply the resulting diagram by x^l where l is the number of loops in the middle rows. For example $n=6$ Note that horizontal and vertical edges can appear in the product of two diagrams via a sequences of edges that starts and ends with a vertical edge and which may have horizontal edges in the middle.



*Corresponding author: **R. MANGAYARKARASI

**E. M. G. Yadava Women's college, Madurai 625 014, Tamilnadu, India

Knot symmetric algebras 1.2

Let S_n denote the symmetric group of order n . Every element of S_n can be represented as Brauer diagram (graph) with $2n$ vertices and without horizontal edges [Br]. Let $\pi \in S_n$ the vertices of π are represented in two rows such that each row contains n vertices. The vertices of each row is indexed with $1, 2, \dots, n$ from left to right in order. Let $E(\pi)$ denote the set of all edges of π .

(i.e) $E(\pi) = \{e_i = (i, \pi(i)); 1 < i < n\}$

Define $S(\pi)$ is a subset of $E(\pi) \times E(\pi)$ such that $S(\pi) = \{(e_i, e_j), i < j\}$. It is obvious that $|S(\pi)| = \frac{n(n-1)}{2}$

Example. 1 $\begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 2 & 1 \end{pmatrix} \in S_4$ is represented by fig 1

For the Fig 1



$E(\pi) = \{e_1=(1,3), e_2=(2,4), e_3=(3,2), e_4=(4,1)\}$

$S(\pi) = \{(e_1, e_2), (e_1, e_3), (e_1, e_4), (e_2, e_3), (e_2, e_4), (e_3, e_4)\}$

Let f_π be a mapping from $S(\pi)$ to $\{-1, 0, 1\}$ such that

$$f_\pi(e_i, e_j) = \begin{cases} 0 & \text{if } \pi(i) < \pi(j) \\ 1 \text{ or } -1 & \text{if } \pi(i) > \pi(j) \end{cases}$$

and $f_\pi(e_i, e_j) + f_\pi(e_j, e_i) = 0$



Knot mapping 1.3. A mapping f_π defined above is called a Knot mapping. Refer the above Fig 2. We have

$e_1 = (1, 2), e_2 = (2, 3), e_3 = (3, 1)$

$f_\pi(e_1, e_2) = 0, f_\pi(e_1, e_3) = 1, f_\pi(e_3, e_1) = -1$

Knot Number 1.4 Define $K(\pi) = \{(e_i, e_j) \in S(\pi); \pi(i) > \pi(j)\}$

Definition 1.5 $|K(\pi)|$ is called Knot number of π . Let x be indeterminate. Define $N(\pi) = \{x^m f_\pi; m \in \mathbb{Z}, f_\pi \text{ is a Knot mapping}\}$

For any two Knot mapping f_π and g_π .

Define $E(f_\pi, g_\pi) = \{(e_i, e_j) \in K(\pi) : f_\pi(e_i, e_j) + g_\pi(e_i, e_j) = 0\}$

Knot Relation 1.6 Define a relation \sim in $N(\pi)$ such that $x^m f_\pi \sim x^l g_\pi$ if

- (i) $m = l$ and $f_\pi = g_\pi$ or
- (ii) $l - m = 2$ $\sum_{(e_i, e_j) \in E(f_\pi, g_\pi)} f_\pi(e_i, e_j)$ This relation is called Knot relation

Knot multiplication 1.7 Let $\overline{N(\pi)} = N(\pi) / \sim$ That is $\overline{N(\pi)}$ is the collection of disjoint equivalence classes with respect to the Knot relation. Define $T_n = \{(\pi, x^m f_\pi) : \pi \in S_n, f_\pi \in \overline{N(\pi)} \text{ and } m \text{ is an integer}\}$

We define multiplication in T_n as follows:

Let $a, b \in T_n$ and $a = (\pi, x^m f_\pi), b = (\sigma, x^l g_\sigma)$.

Define $ab = (\sigma \circ \pi, x^{m+l} h_{\sigma \circ \pi})$ where α and $h_{\sigma \circ \pi}$ are defined as follows:

Let $(e_i, e_j) \in S(\sigma \circ \pi), (u_i, u_j) \in S(\pi), (v_p, v_q) \in S(\sigma), p, q \in \{\pi(i), \pi(j)\}, f_\pi(u_i, u_j) = u$ and $g_\sigma(v_p, v_q) = v$.

Now $\alpha = \sum_{(e_i, e_j) \in S(\sigma \circ \pi)} \alpha(e_i, e_j)$

where $\alpha(e_i, e_j) = (u + v) |uv|$ and $h_{\sigma \circ \pi}(e_i, e_j) = (u + v)(1 - \delta_{u,v})$ where $\delta(u, v) = \begin{cases} 0 & \text{if } u \neq v \\ 1 & \text{if } u = v \end{cases}$

Theorem 1.8. The Knot multiplication is associative in T_n .


Theorem 1.9. FT_n is an algebra

This algebra is called as Knot symmetric algebra

2. KNOT GRAPHS


Let S_n be the symmetric group of order n and $\pi \in S_n$. A knot graph of order n is a special graph which is defined from π as follows.

Definition 2.1 we start with an element $\pi \in S_n$, π can be represented by a graph. Consider two edges $(i, \pi(i))$ and $(j, \pi(j))$ where vertices i and j are in the upper row and i_1 and j_1 are in the lower row. If $i < j, i_1 < j_1$ then edges are as in the Brauer diagram. If $i < j$ and $j_1 < i_1$, then we draw edges in two forms as shown below.




In form 1, we say (i, i_1) is the upper edge than (j, j_1) . In this case we may also say that (j, j_1) is lower than (i, i_1) . In form 2, we say that (j, j_1) is the upper edge than (i, i_1) . In this case we may also say that (i, i_1) is lower than (j, j_1) . The above graph is called Knot graph of order n with respect to π .


Example 1: Knot graph of order 2;



Example 2: knot graph of order 3:



$K(\pi) = \{(e_1, e_3), (e_2, e_3)\}$ and $|K(\pi)| = 2$ Hence number of Knot mappings of π is $2^{|K(\pi)|} = 2^2 = 4$ The four Knot mappings of π is described below:



Definition 2.2. $\pi \in S_n, i < j, \pi(i) > \pi(j)$ we say there is a crossing and the edges are $(i, \pi(i))$ and $(j, \pi(j))$


Remark 2.3. Given $\pi \in S_n$ there are $2^{|K(\pi)|}$ Knot graphs which is equal to $\{G_i(\pi)\}$ where $i=1$ to are $2^{|K(\pi)|}$

Definition 2.4. If $G_i(\pi)$ is a Knot graph with respect to π , then π is called underlying graph of $G_i(\pi)$

Notation 2.5. $S_n(k)$ denote the collection of all Knot graphs

Remark 2.6. $S_1(k) = S_1$.


Remark 2.7. $S_2(k) \neq S_2$, For



Remark 2.8. $S_n \neq S_n(k) \forall n \geq 2$

Notation 2.9. A Knot graph with π is denoted by $G_i(\pi), i=1, 2, \dots, 2^{|K(\pi)|}$ Examples of knot graphs:

When $n = 4$




Notation 2.10: we denote a Knot graph of order n as $G_n(\pi)$

Theorem 2.11. Every generator of Knot symmetric algebra FT_n can be represented by a unique Knot graph of order n and every Knot graph of order n can be represented by a generator of Knot symmetric algebra FT_n .

That is there is a one to one correspondence between the generators of Knot symmetric algebra FT_n and the Knot Graph of order n

Proof: let (π, f_π) be an generator of FT_n . Now $\pi \in S_n$ the vertices of π are represented in two rows such that each row contains n vertices. Let $e_i = (i, \pi(i))$ and $e_j = (j, \pi(j))$ be two edges. If $i < j$ and $\sigma(i) > \sigma(j)$ we draw in such a way that e_i is upper than e_j if $f_\pi(e_i, e_j) = 1$. The diagram is refer in below **Fig 1:** Next we refer the below **Fig 2** as follows: we draw in such a way that e_j is lower than e_i with respect to f_π if $f_\pi(e_i, e_j) = -1$. The diagram of Fig 1 and Fig 2 is drawn as follows:



Thus we get a Knot graph of order n corresponding to (π, f_π) . Now we will prove that every Knot graph of order n represent a generator.

Let G_n be a Knot graph of order n . Now $\pi(G_n) \in S_n$ we denote π instead of $\pi(S_n)$. Define $f_\pi: S(\pi) \rightarrow \{-1, 0, 1\}$ as

$$f_\pi(e_i, e_j) = \begin{cases} 0 & \text{if } \pi(i) < \pi(j) \\ 1 & \text{if } e_i \text{ is upper than } e_j \\ -1 & \text{if } e_i \text{ is lower than } e_j \end{cases}$$

It is obvious that the graph represented by (π, f_π) is G_n

Example. when $n=4$,


$$\text{let } \pi = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 2 & 1 & 3 \end{pmatrix} \text{ and } E(\pi) = \{e_1=(1,4), e_2=(1,2), e_3=(3,1), e_4=(4,3)\}$$

$$S(\pi) = \{(e_1, e_2), (e_1, e_3), (e_1, e_4), (e_2, e_3), (e_2, e_4), (e_3, e_4)\}$$

Let f_π be defined as follows:

$$f_\pi(e_1, e_2) = 1, f_\pi(e_1, e_3) = 1, f_\pi(e_1, e_4) = 1, f_\pi(e_2, e_3) = 1, f_\pi(e_2, e_4) = 0, f_\pi(e_3, e_4) = 0.$$

Now the Knot Graph represented by (π, f_π) is shown below: In this example the edge e_1 is upper than e_2 with respect to f_π



REFERENCES

- [PK] M. Parvathi and M. Kamaraj signed Brauer's Algebra ,Communications in Algebra, 26(3),839-855(1998).
- [Br] R. Brauer, algebras which are connected with the semisimple continuous graphs, Ann of Math,38(1937),854-872.
- [W] H. Wenzl on the structure of Brauer's centralizer algebras, Ann of math (128) (1988), 173-193.
- [PS] M. Parvathi and C. Selvarajsigned Brauer's algebras as centralizer algebras, communication in algebra 2 7(12) 5985 -5998(1999).
- [KM] M. Kamarajand R. Mangayarkarasi, Knot Symmetric Algebras, Research journal of pure algebra-1(6) (2011), 141-151.
- [RBA] The Rook Brauer Algebra Elise G.delmas "The Rook" (2012). Honors project paper 26, Macalester College, edelmas@macalester.edu.

Source of support: Nil, Conflict of interest: None Declared