



ON NORMAL ALMOST DISTRIBUTIVE LATTICES

S. RAMESH\*

Department of Mathematics, GITAM University, Visakhapatnam-530045, INDIA

U. M. SWAMY

Department of Mathematics, University of Gondar, Ethiopia

(Received on: 11-02-13; Revised & Accepted on: 09-03-13)

ABSTRACT

We examine the normality of subADLs of a normal ADL<sub>0</sub>. We obtain necessary and sufficient conditions for an ADL<sub>0</sub> to become normal (relatively normal) in terms of filter congruences and prime filter congruences.

Mathematics Subject Classifications (2000): 06D99.

Keywords: Relatively complemented ADL<sub>0</sub>, Normal ADL<sub>0</sub>, Prime ideals, Minimal prime ideals, Congruences.

INTRODUCTION

The concept of almost distributive lattice with zero (ADL<sub>0</sub>) was introduced by Swamy and Rao [5] in 1980 as a common abstract of ring theoretic and lattice theoretic generalization of Boolean algebras. It is an algebraic structure of type (2, 2, 0) which satisfies all the conditions of a distributive lattice except the commutativity of  $\vee$ ,  $\wedge$  and the right distributivity of  $\vee$  over  $\wedge$ . Rao and Ravi Kumar [3] introduced the concept of the normality of an ADL<sub>0</sub> in 2008. They obtained several equivalent conditions for an ADL<sub>0</sub> to become a normal almost distributive lattice in terms of prime ideals, minimal prime ideal and annihilator ideals.

In this paper, we observe that a subADL<sub>0</sub> of a normal ADL<sub>0</sub> need not be normal. We obtain a necessary and sufficient condition for a subADL<sub>0</sub> of a normal ADL<sub>0</sub> to become a normal subADL<sub>0</sub>. We study the normality and relative normality of an ADL<sub>0</sub> in terms of filter congruences and prime filter congruences

1. PRELIMINARIES

First we recall the definitions and certain necessary properties of almost distributive lattices with zero from [5].

**Definition 1.1:** [5] An Algebra  $(L, \vee, \wedge, 0)$  of type (2, 2, 0) is called an almost distributive lattice with 0 (ADL<sub>0</sub>) if, it satisfies the following conditions.

- (i)  $0 \wedge a = 0$
  - (ii)  $a \vee 0 = a$
  - (iii)  $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$
  - (iv)  $(a \vee b) \wedge c = (a \wedge c) \vee (a \wedge b)$
  - (v)  $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$
  - (vi)  $(a \vee b) \wedge b = b$
- for all  $a, b, c \in L$ .

**Example 1.2:** [5] Let X be a non empty set. Fix  $x_0 \in X$ . For any  $x, y \in X$ , define

$$x \vee y = \begin{cases} x & \text{if } x \neq x_0 \\ y & \text{if } x = x_0 \end{cases} \quad \text{and} \quad x \wedge y = \begin{cases} y & \text{if } x \neq x_0 \\ x & \text{if } x = x_0 \end{cases}$$

Then  $(X, \vee, \wedge, x_0)$  is an almost distributive lattice with  $x_0$  as its “0”

Form here onwards L means almost distributive lattice with ‘0’ as its zero element. For any  $a, b \in L$ , we say that a is less than or equal to b (that is,  $a \leq b$ ) if  $a \wedge b = a$  or equivalently  $a \vee b = b$ . It can be easily verified that ‘ $\leq$ ’ is a partial ordering on L. An element m of L is said to be maximal if  $m \wedge x = x$  for all  $x \in L$ .

\*Corresponding author: S. RAMESH\*

Department of Mathematics, GITAM University, Visakhapatnam-530045, INDIA

**Definition 1.3:** [5] A non empty sub set I of L is said to be an ideal (filter) of L, if it satisfies the following conditions;

- (i) For all  $a, b \in L, a \vee b \in L (a \wedge b \in L)$
- (ii) For all  $a \in L, x \in I, a \wedge x \in I (x \vee a \in L)$

A proper ideal (filter) P of L is said to be a prime ideal (filter) if, for any  $a, b \in L, a \wedge b \in P (a \vee b \in P)$  implies  $a \in P$  or  $b \in P$ . It can be routinely verified that a proper sub set P of L is prime ideal of L if and only if L-P is a prime filter of L.

**Definition 1.4:** [5] A prime ideal P of L is said to be a minimal prime ideal of L, if there is no prime ideal which is properly contained in P. Similarly, a proper filter P of L is said to be maximal filter of L if there is no proper filter containing P. It can be easily verified that a proper ideal P of L is a minimal prime ideal of L if and only if L-P is a maximal filter of L. Since every proper filter contained in a maximal filter, every non-zero element is contained in a maximal filter. Therefore for any non-zero element x of L, there is a minimal prime ideal P of L such that  $x \notin P$ . Hence we have the following.

**Theorem 1.5:** [6] The intersection of all minimal prime ideals of L is equal to  $\{0\}$ .

For any  $x \in L$ , the set  $(x)^* = \{y \in L | x \wedge y = 0\}$  is an ideal of L.

**Theorem 1.6:**[4] A prime ideal P of L is minimal if and only if, for each  $x \in P$ , there exists  $y \notin P$  such that  $x \wedge y = 0$ . (That is,  $(L-P) \cap (x)^*$  is non-empty.)

**Definition 1.7:** [5] L is said to be a relatively complemented if, given  $a, b \in L$ , there exists  $x \in L$  such that  $a \wedge x = 0$  and  $a \vee x = a \vee b$ .

**Theorem 1.8:** [5] L is relatively complemented if and only if every prime ideal is minimal.

## 2. ON NORMAL ALMOST DISTRIBUTIVE LATTICES

An almost distributive lattice with zero is called normal [3] if, every prime ideal contains a unique minimal prime ideal or equivalently, for  $x, y \in L, x \wedge y = 0$  implies  $(x)^* \vee (y)^* = L$ . A non empty subset S of L is said to be a subADL<sub>0</sub>, if it contains "0" and closed under operations  $\vee$  and  $\wedge$ . An almost distributive lattice with zero L is said to be a dense if  $\{0\}$  is a prime ideal of L.

We observe that a subADL<sub>0</sub> of a normal ADL<sub>0</sub> need not be normal. For, consider the following example.

**Example 2.1:** Let  $X = \{a, b, c\}$ . Let  $L = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ . Then L is a subADL<sub>0</sub> of P(X) with respect to the set inclusion and L is not normal.

In this context, we obtain the following.

**Theorem 2.2:** The following are equivalent for any ADL<sub>0</sub> L.

- (i) Every subADL<sub>0</sub> of L is normal
- (ii) For  $x, y \in L - \{0\}, x \wedge y = 0$ , implies  $x \vee y$  is maximal
- (iii) L is dense ADL<sub>0</sub> or L relatively complemented and every chain in L has at most three elements.

**Proof:** (i)  $\Rightarrow$  (ii): Suppose that every subADL<sub>0</sub> of L is normal. Let  $x, y \in L - \{0\}$  such that  $x \wedge y = 0$ . Suppose there is  $z \in L$  such that  $x \vee y < z$ . Then  $L_1 = \{0, x, y, x \vee y, z\}$  is a subADL<sub>0</sub> of L which is not normal. This is a contradiction to our assumption. Therefore  $x \vee y$  is maximal.

(ii)  $\Rightarrow$  (iii): Assume (ii). Suppose L is not dense ADL<sub>0</sub>. Then  $\{0\}$  is not a prime ideal of L. Let P be a prime ideal of L. Suppose P is not a minimal prime ideal of L. Then there is a minimal prime ideal M ( $\neq \{0\}$ ) of L such that  $M \subset P$ . Choose  $x \in M$  such that  $x \neq 0$ . Now, for any  $y \in L, 0 \neq y \in (x)^* \cap P \Rightarrow x \wedge y = 0$  and  $y \in P \Rightarrow x \vee y$  is maximal (by our assumption) and  $x \vee y \in P$  (since  $y \in P$  and  $x \in M \subset P$ ). This is a contradiction to the minimality of M (see Theorem 1.6). Therefore every prime ideal is minimal and hence L is relatively complemented (see Theorem 1.8). Let  $x, y, z \in L - \{0\}$  such that  $0 < x < y < z$ . Since L is relatively complemented, there exists  $t \in L$  such that  $x \wedge t = 0$  and  $x \vee t = x \vee y = y$ . By our assumption, we get that  $y = x \vee t$  is maximal. This is a contradiction. Therefore every chain has at most three elements.

(iii)  $\Rightarrow$  (i): Assume (iii). Let  $L_1$  be a subADL<sub>0</sub> of L. Let  $x, y \in L_1$  such that  $x \wedge y = 0$ . Suppose  $x \neq 0$  and  $y \neq 0$ .

Then  $0 < x < x \vee y$ . Otherwise  $x = x \vee y$  implies  $y = (x \vee y) \wedge y = x \wedge y = 0$ , which is a contradiction. By our assumption,  $x \vee y$  is maximal in  $L_1$  and  $x \vee y \in (y)^*_{L_1} \vee (x)^*_{L_1} = L_1$ . Thus  $L_1$  is normal.

Given a filter  $F$  of  $L$ , define  $\varphi_F = \{(a, b) \in L \times L \mid x \wedge a = x \wedge b \text{ for some } x \in F\}$ . Then  $\varphi_F$  is a congruence on  $L$ .

The following is a routine verification.

**Theorem 2.3:** For any filter  $F$  of  $L$  and  $x \in L$ , we have

- (i)  $\frac{x}{\varphi_F}$  is a maximal in  $\frac{L}{\varphi_F}$  if and only if  $x \in F$
- (ii)  $\frac{x}{\varphi_F} = \frac{0}{\varphi_F}$  if and only if  $(x)^* \cap F \neq \phi$

**Proof:** (i) Suppose  $\frac{x}{\varphi_F}$  is a maximal element in  $\frac{L}{\varphi_F}$ . Since  $F \neq \phi$ , we can choose  $y \in F$ . Then  $\frac{x}{\varphi_F} \wedge \frac{y}{\varphi_F} = \frac{y}{\varphi_F}$ .

That is  $(x \wedge y, y) \in \varphi_F$ . Therefore there exists  $a \in F$  such that  $a \wedge x \wedge y = a \wedge y \in F$  and hence  $x \in F$ . On the other hand, suppose  $x \in F$ . Let  $y \in L$ . The  $x \wedge x \wedge y = x \wedge y$ .

Therefore  $(x \wedge y, y) \in \varphi_F$  and hence  $\frac{x}{\varphi_F} \wedge \frac{y}{\varphi_F} = \frac{y}{\varphi_F}$ . Thus  $\frac{x}{\varphi_F}$  is a maximal element in  $\frac{L}{\varphi_F}$ .

- (iii) Suppose  $\frac{x}{\varphi_F} = \frac{0}{\varphi_F}$ . Then there exists  $y \in F$  such that  $y \wedge x = y \wedge 0 = 0$ . Therefore  $y \in (x)^* \cap F$  and

hence  $(x)^* \cap F \neq \phi$ . On the other hand, suppose  $(x)^* \cap F \neq \phi$ . Choose  $y \in F$  such that  $x \wedge y = 0$ . Then  $y \wedge x = y \wedge 0$ . Therefore  $(x, 0) \in \varphi_F$  and hence  $\frac{x}{\varphi_F} = \frac{0}{\varphi_F}$ .

In the following, for any filter  $F$  of  $L$ , we obtain a one-to-one correspondence between prime ideals of  $L$  disjoint with  $F$  and prime ideal of  $\frac{L}{\varphi_F}$ .

**Theorem 2.4:** Let  $F$  be a filter of  $L$ . For any prime ideals  $P$  of  $L$  with  $P \cap F = \phi$ , let  $\bar{P} = \left\{ \frac{x}{\varphi_F} \in \frac{L}{\varphi_F} \mid x \in P \right\}$ . Then

$\bar{P}$  is a prime ideal of  $\frac{L}{\varphi_F}$ . Also,  $P \mapsto \bar{P}$  is an order isomorphism (with respect to the inclusion ordering) of the set of prime ideals of  $L$  disjoint with  $F$  onto the set of prime ideals of  $\frac{L}{\varphi_F}$ . This map induces a one to one correspondence between minimal prime ideals of  $L$  disjoint with  $F$  and minimal prime ideals of  $\frac{L}{\varphi_F}$ .

**Proof:** Let  $P$  be a prime ideal of  $L$  such that  $P \cap F = \phi$ . Then, it is easily to verify that  $\bar{P} = \left\{ \frac{x}{\varphi_F} \mid x \in P \right\}$  is an ideal

of  $\frac{L}{\varphi_F}$ . Also, we observe that, for any  $a \in L$ ,  $\frac{a}{\varphi_F} \in \bar{P} \Leftrightarrow a \in P$ . Now, for any  $x, y \in L$ ,


$$\frac{x}{\varphi_F} \wedge \frac{y}{\varphi_F} \in \bar{P} \Rightarrow \frac{(x \wedge y)}{\varphi_F} \in \bar{P} \Rightarrow x \wedge y \in P \Rightarrow x \in P \text{ or } y \in P \text{ (since } P \text{ is prime)} \Rightarrow \frac{x}{\varphi_F} \in \bar{P} \text{ or } \frac{y}{\varphi_F} \in \bar{P}.$$

Therefore  $\bar{P}$  is a prime ideal of  $\frac{L}{\varphi_F}$ . Now, for any prime ideals  $P$  and  $Q$  of  $L$  with  $P \cap F = \phi = Q \cap F$ ,

$$P \subseteq Q \Leftrightarrow \bar{P} \subseteq \bar{Q}. \text{ Let } R \text{ be a prime ideal of } \frac{L}{\varphi_F}. \text{ Put } P = \left\{ x \in L \mid \frac{x}{\varphi_F} \in R \right\}. \text{ Then } P \text{ is a prime ideal of } L$$

disjoint with  $F$  and  $\overline{P} = R$ . Thus the map  $P \mapsto \overline{P}$  is an order isomorphism from the set of prime ideals of  $L$  disjoint with  $F$  onto the set of prime ideals of  $\frac{L}{\varphi_F}$ .

Any two ideals  $I$  and  $J$  are said to be co-maximal, if  $I \vee J = L$ . An  $ADL_0$   $L$  is relatively normal [3], if given  $x, y \in L$  with  $x \leq y$ , the interval  $[x, y] = \{z \in L \mid x \leq z \leq y\}$  is normal, or equivalently, any two incomparable prime ideals are co-maximal. In general every relatively normal  $ADL_0$  is normal but not conversely. For consider  $L = \{0, a, b, c, d, 1\}$ .



Then the prime ideals  $P_1 = \{0, a, b\}$ ,  $P_2 = \{0, a, c\}$  are incomparable and are not co-maximal.

The following is a consequence of the above.

**Theorem 2.5:**  $L$  is normal (relatively normal) if and only if, for any filter  $F$  of  $L$ ,  $\frac{L}{\varphi_F}$  is normal (relatively normal).

Note that any dense  $ADL_0$  is normal, because  $\{0\}$  is the unique prime ideal of  $L$ . Since the intersection of all minimal prime ideals is  $\{0\}$  in any  $ADL_0$  (see Theorem 1.5), it follows that an  $ADL_0$   $L$  is dense if and only if it has only one minimal prime ideal which is  $\{0\}$ . The following result is another characterization of the normality of an  $ADL_0$ .

**Theorem 2.6:**  $L$  is normal if and only if, for any prime filter  $P$  of  $L$ ,  $\frac{L}{\varphi_P}$  is a dense  $ADL_0$ .

**Proof:** Suppose that  $L$  is normal. Let  $P$  be a prime filter of  $L$  and  $I = L - P$ . Then  $I$  is a prime ideal of  $L$  and hence  $I$  contains a unique minimal prime ideal. Say  $M$ . Then  $M$  is the only minimal prime ideal of  $L$  which is disjoint with  $P$ .

Therefore, by Theorem 2.4,  $\frac{L}{\varphi_P}$  has a unique minimal prime ideal which implies that  $\left\{ \frac{0}{\varphi_P} \right\}$  is a prime ideal in  $\frac{L}{\varphi_P}$

and hence  $\frac{L}{\varphi_P}$  is dense. Conversely, suppose that  $\frac{L}{\varphi_P}$  is a dense  $ADL_0$ , for any prime filter  $P$  of  $L$ . That is, the zero

ideal in  $\frac{L}{\varphi_P}$  is prime. Let  $I$  be a prime ideal of  $L$ . Then  $P = L - I$  is a prime filter of  $L$ . By our assumption,  $\frac{L}{\varphi_P}$  has

only one minimal prime ideal namely,  $\left\{ \frac{0}{\varphi_P} \right\}$ . By Theorem 2.4, there is only one minimal prime ideal of  $L$  disjoint with  $P$  (or, equivalently, contained in  $I$ ). Thus  $L$  is normal.

Given a congruence  $\theta$  on  $L$ , we say that  $\frac{L}{\theta}$  is an almost chain, if for any  $x, y \in L$ ,  $\frac{x}{\theta} \wedge \frac{y}{\theta} = \frac{y}{\theta}$  or  $\frac{y}{\theta} \wedge \frac{x}{\theta} = \frac{x}{\theta}$ .

**Theorem 2.7:**  $L$  is relatively normal if and only if  $\frac{L}{\varphi_P}$  is an almost chain, for any prime filter  $P$  of  $L$ .

**Proof:** Suppose that  $L$  is relatively normal. Let  $P$  be a prime filter of  $L$ . Suppose that  $a$  and  $b$  are elements in  $L$  such that  $\frac{a}{\theta} \wedge \frac{b}{\theta} \neq \frac{b}{\theta}$  and  $\frac{b}{\theta} \wedge \frac{a}{\theta} \neq \frac{a}{\theta}$ . Then there exists a prime ideal  $R$  and  $S$  of  $\frac{L}{\varphi_P}$  such that

$$\frac{(a \wedge b)}{\varphi_P} \in R, \frac{b}{\varphi_P} \notin R, \frac{(b \wedge c)}{\varphi_P} \in S \text{ and } \frac{a}{\varphi_P} \notin S.$$

Put  $A = \left\{ x \in L \mid \frac{x}{\varphi_P} \in R \right\}$  and  $B = \left\{ x \in L \mid \frac{x}{\varphi_P} \in S \right\}$ . Then A and B are prime ideals of L which are disjoint from P and hence A and B are contained in the prime ideal  $L - P$ . This implies that A and B are not co-maximal. Also, we have

$$a \wedge b \in A, b \notin A \text{ \& } a \in A \text{ and } b \wedge a \in B, a \notin B \text{ \& } b \in B$$

So that A and B are incomparable. This is a contradiction. Thus  $\frac{L}{\varphi_P}$  is an almost chain.

Conversely, suppose that, for any prime filter P of L,  $\frac{L}{\varphi_P}$  is an almost chain. Let I and J be two incomparable prime ideals of L such that  $I \vee J \neq L$ . Since every proper ideal is contained in a prime ideal (by the Zorn's lemma), there exists a prime ideal K of L such that  $I \vee J \subseteq K$ . Put  $P = L - K$ . Then P is a prime filter of L. Choose  $x \in I - J$  and  $y \in J - I$ . Since  $\frac{L}{\varphi_P}$  is an almost chain, with out loss of generality we can suppose that  $\frac{x}{\varphi_P} \wedge \frac{y}{\varphi_P} = \frac{y}{\varphi_P}$ . Then  $(x \wedge y, y) \in \varphi_P$ . Therefore  $t \wedge x \wedge y = t \wedge y$  for some  $t \in P = L - K$ . Since  $x \in I$ , we get that  $t \wedge y \in I$ . Which is a contradiction, since  $t \notin I$  and  $y \notin I$ . Therefore any two incomparable prime ideal are co-maximal. Hence L is relatively normal.

#### REFERENCE

1. Birkhoff, G., *Lattice Theory*, American Mathematical Society, Colloquium Publications, 1967.
2. Cornish, W.H., *Normal lattices*, J. Austral Math. Soc., Vol.14 (1971), 200-215.
3. Rao, G.C., Ravi Kumar, S., *Normal almost distributive lattices*, Southeast Asian Bulletin of Mathematics, Vol.32 (2008), 831-841.
4. Rao, G.C., Ravi Kumar, S., *Minimal prime ideals in almost distributive lattices*, Int. J. Contemp. Math. Sciences, Vol.4(2009),475-484.
5. Swamy, U.M., Rao, G.C., *Almost distributive lattices* J. Austral math. Soc. Vol.31 (1981), 77-91.
6. Swamy, U.M., Ramesh, S., Shanthi Sundar Raj, Ch., *Prime ideal characterization of Stone almost distributive lattices*, Vol.3 (2010), 357-367.

**Source of support: Nil, Conflict of interest: None Declared**