



COMMON FIXED POINT THEOREMS IN FUZZY METRIC SPACE USING INTEGRAL TYPE

M. Ramana Reddy^{1*} and Prof. V. Dharmiah²

¹*Asst. Professor Of Mathematics, Sreenidhi Institute Of Science & Technology, Hyderabad – (A.P.), India.*

²*Professor of Mathematics, Osmania University University, Hyderabad – (A.P.), India.*

(Received on: 12-10-13; Revised & Accepted on: 22-10-13)

ABSTRACT

In this paper, the concept of common fixed point theorem in fuzzy metric space using integral type.

Key Words: contraction condition, fuzzy metric space, weakly compatible mapping, common fixed point

AMS subject classification: 47H10, 54H25.

1. INTRODUCTION

Zadseh in 1965 was introduced fuzzy set. This concept in topology and analysis many authors have developed the theory of fuzzy set and applications. Recently many authors have also studied the fixed point theory in fuzzy metric spaces ([11], [12][15], [22], [28], [29]. Grabiec[15] followed kramsoil and michalek[17] and obtained the banasch's fixed theorem using fuzzy sets. jungck[6] established common fixed point theorems for commuting maps generalizing the banach's fixed point theorems. sessa [25] defined a generalization of commutativity which is called weak commutativity. The study of common fixed point of mappings satisfying contractive type conditions has been a very active field of research during recent years. Several authors [5], [16], [17], [32] proved some fixed point theorems for various generalizations of contraction mapping in probabilistic and fuzzy metric metric space. In this paper, we prove some common fixed point theorems for mappings by using contracting condition of integral type for class of weakly compatible maps in non complete intuitionistic fuzzy metric spaces

2. PRELIMINARIES

Definition 2.1:[5].A binary operation $*$: $[0,1] \times [0,1] \rightarrow [0,1]$ is called a continuous t- norm if $([0,1],*)$ is an abelian topological monoid with the unit 1 such that $a*b \leq c*d$ whenever $a \leq c$ and $b \leq d$ for all $a,b,c,d \in [0,1]$. Examples of t- norms are $a*b = ab$ and $a*b = \min\{a, b\}$

Definition 2.2 ([21]): the 3 –tuples $(X, M,*)$ is called a fuzzy metric space if X is an arbitrary set * is a continuous t- norm, and M is fuzzy set in $X^2 \times [0,\infty]$ satisfying the following conditions for all $x, y, z \in X$ and $t, s > 0$.

1. $M(x, y, 0) = 0$
2. $M(x, y, t) = 1 \forall t > 0 \Leftrightarrow x = y$
3. $M(x, y, t) = M(y, x, t)$
4. $M(x, y, t) * M(y, z, s) \leq M(x, z, t+s)$ and
5. $M(x, y, *): [0, \infty) \rightarrow [0,1]$ is left continuous.
6. $\lim_{t \rightarrow \infty} M(x, y, t) = 1$

In what follow $(X, M,*)$ will denote a fuzzy metric space. Note that $M(x, y, t)$ can be thought as the degree of nearness between x and y with respect to t. We identify $x = y$ with $M(x, y, t) = 1$ for all t. 0 and $M(x, y, t) = 0$ with ∞ and we can find some topological properties and examples of fuzzy metric spaces in George and veeramani [3].

In the following example, we know that every metric induces a fuzzy metric.

$$M(x, y, t) = \frac{t}{t + d(x, y)}, \tag{A}$$

***Corresponding author: M. Ramana Reddy^{1*}**
¹Sreenethi Engineering College, Hyderabad, India.

Then $M(x, y, *)$ is a fuzzy metric space. We call the fuzzy metric M induced by the metric d the standard fuzzy metric. On the other hand note that there exists no metric on x satisfying (A)

Lemma 2.1([15]): for all $x, y \in X$, $M(x, y, *)$ is non decreasing.

Definition 2.3: ([15]) Let $M(x, y, *)$ is a fuzzy metric space:

- (i). A sequence $\{X_n\}$ in X is said to be convergent to a point $x, \in X$ (denoted by $\lim_{t \rightarrow \infty} X_n = x$) if for $t > 0$ and $\lim_{t \rightarrow \infty} M(x, y, t) = 1$
- (ii) A sequence $\{X_n\}$ in X is said to be Cauchy sequence if, for all $t > 0$ and $p > 0$, $\lim_{t \rightarrow \infty} M(x_{n+p}, x_n, t) = 1$
- (iii). A fuzzy metric space in which every Cauchy sequence is convergent is said to be complete.

Note 2.1: since $*$ is continuous, it follow from definition 2.2(4) that the limit of the sequence in fuzzy metric space is uniquely determined

Lemma 2.2 ([23] [33]): Let $\{y_n\}$ be a sequence in a fuzzy metric space $M(x, y, *)$ with the condition Definition 2.2(6) . If there exisats a number $k \in (0, 1)$ such that $M(y_{n+2}, y_{n+1}, kt) \geq M(y_{n+1}, y_n, t)$ for all $t > 0$ and $n = 1, 2, \dots$. Then $\{y_n\}$ is a Cauchy sequence in X .

Lemma 2.3 ([23][33]): if for all $x, y \in X$, $t > 0$ and for a number $k \in (0, 1)$

$$M(x, y, kt) \geq M(x, y, t)$$

Then $x = y$.

Definition 2.4 ([23]): Let A and B be mappings from a fuzzy metric space $M(x, y, *)$ into it self. The mapping A and B are said to be compatible if $\lim_{t \rightarrow \infty} M(ABx_n, BAx_n, t) = 1$ For all $t > 0$, whenever $\{X_n\}$ is a sequence in X such that $\lim_{t \rightarrow \infty} Ax_n = \lim_{t \rightarrow \infty} Bx_n = Z$ for some $z \in X$.

Definition 2.5 ([33]): Let A and B be mappings from a fuzzy metric space $M(x, y, *)$ into it self. The mapping A and B are said to be compatible of type (α) if for all $t > 0$

$$\text{if } \lim_{t \rightarrow \infty} M(ABx_n, BBx_n, t) = 1 \text{ and } \lim_{t \rightarrow \infty} M(BAx_n, AAx_n, t) = 1$$

For all $t > 0$, whenever $\{X_n\}$ is a sequence in X such that $\lim_{t \rightarrow \infty} Ax_n = \lim_{t \rightarrow \infty} Bx_n = Z$ for some $z \in X$.

Definition 2.6 ([34]): Let A and B be mappings from a fuzzy metric space $M(x, y, *)$ into it self . The mapping A and B are said to be compatible of type (β) if for all $t > 0$

$$\text{if } \lim_{t \rightarrow \infty} M(AAx_n, BBx_n, t) = 1 \text{ when ever For all } t > 0, \text{ whenever } \{X_n\} \text{ is a sequence in } X \text{ such that } \lim_{t \rightarrow \infty} Ax_n = \lim_{t \rightarrow \infty} Bx_n = Z \text{ for some } z \in X.$$

Definition 2.6 ([9]): two self mappings S and T are said to be weakly compatible if they commute at their coincidence points, i.e, if $Tu = Su$ for $u \in X$. $TSu = Stu$

Examples 2.2: Let $X [0, 2]$ with the metric d define by $d(x, y) = |x - y|$. For each $t \in (0, \infty)$

$$M(x, y, t) = \frac{t}{t + d(x, y)},$$

And define $M(x, y, 0) = 0$ $x, y \in X$

Clearly, $M(x, y, *)$ is a fuzzy metric space on X where $*$ is defined by $a * b = ab$ or $a * b = \min\{a, b\}$

$$AX = \begin{cases} x, & \text{if } x \in [0, \frac{1}{4}] \\ \frac{1}{4}, & \text{if } x \geq \frac{1}{4} \end{cases} \quad Bx = \frac{x}{1+x}$$

For all $x \in [0, 2]$. Consider the sequence $\{X_n = 1/3 + 1/n: n \geq 1\}$ in X . Then

$$\lim_{n \rightarrow \infty} Ax_n = \frac{1}{4}, \lim_{n \rightarrow \infty} Bx_n = \frac{1}{4}$$

$$\lim_{t \rightarrow \infty} M(ABx_n, BAx_n, t) = \frac{1}{t + \frac{1}{4} + \frac{1}{5}} \neq 1$$

and

$$\lim_{t \rightarrow \infty} M(BAx_n, AAx_n, t) = \frac{1}{t + \frac{t-1}{5}} \neq 1. \text{ Thus A and B are}$$

non compatible, but A and B are commuting at their. Coincidencce point, That is weakly compatible at $x = 0$. also Thus A and B are not compatible of type (β) . in view of this example, we observe weakly compatible maps need not be compatible. or compatible of type (α) and type (β)

3. MAIN THEOREM

Theorem 3.1: Let $(X, M, *)$ be a fuzzy metric space with continuous t – norm $*$ defined by $t*t$ for all $t \in [0, 1]$. Let A, B, S, T, P and Q be mappings from X into itself such that

(a) $P(X) \subset ST(X), Q(X) \subset AB(X)$

(b) there exists a constant $k \in (0, 1)$ such that

$$\int_0^{M(Px, Qy, qt)} \varphi(t) dt \geq \int_0^{m(x, y, t)} \varphi(t) dt$$

Where $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a lebesque – integrable mapping which is summable, non negative, and such that

$$\int_0^\varepsilon \varphi(t) dt > 0 \text{ for each } \varepsilon > 0$$

Where $m(x, y, t) = \min\{M(ABx, Sty, t), M(Px, ABX, t), M(Qy, Sty, t), M(Px, ABX, t), M(Px, Sty, t)\}$ For all $x, y \in X$ and $t > 0$

(c) $P(x), Q(x), AB(x), ST(x)$ is complete subspace of X, then

- (i). P and AB have a coincidence point, and
- (ii). Q and St have a coincidence point.

(d) $AB=BA, ST=TS, PB=BP, QT=TQ,$

(e). the pair $\{P, AB\}$ is weakly compatible ,

Then A, B, S, T, P and Q have a unique common fixed point in X.

Proof: let $x_0 \in X$, from (a) there exist $x_1, x_2 \in X$ such that

$$Px_0 = STx_1 \text{ And } Qx_1 = ABx_2$$

Inductively, we can construct sequence $\{X_n\}$ and $\{y_n\}$ in X such that

$$Px_{2n-2} = STx_1 = y_{2n-1} \text{ and}$$

$$Qx_{2n-1} = ABx_{2n} = y_{2n} \text{ for } n=1, 2, 3, \dots$$

Step - 1: put $x=x_{2n}$ and $y=y_{2n+1}$ in (e), we get

$$\begin{aligned} \int_0^{M(Px_{2n}, Qx_{2n+1}, qt)} \varphi(t) dt &\geq \int_0^{M(ABx_{2n}, STx_{2n+1}, t)M(Px_{2n}, ABx_{2n}, t)M(Qx_{2n+1}, STx_{2n+1}, t)M(Px_{2n}, STx_{2n+1}, t)} \varphi(t) dt \\ &= \int_0^{M(y_{2n}, y_{2n+1}, t)M(y_{2n+1}, y_{2n}, t)M(y_{2n+2}, y_{2n+1}, t)M(y_{2n+1}, y_{2n+1}, t)} \varphi(t) dt \\ &\geq \int_0^{M(y_{2n}, y_{2n+1}, t)M(y_{2n+1}, y_{2n+2}, t)} \varphi(t) dt \end{aligned}$$

Form lemma 2, 1 and 2, 2, we have

Similarly, we have

$$\int_0^{M(y_{2n+2}, y_{2n+3}, qt)} \varphi(t) dt \geq \int_0^{M(y_{2n+1}, y_{2n+2}, t)} \varphi(t) dt$$

Thus, we have

$$\int_0^{M(y_{n+1}, y_{n+3}, qt)} \varphi(t) dt \geq \int_0^{M(y_n, y_{n+1}, t)} \varphi(t) dt \text{ for } n=1, 2, \dots$$

$$\begin{aligned} \int_0^{M(y_n, y_{n+1}, t)} \varphi(t) dt &\geq \int_0^{M(y_n, y_{n+1}, \frac{t}{q})} \varphi(t) dt \\ &\geq \int_0^{M(y_{n-2}, y_{n-1}, \frac{t}{q^2})} \varphi(t) dt \\ &\dots\dots\dots \\ &\geq \int_0^{M(y_1, y_2, \frac{t}{q^n})} \varphi(t) dt \rightarrow 1 \text{ as } n \rightarrow \infty \end{aligned}$$

And hence $\int_0^{M(y_n, y_{n+1}, t)} \varphi(t) dt \rightarrow 1$ as $n \rightarrow \infty$ for any $t > 0$.

For each $\varepsilon > 0$ and $t > 0$, we choose $n_0 \in \mathbb{N}$ such that $\int_0^{M(y_n, y_{n+1}, t)} \varphi(t) dt > 1 - \varepsilon$ for all $n > n_0$

For $m, n \in \mathbb{N}$, we suppose $m \geq n$. Then we have

$$\begin{aligned} \int_0^{M(y_n, y_m, t)} \varphi(t) dt &\geq \int_0^{M(y_n, y_{n+1}, \frac{t}{m-n}) M(y_{n+1}, y_{n+2}, \frac{t}{m-n}) \dots M(y_{m-1}, y_m, \frac{t}{m-n})} \varphi(t) dt \\ &\geq (1 - \varepsilon) * (1 - \varepsilon) * \dots * (1 - \varepsilon) (m - n) \text{ times} \\ &\geq (1 - \varepsilon) \end{aligned}$$

And hence $\{y_n\}$ is a Cauchy sequence in X

Since $(X, M, *)$ is complete, $\{y_n\}$ converges to some point $z \in X$. Also its subsequence converges to the same point

i. e. $z \in X$

i.e. $\{Qx_{2n+1}\} \rightarrow Z$ and $\{STx_{2n+1}\} \rightarrow Z$ (1)

$\{Px_{2n}\} \rightarrow Z$ and $\{ABx_{2n}\} \rightarrow Z$ (2)

Case - I: suppose AB is continuous, we have

$(AB)^2 x_{2n} \rightarrow ABZ$ and

$ABx_{2n} \rightarrow ABZ$

As (P, AB) is compatible pair, we have $PABx_{2n} \rightarrow ABZ$

Step -2: put $x = ABx_{2n}$ and $y = x_{2n+1}$ in (e), we get

$$\int_0^{M(PABx_{2n}, Qx_{2n+1}, qt)} \varphi(t) dt \geq \int_0^{M(ABABx_{2n}, STx_{2n+1}, t) M(PABx_{2n}, ABABx_{2n}, t) M(Qx_{2n+1}, STx_{2n+1}, t) M(PABx_{2n}, STx_{2n+1}, t)} \varphi(t) dt$$

Taking $n \rightarrow \infty$, we get

$$\int_0^{M(ABZ, Z, qt)} \varphi(t) dt \geq \int_0^{M(ABZ, Z, t)M(ABZ, ABZ, t)Mz, z, t)M(ABZ, z, t)} \varphi(t) dt$$

$$\int_0^{M(ABZ, Z, qt)} \varphi(t) dt \geq \int_0^{M(ABZ, Z, t), M(ABZ, Z, t)} \varphi(t) dt$$

$$\int_0^{M(ABZ, Z, qt)} \varphi(t) dt \geq \int_0^{M(ABZ, Z, t)} \varphi(t) dt$$

Therefore, by using lemma 2.2., we get $ABz=z$

Step - 3: put $x=z$ and $y = x_{2n+1}$ In (b), we have

$$\int_0^{M(z, z, t)M(Pz, z, t)} \varphi(t) dt \geq \int_0^{M(z, z, t)M(Pz, z, t)M(z, z, t)M(Pz, z, t)} \varphi(t) dt \geq \int_0^{M(Pz, z, t)M(Pz, z, t)} \varphi(t) dt$$

$$\int_0^{M(Pz, z, t)} \varphi(t) dt \geq \int_0^{M(Pz, z, t)} \varphi(t) dt$$

Therefore by using lemma 2.2., we get $Pz=z$, therrefre $ABz=Pz=z$

Step - 4: putting $x=Bz$ and $y = x_{2n+1}$ in condition (e) we get

$$\int_0^{M(PBz, Qx_{2n+1}, qt)} \varphi(t) dt \geq \int_0^{M(ABBz, STx_{2n+1}, t)M(Pz, ABBz, t)M(Qx_{2n+1}, STx_{2n+1}, t)M(PBz, STx_{2n+1}, t)} \varphi(t) dt$$

As $BP=PB$, $AB=BA$, so we have $P(Bz)=B(Pz)=Bz$ and

$$(AB)(Bz) = (BA)(Bz) = B(ABZ) = Bz$$

Taking $n \rightarrow \infty$ and using (1), we get

$$\int_0^{M(Bz, z, qt)} \varphi(t) dt \geq \int_0^{M(Bz, z, t)M(Bz, Bz, t)M(z, z, t)M(Bz, z, t)} \varphi(t) dt \geq \int_0^{M(Bz, z, t)M(Bz, z, t)} \varphi(t) dt$$

$$\int_0^{M(Bz, z, qt)} \varphi(t) dt \geq \int_0^{M(Bz, z, qt)} \varphi(t) dt$$

Therefore, by using lemma 2.2, we get

$$Bz = z$$

And also we have $ABz=z$

$$\Rightarrow Az=z, \text{ Therefore } Az=Bz=Pz=z$$

(4)

Step - 5: As $P(X) \subseteq ST(X)$, there exists $u \in X$ such that

$Z=PZ=STU$. Putting $x=x_{2n}$ and $y=u$ in (e) we get

$$\int_0^{M(Px_{2n}, Qu, qt)} \varphi(t) dt \geq \int_0^{M(ABx_{2n}, STu, t)M(Px_{2n}, ABx_{2n}, t)M(Qu, STu, t)M(Px_{2n}, STu, t)} \varphi(t) dt$$

Taking $n \rightarrow \infty$ and using (1) and (2), we get

$$\int_0^{M(z,Qu,qt)} \varphi(t) dt \geq \int_0^{M(z,z,t)M(z,z,t)M(Qu,z,t)M(z,z,t)} \varphi(t) dt$$

$$\geq \int_0^{M(Qu,z,t)} \varphi(t) dt$$

$$\int_0^{M(z,Qu,qt)} \varphi(t) dt \geq \int_0^{M(z,Qu,t)} \varphi(t) dt$$

Therefore, by using lemma 2.2. we get $Qu=z$ hence $Stu=z=Qu$

Thus $Qz=STz$,

Step - 6: putting $x=x_{2n}$ and $y=z$ in (e)

$$\int_0^{M(Px_{2n},Qu,qt)} \varphi(t) dt \geq \int_0^{M(ABx_{2n},STu,t)M(Px_{2n},ABx_{2n},t)M(Qu,STu,t)M(Px_{2n},STu,t)} \varphi(t) dt$$

Taking $n \rightarrow \infty$ and using (2) and step 5, we get

$$\int_0^{M(z,Qz,qt)} \varphi(t) dt \geq \int_0^{M(z,z,t)M(z,z,t)M(Qz,Qz,t)M(z,qz,t)} \varphi(t) dt$$

$$\geq \int_0^{M(z,Qz,qt)M(z,Qz,qt)} \varphi(t) dt$$

$$\int_0^{M(z,Qz,qt)} \varphi(t) dt \geq \int_0^{M(z,Qz,t)} \varphi(t) dt$$

Therefore, by using lemma 2.2, we get $Qz=z$

Step - 7: putting $x=x_{2n}$ and $y= Tz$ in (e), we get

$$\int_0^{M(Px_{2n},QTz,qt)} \varphi(t) dt \geq \int_0^{M(ABx_{2n},STTz,t)M(Px_{2n},ABx_{2n},t)M(QTz,STTz,t)M(Px_{2n},STTz,t)} \varphi(t) dt$$

As $QT=TQ$ and $ST=Ts$, we have $QTz=TQz =Tz$ and $ST(Tz)=T(STz)=TQz=Tz$

Taking $n \rightarrow \infty$, we get

$$\int_0^{M(Pz,,Tz,qt)} \varphi(t) dt \geq \int_0^{M(z,Tz,t)M(z,z,t)M(Tz,Tz,t)M(z,Tz,t)} \varphi(t) dt$$

$$\geq \int_0^{M(z,,Tz,t)M(z,,Tz,t)} \varphi(t) dt$$

$$\int_0^{M(z,,Tz,qt)} \varphi(t) dt \geq \int_0^{M(z,,Tz,t)} \varphi(t) dt$$

Therefore, by using lemma 2.2. we get $Tz=z$ now $STz=Tz=z$ implies $Sz=z$ hence $Sz=Tz=Qz=Z$ (5)

Combining (4) and (5) we get

$$Az = Bz = Pz = Qz = Tz = Sz = z$$

Hence, z is the common fixed point of A, B, S, T, P and Q

Case - II: suppose P is continuous

As P is continuous

$$P^2 x_{2n} \rightarrow Pz \text{ and } P(AB) x_{2n} \rightarrow Pz$$

As (P, AB) is compatible pair. (AB) P x_{2n} → Pz

Step - 8: putting x= Px_{2n} and y=x_{2n+1} in condition (e), we have

$$\int_0^{M(P Px_{2n}, Q x_{2n+1}, qt)} \varphi(t) dt \geq \int_0^{M(AB Px_{2n}, ST x_{2n+1}, t) M(P Px_{2n}, AB Px_{2n}, t) M(Q x_{2n+1}, ST x_{2n+1}, t) M(P Px_{2n}, ST x_{2n+1}, t)} \varphi(t) dt$$

Taking n → ∞, we get

$$\begin{aligned} \int_0^{M(Pz, z, qt)} \varphi(t) dt &\geq \int_0^{M(Pz, z, t) M(Pz, Pz, t) M(z, z, t) M(Pz, z, t)} \varphi(t) dt \\ &\geq \int_0^{M(Pz, z, t) M(Pz, z, t)} \varphi(t) dt \end{aligned}$$

$$\int_0^{M(Pz, z, qt)} \varphi(t) dt \geq \int_0^{M(Pz, z, t)} \varphi(t) dt$$

Therefore by using lemma 2.2, we have Pz=z further, using step 5, 6, 7 we get Qz=STz=Sz=Tz=z

Step - 9: Q(X) ⊆ AB(X), there exists w belongs to X such that

$$Z=Qz=ABw$$

Put x=w and y=x_{2n+1} in (e), we have

$$\int_0^{M(Pw, Q x_{2n+1}, qt)} \varphi(t) dt \geq \int_0^{M(ABw, ST x_{2n+1}, t) M(Pw, ABw, t) M(Q x_{2n+1}, ST x_{2n+1}, t) M(Pw, ST x_{2n+1}, t)} \varphi(t) dt$$

Taking n → ∞, we get

$$\begin{aligned} \int_0^{M(Pw, z, qt)} \varphi(t) dt &\geq \int_0^{M(z, z, t) M(Pw, z, t) M(z, z, t) M(Pw, z, t)} \varphi(t) dt \\ &\geq \int_0^{M(Pw, z, t) M(Pw, z, t)} \varphi(t) dt \end{aligned}$$

$$\int_0^{M(Pw, z, qt)} \varphi(t) dt \geq \int_0^{M(Pw, z, t)} \varphi(t) dt$$

Therefore, buy using ABw=Pw=z

As (P, AB) is compatible pair. We have Pz=ABz

Also, from step4, we get Bz=z.

Thus Az = Bz =Pz z and we see that z is the common fixed point in this case also.

Uniqueness: Let u be another common fixed point of A,B,S,T,P and Q put x=z and y=uy in (e) we get

$$\int_0^{M(Pz, Qu, qt)} \varphi(t) dt \geq \int_0^{M(ABzSTu, t) M(Pz, ABz, t) M(Qu, STu, t) M(Pz, STu, t)} \varphi(t) dt$$

Taking $n \rightarrow \infty$, we get

$$\int_0^{M(z,u,qt)} \varphi(t) dt \geq \int_0^{M(z,u,t)M(z,z,t)M(u,u,t)M(z,u,t)} \varphi(t) dt$$

$$\geq \int_0^{M(z,u,t)M(z,u,t)} \varphi(t) dt$$

$$\int_0^{M(z,u,qt)} \varphi(t) dt \geq \int_0^{M(z,u,t)} \varphi(t) dt$$

Therefore by using lemma (2.2), we get $z=u$

Therefore z is the unique common fixed point of self maps A, B, S, T, P and Q .

Corollary 3.1: Let $(X, M, *)$ be a fuzzy metric space with continuous t – norm $*$ defined by $t*t$ for all $t \in [0,1]$. Let A, S, P and Q be mappings from X into itself such that

- (a) $P(X) \subset S(X), Q(X) \subset A(X)$
- (b) there exists a constant $k \in (0,1)$ such that

$$\int_0^{M(Px,Qy,qt)} \varphi(t) dt \geq \int_0^{m(x,y,t)} \varphi(t) dt$$

Where $\varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a Lebesgue – integrable mapping which is summable, non negative, and such that

$$\int_0^\varepsilon \varphi(t) dt > 0 \text{ for each } \varepsilon > 0$$

Where $m(x,y,t) = \min\{M(Px,Qy,qt), M(Ax,Sy,t), M(Px,Ax,t), M(Qy,Sy,t), M(Px,Sy,t)\}$

For all $x, y \in X$ and $t > 0$

- (c). $P(x), AB(x)$, or $ST(x)$ is complete subspace of X , then
 - (i). P and AB have a coincidence point, and
 - (ii). Q and St have a coincidence point.
- (d). the pair $\{P, A$ and $\{Q, S\}$ are compatible, Then A, S, P and Q have a unique common fixed point in X .

REFERENCES

- [1] A. Altun, D. Turkoglu and B. E. Rhoades, Fixed points of weakly compatible maps satisfying a general contractive condition of integral type, Fixed Point Theory and Applications, Vol. 2007, Article ID 17301, 9 pages.
- [2] A. Branciari, A fixed point theorem for mappings satisfying a general contractive condition of integral type, International J. of Mathematics and Mathematical Sciences, 29(2002), 531-536.23
- [3] A. George and P. Veeramani, On some results in fuzzy metric spaces, Fuzzy Sets and Systems, 64(1994), 395-399.
- [4] B. E. Rhoades, Two fixed point theorems for mappings satisfying a general contractive condition of integral type, International J. of Mathematics and Mathematical Sciences, 63(2003), 4007-4013.
- [5] B. Schweizer and A. Sklar, Statistical metric spaces, Pacific Journal Math., 10(1960), 314-334.
- [6] G. Jungck, Commuting mappings and fixed points, Amer. Math. Monthly, 83(1976), 261-263.
- [7] G. Jungck, Commuting mappings and fixed points, Internat. J. Math. and Math. Sci., 9(1986), No. 4, 771-779.
- [8] G. Jungck, P.P. Murthy and Y. J. Cho, Compatible mappings of type (A) and common fixed points, Math. Japon., 38(1993), No.2, 381-390.

- [9] G. Jungck and B.E. Rhoades, Fixed point for set valued functions without continuity, *Ind. J. Pure and Appl. Math.*, 29(3)(1998), 227-238.
- [10] H.K. Pathak, Y.J. Cho, S.S. Chang and S.M. Kang, Compatible mappings of type (P) and fixed point theorems in metric spaces and probabilistic metric spaces, *Novi Sad J. Math.*, 26(1996), No.2, 87-109.
- [11] J. X. Fang, On fixed points theorems in fuzzy metric spaces, *Fuzzy Sets and Systems*, 46(1992), 107-113.
- [12] J.S. Jung, Y.J. Cho and J.K. Kim, Minimization theorems for fixed point theorems in fuzzy metric spaces and applications, *Fuzzy Sets and Systems*, 61(1994), No.2, 199-207.
- [13] L. A. Zadeh, Fuzzy sets, *Inform control*, 8(1965), 338-353.
- [14] M. A. Erceg, Metric spaces in fuzzy set theory, *J. Math. Anal. Appl.*, 69(1979), 205-230.
- [15] M. Grabiec, Fixed point in fuzzy metric spaces, *Fuzzy Sets and Systems*, 27(1988), 385-389.
- [16] O. Hadzic, Fixed point theory in probabilistic metric spaces, *Navi Sad: Serbian Academy of Science and Arts*, 1995.24
- [17] O. Karmosil and J. Michalek, Fuzzy metric and statistical metric spaces, *Kybernetika*, 11(1975), 326-334.
- [18] O. Kaleva and S. Seikkala, On fuzzy metric spaces, *Fuzzy Sets and Systems*, 12(1984), 225-229.
- [19] P. Vijayaraju, B. E. Rhoades and R. Mohanraj, A fixed point theorem for a pair of maps satisfying a general contractive condition of integral type, *International J. of Mathematics and Mathematical Sciences*, 15(2005), 2359-2364.
- [20] R. Kannan, Some results on fixed points, *Bull. Calcutta Math. Soc.*, 60(1968), 71-76.
- [21] R. Vasuki, Common fixed point theorem in a fuzzy metric space, *Fuzzy Sets and Systems*, 97(1998), 395-397.
- [22] S.S. Chang, Y.J. Cho, B.S. Lee, J.S. Jung and S.M. Kang, Coincidence point theorems and minimization theorems in fuzzy metric spaces, *Fuzzy Sets and Systems*, 88(1997), No.1, 119-127.
- [23] S.N. Mishra, N. Sharma and S. L. Singh, Common fixed points of maps on fuzzy metric spaces, *Internat. J. Math. and Math. Sci.*, 17(1994), No. 2, 253-258.
- [24] S. Muralisankar and G. Kalpana, Common fixed point theorem in intuitionistic fuzzy metric spaces using general contractive condition of integral type, *Int. J. Contemp. Math. Sciences*, 4(2009), No.11, 505-518.
- [25] S. Sessa, On a weak commutativity condition of mappings in fixed point considerations, *Publ. Inst. Math.*, 32(1982), No.46, 149-153.
- [26] S. Sedghi, N. Shobe and A. Aliouche, Common fixed point theorems in intuitionistic fuzzy metric spaces through condition of integral type, *Applied Mathematics and Information sciences*, 2(1)(2008), 61-82.
- [27] S. Sedghi, N. Shobe and A. Aliouche, Common fixed point theorems in fuzzy metric spaces through conditions of integral type, *Journal of Fuzzy Mathematics*, 17(3) (2009), 687-698.
- [28] Sushil Sharma, On fuzzy metric spaces, *South East Asian Bulletin of Math.*, 26(2002), No.1, 145-157.
- [29] Sushil Sharma, Common fixed point theorems in fuzzy metric spaces, *Fuzzy Sets and Systems*, 127(2002), 345-352.25
- [30] Sushil Sharma and B. Deshpande, Common fixed points without continuity in fuzzy metric spaces, *J. Korean Soc. Educ. Ser. B*, 12(2005), No.4, 289-306.
- [31] T. Suzuki, Meir-Keeler contractions of integral type are still Meir-Keeler contractions, *International J. of Mathematics and Mathematical Sciences*, Vol.2007, Article ID 39281, 6 pages.
- [32] V. Gregori and A. Sapena, On fixed point theorem in fuzzy metric spaces, *Fuzzy Sets and Systems*, 125(2002), 245-252.

- [33] Y.J. Cho, Fixed point in fuzzy metric spaces, Journal of Fuzzy Mathematics, 5(1997), No.4, 949 –962.
- [34] Y.J. Cho, H.K. Pathak, S.M. Kang and J.S. Jung, Common fixed points of compatible maps of type (b) on fuzzy metric spaces, Fuzzy Sets and Systems, 93(1998), 99-11.
- [35] Z.K. Deng, Fuzzy pseudo metric spaces, J. Math. Anal. Appl., 86(1982), 84-75.

Source of Support: Nil, Conflict of interest: None Declared