



**A FIELD ALGEBRA STRUCTURE
ON THE REPRESENTATION OF THE AFFINE LIE ALGEBRA**

Tianzeng Li & Yu Wang*

School of Science, Sichuan University of Science and Engineering, Zigong 643000, China.

(Received on: 26-10-13; Revised & Accepted on: 25-11-13)

ABSTRACT

The vertex operator structure on the representation v_α of the affine algebra associated with $SL(n, \mathbb{C})$ is studied by using the representation theory of Lie algebra. Moreover, it is proved that v_α is a field algebra according to calculus methods of formal distributions.

Key words: field algebra; vertex operator; n-th product.

1. INTRODUCTION

The physicists brought forward the concept of vertex operation algebra in studying the theory of field and string. It is important in studying representation theory and finite group. Meurmen and Lepowsky solved the Guss of McKay-Thompson with this theory. And Borchers used the vertex operation algebra and Kac-Moody Lie algebra to solve the famous problem of the Monstrous Moonshine Conjecture and won fields award in 1998. Frenkel and Kac^[1,2] had constructed the level-one representations of affine Kac-Moody algebras $A_n^{(1)}, D_n^{(1)}, E_6^{(1)}, E_7^{(1)}, E_8$ by means of vertex operators in 1981. In addition, Xu and Jiang^[3] have introduced another set of vertex operators in 1990 which are constructed for the level-one representations of the cases $B_n^{(1)}$ and $G_2^{(1)}$. Xu^[4] gave the level-one representations of the affine Lie algebras with first kind in 1991.

Using these vertex operators to construct a field algebra has been an important subject of study. The representations V_α of Kac-Moody Lie algebra associated to $SL(2, \mathbb{C})$ are constructed, which are based on a certain untwisted or twisted vertex operators, and it is proofed to be a vertex operator algebra in Ref.[5]. In this paper, we use the vertex operators of affine Lie algebra $SL(n, \mathbb{C})$ with first kind to construct field algebras.

2. THE VERTEX REPRESENTATION V_α OF $SL(n, \mathbb{C})$

In this section, we briefly introduce the structure of V_α and vertex operators $Y(v, z)$ on V_α . Let $B(x, y)$ be the Killing form of a finite dimensional complex simple Lie algebra $SL(n, \mathbb{C})$. Let \mathfrak{h} ($\dim(\mathfrak{h}) = n$) be a Cartan subalgebra of $SL(n, \mathbb{C})$ and Δ be the root system. Then $SL(n, \mathbb{C}) = \mathfrak{h} + \sum_{\alpha \in \Delta} g_\alpha$, g_α is the root subspace decomposition of by the Cartan subalgebra \mathfrak{h} . Denote $\Pi = \{ \alpha_1, \dots, \alpha_n \} \subset \mathfrak{h}^*$ a simple root system, where \mathfrak{h}^* is the dual space of \mathfrak{h} .

The root lattice

$$L = \{ \alpha = \sum_{i=1}^n m_i \alpha_i \mid m_1, \dots, m_n \in \mathbb{Z} \} \subset \mathfrak{h}_r^* \tag{1}$$

is an abelian addition group in the real linear space \mathfrak{h}_r^* . Then \mathfrak{h}_r^* has an inner product

$(x, y) = c_0 B(x, y)$, $\forall x, y \in \mathfrak{h}_r^*$, where c_0 is a positive constant. The group algebra $\mathbb{C}(e^L)$ of L is an abelian associative algebra with the basis $\{ e^\alpha \mid \alpha \in L \}$, where $e^0 = 1$ and $e^\alpha e^\beta = e^{\alpha+\beta}$. Denote

$h_i(m) = t^m \otimes \alpha_i$, $m \in \mathbb{Z}$, $1 \leq i \leq n$, $X_m(\alpha) = t^m \otimes e_\alpha$, $\forall \alpha \in \Delta$, where t is a complex parameter. Let S^- be the complex linear space spanned by the basis $1, h_i(-m)$, $m \in \mathbb{Z}^+$, $1 \leq i \leq n$. Denote $S(S^-)$ be the symmetric tensor algebra over \mathbb{C} generated by S^- with the product \vee . Then $S(S^-)$ is a commutative associated algebra with the unit element 1 and has a basis $1, h_{i_1}(-m_1) \vee \dots \vee h_{i_s}(-m_s)$, $1 \leq i_1 \leq \dots \leq i_s \leq n, m_1, \dots, m_s \in \mathbb{Z}^+, s \in \mathbb{Z}^+$

***Corresponding author: Yu Wang*, E-mail: 281505815@qq.com**

Let $V = S(S^-) \otimes C(e^L)$. The formal linear combination of finite or infinite elements of the basis forms a complete space V_Q of V . It is well known that V_Q is an associative algebra with

$$(h_{i_1}(-m_1) \vee \dots \vee h_{i_s}(-m_s)) \otimes e^\alpha = (1 \otimes e^\alpha) \prod_{k=1}^s (h_{i_k}(-m_k) \otimes e^0). \tag{2}$$

Hence the representation space V_Q has a basis

$$1 \otimes e^\beta, (h_{i_1}(-m_1) \vee \dots \vee h_{i_s}(-m_s)) \otimes e^\beta = (1 \otimes e^\beta) \prod_{k=1}^s (h_{i_k}(-m_k) \otimes e^0),$$

where $\beta \in L, 1 \leq i_1 \leq \dots \leq i_s \leq n, m_1, \dots, m_s \in \mathbb{Z}^+, s = 1, 2, \dots$.

For any $u \otimes e^\beta \in V_Q$, the degree of $u \otimes e^\beta$ is defined by

$$\deg(u \otimes e^\beta) = \deg(u) + \frac{1}{2}(\beta, \beta),$$

where $\deg(u)$ is defined by

$$\deg(1) = 0, \deg[h_{i_1}(-m_1) \vee \dots \vee h_{i_s}(-m_s)] = \sum_{k=1}^s m_k.$$

Now, we introduce some linear operators acting on V_Q for the definition of the vertex operator representation of the affine Lie algebra, vertex operator algebra and vertex algebra.

(I) Let $D : V_Q \rightarrow V_Q$ be a linear operator, which is defined by

$$D(v \otimes e^\beta) = \text{deg } v \otimes \alpha^\beta(v \otimes e^\beta). \tag{3}$$

(II) Let $\partial_{h_{i_1}(m_1)}, 1 \leq i \leq n, m_i \in \mathbb{Z}$ be the linear differential operators acting on the linear space $S(S^-)$, which are defined by

$$\partial_{h_{i_1}(m_1)}(h_{i_2}(-m_2)) = m_1 \delta_{m_1, -m_2}(\alpha_{i_1}, \alpha_{i_2}), m_1, m_2 \in \mathbb{Z}^+. \tag{4}$$

Let $\alpha_i(m_i)$ be a linear operator acting on V_Q , which is defined by

$$\begin{cases} \alpha_i(-m_i)(v \otimes e^\beta) = (h_{i_1}(-m_1) \vee v) \otimes e^\beta, & \text{when } m_i \in \mathbb{Z}^+, \\ \alpha_i(0)(v \otimes e^\beta) = (\alpha_i, \beta)(v \otimes e^\beta), \\ \alpha_i(m_i)(v \otimes e^\beta) = \partial_{h_{i_1}(m_1)}(v) \otimes e^\beta, & \text{when } m_i \in \mathbb{Z}^-. \end{cases}$$

where $v \in S(S^-), e^\beta \in C(e^L)$.

Lemma: 1 $[\alpha_i(m_i), \alpha_j(q_j)] = m_i \delta_{m_i, -q_j}(\alpha_i, \alpha_j) \text{id}, m_i, q_j \in \mathbb{Z}$.

Proof: This formula has been easily proved in Refs.[6,7].

Let $\alpha = \sum_{i=1}^n a_i \alpha_i \in L$. The linear operator $\alpha(m)$ acting on V is defined by $\alpha(m) = \sum_{i=1}^n a_i \alpha_i(m), \forall m \in \mathbb{Z}$.

By the induction, we have

Lemma: 2 Let $\alpha(m) = \sum_{i=1}^n a_i \alpha_i(m), \beta(q) = \sum_{j=1}^n b_j \alpha_j(q)$, then

$$[\alpha(m), \beta(q)] = m \delta_{m+q, 0}(\alpha, \beta) \text{id} \tag{5}$$

Particularly,

$$\exp(\alpha(m)) \exp(\beta(-m)) = \exp(m(\alpha, \beta)) \exp(\beta(-m)) \exp(\alpha(m))$$

(III) The mapping $\varepsilon : L \times L \rightarrow \{e^{i\theta} | 0 \leq \theta < 2\pi\}$ is called the ε -mapping, if ε satisfies the following conditions:

- (i) $\varepsilon(0, \beta) = \varepsilon(\beta, 0) = 1, \forall \beta \in L$;
- (ii) $\varepsilon(\alpha, \beta) = (-1)^{(\alpha, \beta)} \varepsilon(\beta, \alpha), \forall \alpha, \beta \in L$;
- (iii) $\varepsilon(\beta + \gamma, \alpha) \varepsilon(\beta, \gamma) = \varepsilon(\beta, \alpha + \gamma) \varepsilon(\gamma, \alpha), \forall \alpha, \beta, \gamma \in L$.

(IV) Let $x(\alpha, z) : \mathbb{C}(e^t) \rightarrow \mathbb{C}(e^t)$ be a linear operator, which is defined by

$$x(\alpha, z)(v \otimes e^\gamma) = \varepsilon(\alpha, \gamma) z^{(\alpha, \gamma)} (v \otimes e^{\alpha+\gamma}) \tag{6}$$

Let $X(\alpha, z) = \sum_{m=-\infty}^{\infty} X_m(\alpha) z^{-m}$ be the Laurent series of z . Then

$$X(\alpha, z)(v \otimes e^\gamma) = E^+(\alpha, z) E^-(\alpha, z) x(\alpha, z)(v \otimes e^\gamma) = \text{ex} \sum_{m=1}^{\infty} \frac{z^m}{m} \alpha(-m) \text{ex} \sum_{m=1}^{\infty} \frac{-z^{-m}}{m} \alpha(m) \varepsilon(\alpha, \gamma) z^{(\alpha, \gamma)} (v \otimes e^{\alpha+\gamma}).$$

Theorem: 1 The vertex operator representation (ρ, V_ρ) of affine Lie algebras with the first kind can be defined on the generators by

$$\begin{cases} \rho(c) = \text{id}, \\ \rho(d) = D, \\ \rho(t^m \otimes \alpha_i) = \alpha_i(m), \quad 1 \leq i \leq n, \quad m \in \mathbb{Z}, \\ \rho(t^m \otimes e_\alpha) = X_m(\alpha), \quad \alpha \in \Delta, \quad m \in \mathbb{Z}. \end{cases}$$

This theorem is proofed in Pre[8]. In this case, the multiplication table of the affine Lie algebra is

$$[\text{id}, D] = [\text{id}, \alpha_i(m)] = [\text{id}, X_m(\alpha)] = 0, [D, \alpha_i(m)] = m\alpha_i(m),$$

$$[D, X_m(\alpha)] = mX_m(\alpha), [\alpha_i(m), \alpha_j(k)] = m\delta_{m,-k}(\alpha_i, \alpha_j)\text{id},$$

$$[\alpha_i(m), X_k(\alpha)] = (\alpha, \alpha_i) X_{m+k}(\alpha),$$

$$[X_m(\alpha), X_k(-\alpha)] = \varepsilon(\alpha, -\alpha)(\alpha(m+k) + m\delta_{m,-k}\text{id}),$$

$$[X_m(\alpha), X_k(\beta)] = 0, \alpha, \beta \in \Delta, \alpha + \beta \notin \Delta \cup \{0\}, [X_m(\alpha), X_k(\beta)] = \varepsilon(\alpha, \beta) X_{m+k}(\alpha + \beta), \alpha, \beta, \alpha + \beta \in \Delta$$

3. THE STRUCTURE OF FIELD ALGEBRA ON V_ρ

Definition: 1 A complex linear space V is called a field algebra, if there exist a set of linear operators (every linear operator is called a field) for v :

$$Y(v, z) = \sum_{m \in \mathbb{Z}} v_{(m)} z^{-m-1} \in \text{End}V[[z, z^{-1}]], \tag{7}$$

such that given any $v, w \in V$, there is a positive integer $m_0 = m_0(v, w)$, such that $v_{(m)}(w) = 0, \forall m > m_0$. And there is a fixed vector $|0\rangle \in V$, which is called by the vacuum vector, such that

(i) (vacuum)

$$Y(|0\rangle, z) = \text{id}_V, \quad Y(v, z)|0\rangle_{|z=0} = v.$$

(ii) (translation covariance)

$T \in \text{End}(V)$ is defined by $T(v) = v_{(-2)}|0\rangle, \forall v \in V$. T is called the infinitesimal translation operator,

if T is a derivation on V and satisfies the condition: $\text{ad}(T) = \partial_z$ acting on any linear operator $Y(u \otimes e^\gamma, z)$.

(iii) (weak locality) There exists a positive integer N , such that

$$\text{Res}_z (z-w)^N [Y(u, z), Y(v, w)] = 0, \forall u, v \in V.$$

Define the map

$$\begin{aligned} Y(\cdot, z) : V_\rho &\longrightarrow (\text{End}V_\rho)[[z, z^{-1}]] \\ v &\longrightarrow \sum_{n \in \mathbb{Z}} v_n z^{-n-1} \end{aligned}$$

by the following way:

$$Y(1, z) = Id_{V_\alpha}; \quad Y(h_i(-1) \otimes 1, z) = H_i(z) = \sum_{m \in \mathbb{Z}} H_i(m) Z^{-m-1};$$

$$Y(h_i(-m) \otimes 1, z) = \partial^{(m-1)} H_i(z); \quad Y(1 \otimes e^\alpha) = X(\alpha, z) = E^+(\alpha, z) E^-(\alpha, z) x(\alpha, z);$$

$$Y(h_{i_1}(-m_1) \vee \dots \vee h_{i_s}(-m_s) \otimes 1, z) =: \partial^{(m_1-1)} H_{i_1}(z) \dots \partial^{(m_s-1)} H_{i_s}(z); \quad Y(h_{i_1}(-m_1) \vee \dots \vee h_{i_s}(-m_s) \otimes e^\beta, z) =: \partial^{(m_1-1)} H_{i_1}(z) \dots \partial^{(m_s-1)} H_{i_s}(z) Y(1 \otimes e^\beta, z);$$

where $\cdot \vee \cdot$ is the normal order of fields or operators, $\partial^{(m)} = \frac{\partial^m}{m!}$ is a differential operator, and the vacuum vector $|0\rangle = 1 \otimes e^0$.

In the following, we shall check that $(V, Y(v \otimes e^\alpha, z))$ is a field algebra, i.e., it satisfies the three axioms of the field algebra.

3.1. The vacuum axiom

- (1) $Y(1 \otimes 1, z) = Id_V$;
- (2) $Y(h_i(-m) \otimes 1, z) 1 \otimes 1|_{z=0} = h_i(-m) \otimes 1$;
- (3) $Y(v \otimes e^\alpha, z) 1 \otimes 1|_{z=0} = v \otimes e^\alpha$.

This formula can be easily proved by the Theorem 1.

3.2. The translation covariance axiom

Let $v = \prod_{k=1}^i h_{i_k}(-m_k)$, $v_k = \prod_{1 \leq j \leq s, j \neq k} h_{i_j}(-m_j)$, $m_i \in \mathbb{Z}^+$. Then

$$T(v \otimes e^\gamma) = \gamma(-1)(v \otimes e^\gamma) + \sum_{k=1}^i m_k h_{i_k}(-1+m_k)(v_k \otimes e^\gamma).$$

T is a derivation which acts on V_Q . Particularly,

$$T(1 \otimes e^\gamma) = \gamma(-1)(1 \otimes e^\gamma), \quad T(h_i(-m) \otimes 1) = m h_i(-m-1) \otimes 1.$$

Lemma: 3 Given $i = 1, 2, \dots, n$, $m \in \mathbb{Z}_+, m > 0$, $\alpha \in L$, then

- (i) $\text{ad}(T)\alpha(-m) = m\alpha(-1+m)$;
- (ii) $\text{ad}(T)\alpha(1)(v \otimes e^{\alpha+\gamma}) = -(\alpha, \alpha + \gamma)(v \otimes e^{\alpha+\gamma})$;
- (iii) $\text{ad}(T)\alpha(m) = -m\alpha(m-1)$;
- (iv) $\text{ad}(T)x(\alpha, z) = \alpha(-1)x(\alpha, z)$;
- (v) $\text{ad}(T)E^+(\alpha, z) = E^+(\alpha, z) \sum_{m=2}^{\infty} z^{m-1} \alpha(-m)$;
- (vi) $\text{ad}(T)E^-(\alpha, z) = [\sum_{m=1}^{\infty} z^{-m-1} \alpha(m) + z^{-1}(\alpha, \alpha + \gamma)]E^-(\alpha, z)$;

$$E^-(\alpha, z)[T, x(\alpha, z)](v \otimes e^\gamma) = [\alpha(-1) - z^{-1}(\alpha, \alpha)]E^-(\alpha, z)x(\alpha, z)(v \otimes e^\gamma).$$

Proof: The formulas (i-iii, v) is easily proved[11]. Now, we only check the formulas (iv) and (v).

$$\begin{aligned} \text{ad}(T)E^+(\alpha, z) &= \text{ad}(T) \prod_{n=1}^{\infty} \exp \frac{1}{m} z^n \alpha(-m) \\ &= \sum_{j=1}^{\infty} \prod_{n=1}^{j-1} \exp \frac{1}{m} z^n \alpha(-m) \text{ad}(T) \cdot \exp \frac{1}{j} z^j \alpha(-j) \prod_{n=j+1}^{\infty} \exp \frac{1}{m} z^n \alpha(-m) \\ &= E^+(\alpha, z) \sum_{j=1}^{\infty} z^j \alpha(-1+j). \end{aligned}$$

Hence (iv) holds.

Lemma: 4 The infinitesimal translation operator T satisfies the translation covariance axioms

$$[T, Y(v \otimes e^\gamma)] = \partial_z Y(v \otimes e^\gamma). \tag{8}$$

Proof: Since ∂_z is a differential operator acting on $Y(u \otimes e^\alpha)$ about z , then

$$\begin{aligned} \partial_z(Y(1 \otimes e^\alpha, z))(v \otimes e^\gamma) &= \partial_z(E^+(\alpha, z)E^-(\alpha, z)x(\alpha, z))(v \otimes e^\gamma) \\ &= \partial_z(E^+(\alpha, z))E^-(\alpha, z)x(\alpha, z) + E^+(\alpha, z)\partial_z(E^-(\alpha, z))x(\alpha, z) + E^+(\alpha, z)E^-(\alpha, z)\partial_z(x(\alpha, z))(v \otimes e^\gamma) \\ &= E^+(\alpha, z)[\sum_{m=1}^{\infty} z^{-m-1}\alpha(-m) + \sum_{m=1}^{\infty} z^{-m-1}\alpha(m) + z^{-1}(\alpha, \gamma)]E^-(\alpha, \sqrt{z})(\partial_z x(\alpha, \sqrt{z}))(v \otimes e^\gamma). \end{aligned}$$

By the formulas (iv), (v) and (vi) of Lemma 3, we have

$$\begin{aligned} [TY(1 \otimes e^\alpha, z)](v \otimes e^\gamma) &= [TE^+(\alpha, z)E^-(\alpha, z)x(\alpha, z)](v \otimes e^\gamma) \\ &= [[TE^+(\alpha, z)]E^-(\alpha, z)x(\alpha, z) + E^+(\alpha, z)[TE^-(\alpha, z)]x(\alpha, z) + E^+(\alpha, z)E^-(\alpha, z)[Tx(\alpha, z)]](v \otimes e^\gamma) \\ &= E^+(\alpha, z)[\sum_{m=1}^{\infty} z^{-m-1}\alpha(-m) + \sum_{m=1}^{\infty} z^{-m-1}\alpha(m) + z^{-1}(\alpha, \gamma)]E^-(\alpha, z)x(\alpha, z)(v \otimes e^\gamma), \end{aligned}$$

Therefore

$$[TY(1 \otimes e^\alpha, z)] = \partial_z(Y(1 \otimes e^\alpha, z)).$$

It is easy to see that

$$[TY(h_i(-m) \otimes e^0, z)] = \partial_z(Y(h_i(-m) \otimes e^0, z))$$

By the formulas (ii), (iv) and (vi) of Lemma 3. By the induction, we have

$$[TY(v \otimes e^0, z)] = \partial_z(Y(v \otimes e^0, z)), \text{ and}$$

$$[TY(u \otimes e^\alpha, z)] = \partial_z(Y(u \otimes e^\alpha, z)),$$

where $u = \prod_{k=1}^s h_k(-m_k)$. Then

$$\begin{aligned} \text{ad}(T)Y(h_i(-m_0) \vee u \otimes e^\alpha, z) &= \text{ad}(T) : \partial^{(m_0-1)} H_i(z)Y(u \otimes e^\alpha) : \\ &= \text{ad}(T)[\partial^{(m_0-1)} H_i(z)_+ Y(u \otimes e^\alpha)] + \text{ad}(T)[Y(u \otimes e^\alpha) \partial^{(m_0-1)} H_i(z)_-] \\ &= [\text{ad}(T) \partial^{(m_0-1)} H_i(z)_+]Y(u \otimes e^\alpha) + \partial^{(m_0-1)} H_i(z)_+ [\text{ad}(T)Y(u \otimes e^\alpha)] \\ &\quad + [\text{ad}(T)Y(u \otimes e^\alpha)] \partial^{(m_0-1)} H_i(z)_- + Y(u \otimes e^\alpha) [\text{ad}(T) \partial^{(m_0-1)} H_i(z)_-] \\ &= \partial_z Y(h_i(-m_0) \vee u \otimes e^\alpha) + [(\text{ad}(T) - \partial_z) \partial^{(m_0-1)} H_i(z)_+]Y(u \otimes e^\alpha) + Y(u \otimes e^\alpha) [(\text{ad}(T) - \partial_z) \partial^{(m_0-1)} H_i(z)_-] \end{aligned}$$

Hence we need to prove that

$$[(\text{ad}(T) - \partial_z) \partial^{(m_0-1)} H_i(z)_+]Y(u \otimes e^\alpha) + Y(u \otimes e^\alpha) [(\text{ad}(T) - \partial_z) \partial^{(m_0-1)} H_i(z)_-] = 0.$$

From the formulas (i) and (iii), it is easy to prove

$$[(\text{ad}(T) - \partial_z) \partial^{(m_0-1)} H_i(z)_+] = 0, [(\text{ad}(T) - \partial_z) \partial^{(m_0-1)} H_i(z)_-] = 0.$$

Hence (10) holds.

3.3. The weak locality axiom

In this subsection, we check the locality axiom. Namely, we will prove the following formula holds:

$$\text{Res}_z(z-w)^N [Y(u \otimes e^\alpha, z), Y(v \otimes e^\beta)] = 0, \quad \forall u \otimes e^\alpha, v \otimes e^\beta \in V_0, z, w \in \mathbb{C}, N \gg 0.$$

Lemma: 5 $Y(h_i(-1) \otimes 1, z)$ and $Y(h_j(-1) \otimes 1, w)$ are weak local, i.e.,

$$\text{Res}_z(z-w)^N [Y(h_i(-1) \otimes 1, z), Y(h_j(-1) \otimes 1, w)] = 0, \quad N \gg 0.$$

Proof: Notice that

$$[Y(h_i(-1) \otimes 1, Y(h_j(-1) \otimes 1, w))] = \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} [H_i(m) H_j(n)] z^{-m-1} w^{-n-1} = (\alpha_i, \beta_j) \sum_{m \in \mathbb{Z}} m z^{-m-1} w^{-m-1} = (\alpha_i, \beta_j) \partial_w \delta(z-w).$$

By the formula (iv) of Lemma 3, it is easy to prove

$$(z-w)^2 [Y(h_i(-1) \otimes 1 Y(h_j(-1) \otimes 1, w)] = (\alpha_i, \beta_j)(z-w)^2 \partial_- \delta(z-w) = 0,$$

then

$$\text{Res}_z (z-w)^2 [Y(h_i(-1) \otimes 1 Y(h_j(-1) \otimes 1, w)] = \text{Res}_z (\alpha_i, \beta_j)(z-w)^2 \partial_- \delta(z-w) = 0$$

Lemma: 6^[12] $Y(h_i(-m_i) \otimes 1, z)$ and $Y(h_j(-n_j) \otimes 1, z)$ are weak local, i.e.,

$$\text{Res}_z (z-w)^N [Y(h_i(-m_i) \otimes 1, z), Y(h_j(-n_j) \otimes 1, w)] = 0, N > m_i + n_j.$$

Proof: Since

$$\begin{aligned} Y(h_i(-m_i) \otimes 1, z) \cdot Y(h_j(-n_j) \otimes 1, w) &= Y(h_j(-n_j) \otimes 1, w) \cdot Y(h_i(-m_i) \otimes 1, z) \\ &+ (\alpha_i, \alpha_j) \sum_{m=0}^{\infty} z^m w^{-m-m_i-n_j} \cdot \frac{(m+m_i-1)! (m+m_i+n_j-1)!}{m!(m_i-1)! (m-m_i)!(n_j-1)!} (-m-m_i) a_{n_j}(n_j) \\ &+ (\alpha_i, \alpha_j) \sum_{n=0}^{\infty} z^{-n-m_i-n_j} w^n \cdot \frac{(n+m_i+n_j-1)! (n+n_j-1)!}{(n+n_j)!(m_i-1)! n!(n_j-1)!} (n+n_j) a_{m_i}(m_i), \end{aligned}$$

then

$$\begin{aligned} &Y(h_i(-m_i) \otimes 1, z) \cdot Y(h_j(-n_j) \otimes 1, w) - Y(h_j(-n_j) \otimes 1, w) \cdot Y(h_i(-m_i) \otimes 1, z) \\ &= (\alpha_i, \alpha_j) \left(\sum_{n=0}^{\infty} z^{-n-m_i-n_j} w^n \frac{(n+m_i+n_j-1)!}{(m_i-1)! n!(n_j-1)!} a_{m_i}(m_i) - \sum_{m=0}^{\infty} z^m w^{-m-m_i-n_j} \frac{(m+m_i+n_j-1)!}{(n_j-1)! m!(m_i-1)!} a_{n_j}(n_j) \right) \\ &= (\alpha_i, \alpha_j) \frac{(m_i+n_j-1)!}{(m_i-1)!(n_j-1)!} \left[\sum_{n=0}^{\infty} z^{-n-m_i-n_j} w^n \cdot \frac{(n+m_i+n_j-1)!}{n!(m_i+n_j-1)!} a_{m_i}(m_i) - \sum_{m=0}^{\infty} z^m w^{-m-m_i-n_j} \frac{(m+m_i+n_j-1)!}{m!(m_i+n_j-1)!} a_{n_j}(n_j) \right] \end{aligned}$$

By the formula (iv) of Lemma 3, we have

$$(z-w)^{m_i+n_j} [Y(h_i(-m_i) \otimes 1, z), Y(h_j(-n_j) \otimes 1, w)] = 0,$$

then

$$\text{Res}_z (z-w)^{m_i+n_j} [Y(h_i(-m_i) \otimes 1, z), Y(h_j(-n_j) \otimes 1, w)] = 0, N > m_i + n_j.$$

Lemma: 7 $Y(h_i \otimes 1, z)$ and $Y(1 \otimes e^\alpha, w)$ are weak local, i.e.,

$$\text{Res}_z (z-w)^N [Y(h_i \otimes 1, z), Y(1 \otimes e^\alpha, w)] = 0, N > 1.$$

Proof: Since

$$Y(h_i \otimes 1, z) = H_i(z) = \sum_{m \in \mathbb{Z}} H_i(m) z^{-m-1},$$

then $[H_i(m), Y(1 \otimes e^\alpha, w)] = w^m (\alpha_i, \alpha) Y(1 \otimes e^\alpha, w)$, and $[Y(h_i \otimes 1, z) Y(1 \otimes e^\alpha, w)] = \delta(z-w) Y(1 \otimes e^\alpha, w)$.

Therefore,

$$(z-w)[Y(h_i \otimes 1, z) Y(1 \otimes e^\alpha, w)] = 0, \text{ and } \text{Res}_z (z-w)[Y(h_i \otimes 1, z) Y(1 \otimes e^\alpha, w)] = 0$$

Lemma: 8 $Y(1 \otimes e^\alpha, z)$ and $Y(1 \otimes e^\beta, w)$ are weak local, i.e.

$$\text{Res}_z (z-w)^N [Y(1 \otimes e^\alpha, z), Y(1 \otimes e^\beta, w)] = 0, N > 1.$$

Proof: Since

$$Y(1 \otimes e^\alpha, z) = E^+(\alpha, z) E^-(\alpha, z) x(\alpha, z),$$

We have

$$\begin{aligned} Y(1 \otimes e^\alpha, z) Y(1 \otimes e^\beta, w) &= x(\alpha, z) x(\beta, w) E^+(\alpha, z) E^-(\alpha, z) E^+(\beta, z) E^-(\beta, z) \\ &= z^{-(\alpha, \beta)} (z-w)^{(\alpha, \beta)} x(\alpha, z) x(\beta, w) E^+(\alpha, z) E^+(\beta, z) E^-(\alpha, z) E^-(\beta, z). \end{aligned}$$

By the same way, we get

$$\begin{aligned} Y(1 \otimes e^\beta, z) Y(1 \otimes e^\alpha, w) &= x(\beta, w) x(\alpha, z) E^+(\beta, w) E^-(\beta, w) E^+(\alpha, z) E^-(\alpha, z) \\ &= w^{-(\alpha, \beta)} (w-z)^{(\alpha, \beta)} x(\beta, w) x(\alpha, z) E^+(\alpha, z) E^+(\beta, z) E^-(\alpha, z) E^-(\beta, z). \end{aligned}$$

From the definition of mapping ε [13], we have

$$z^{-(\alpha,\beta)}(z-w)^{(\alpha,\beta)}x(\alpha,z)x(\beta,w) = w^{-(\alpha,\beta)}(w-z)^{(\alpha,\beta)}x(\beta,w)x(\alpha,z),$$

i.e. $\text{Res}_z(z-w)[Y(1 \otimes e^\alpha, z), Y(1 \otimes e^\beta, w)] = 0$.

Lemma: 9^[13] If $a(z)$, $b(z)$ and $c(z)$ are pairwise mutually local fields, then $a(z)$, $b(z)$ and $c(z)$ are mutually local fields for all $n \in \mathbb{Z}$ (resp. $n \in \mathbb{Z}_+$). In particular $a(z)b(z)$ and $c(z)$ are mutually local fields provided that $a(z)$, $b(z)$ and $c(z)$ are.

Lemma: 10 For any $u \otimes e^\alpha, v \otimes e^\beta \in V_Q$, there exists a nonnegative integer N satisfies

$$\text{Res}_z(z-w)^N [Y(u \otimes e^\alpha, z), Y(v \otimes e^\beta, w)] = 0, \quad N \gg 0.$$

Proof: From the Lemma 5-8, we know that $Y(h_i(-1) \otimes 1, z)$, $Y(h_j(-m) \otimes 1)$ and $Y(1 \otimes e^\alpha)$ are pairwise mutually local fields. Then by the Lemma 9, we can easily prove Lemma 10.

From the fact that we have checked that $(V, Y(v \otimes e^\alpha, z))$ satisfies the three axioms. It follows

Theorem 2: $(V, Y(v \otimes e^\alpha, z))$ is field algebra.

5. ACKNOWLEDGMENT

This paper was supported by Scientific Research Fund of Sichuan Provincial Education Department grant 11ZA261, 12ZB294 and Sichuan University of Science and Engineering grant 2012PY17.

REFERENCES

- 1) Frenkel I B, Kac V G. *Basic representations of affine Lie algebras and dual resonance models* [J]. Invent Math, 1980, 62(1): 23-66.
- 2) Kac V G, Kazhdan D, Lepowsky J, Wilson R. *Realization of the basic representation of the Euclidean Lie algebras*[J]. Advances in Math, 1981, 42: 83-112.
- 3) Xu Y C, Jiang C P. *Vertex operators of G_2^1 and B_n^1* [J]. J Phys A Math Gen, 1990, 23: 3105-3121.
- 4) Xu Y C. *Vertex operators of affine Lie algebras with first kind*[C]// Proceeding of the SEAMS Conference on Ordered Structures and Algebra of computer Languages, June 1991.
- 5) Chu Y J, Cheng J F, Zheng Z J. *A vertex algebra structure on the representation V_Q of untwisted affine lie algebra $\widehat{sl}(2, \mathbb{C})$* [J].Journal of Henan University (Natural Science), 2007, 37(6):551-557.
- 6) Li H S. *Modules-at-infinity for quantum vertex algebras* [J]. Commun Math Phys, 2008, 282: 819-864.
- 7) Berman S, Dong C, Tan S. *Representations of a class of lattice type vertex algebras* [J]. J Pure Appl Algebra, 2002, 176:27-47.
- 8) Frankel I B, Lepowsky J, Meurman A. *Vertex operator algebras and the Monster* [M]. New York. Academia Press 1988.
- 9) Lu K P, Chu Y J, Zheng Z J. *P-twisted affine Lie algebra and its realizations by twisted vertex operators* [J]. Science In China [J].Series A Mathematics, 2005(48):295-305.
- 10) Frenkel I B, Jing N, Wang W. *Vertex representations via finite groups and the Mckay correspondence* [J]. IMRN, 4 (2000), 195-222.
- 11) Kac V G. *Vertex Algebras for Beginners* [M]. American Mathematical Society, 1996.
- 12) Wang F X, Jin H X, Chang X. *Relevance Vector Ranking for Information Retrieval* [J]. JCIT: Journal of Convergence Information Technology, 2010, 15(9):118-125.
- 13) Ding D S, Luo X P, Chen J F, etc. *A Convergence Proof and Parameter Analysis of Central Force Optimization Algorithm* [J]. JCIT: Journal of Convergence Information Technology, 2011, 6(10): 16-23.

Source of Support: The Opening Project of Sichuan Province University Key Laboratory of Bridge Non-destruction Detecting and Engineering Computing 2013QYJ01, Sichuan University of Science and Engineering grant 2012PY17, 2012KY06, Scientific Research Fund of Sichuan Provincial Education Department grant 11ZA261, 12ZB294. Conflict of interest: None Declared.