

COMMUTATIVITY OF ALTERNATIVE PERIODIC RINGS WITH $[(yx)^m x^m - x^m (xy)^m, x] = 0$

Y. S. Kalyan Chakravarthy* and K. Suvarna

Department of Mathematics, S. K. University, Ananthapuramu – 515001, India.

(Received on: 25-02-14; Revised & Accepted on: 23-05-14)

ABSTRACT

Let R be an alternative periodic ring. In this paper, we prove that if R is an $m(m+1)$ -torsion free alternative periodic ring satisfying the properties $[(yx)^m x^m - x^m (xy)^m, x] = 0$ for all x, y in $R \setminus N(R)$ and $N(R)$ is commutative, then R is commutative.

AMS Mathematics subject classification: 17.

Key words: Alternative rings, Periodic rings, Center.

INTRODUCTION

M. A. Khan[4] shown that R is commutative if R satisfies the conditions $[x^m, y^m] = 0$, for all x, y in $R \setminus N(R)$ and $[x, [x, y^m]] = 0$, for all x, y in $R \setminus N(R)$. He also shown that the result is valid if $[x, [x, y^m]] = 0$ is replaced by $[(yx)^m x^m - x^m (xy)^m, x] = 0$, for all x, y in $R \setminus N(R)$. In this paper, we prove that if R is an $m(m+1)$ -torsion free alternative periodic ring satisfying the properties $[(yx)^m x^m - x^m (xy)^m, x] = 0$ for all x, y in $R \setminus N(R)$ and $N(R)$ is commutative, then R is commutative.

PRELIMINARIES

Throughout this section R denotes an alternative periodic ring, $Z(R)$ the center of R , $U(R)$ the group of units of R , $J(R)$ the Jacobson radical of R , $N(R)$ the set of all nilpotent elements of R and $C(R)$ the commutator ideal of R .

In order to prove our result, we first state the well-known results:

Lemma: 1[1, Theorem 1] Let R be a periodic ring such that $N(R)$ is commutative. If for each a in $N(R)$ and x in R there exists an integer $m = m(x, a) \geq 1$ such that $[x^m, [x^m, a]] = 0$ and $[x^{m+1}, [x^{m+1}, a]] = 0$, then R is commutative. In particular, if R is a periodic ring such that $N(R)$ is commutative and $[x, [x, a]] = 0$ for all a in $N(R)$ and x in R , then R is commutative.

Lemma: 2[2, Lemma 4] Let R be a periodic ring and let $f: R \rightarrow S$ be a homomorphism of R onto S , then the nilpotents of S coincide with $f(N(R))$, where $N(R)$ is the set of nilpotents of R .

Lemma: 3[4] Let R satisfy the conditions

- (i) $[x^m, y^m] = 0$, for all x, y in $R \setminus N(R)$
- (ii) $[(yx)^m x^m - x^m (xy)^m, x] = 0$, for all x, y in $R \setminus N(R)$ and
- (iii) for any x, y in R , $m[x, y] = 0$ implies $[x, y] = 0$, then R is commutative.

MAIN RESULT

Theorem: 1 Let $m \geq 1$ be a fixed positive integer and let R be an alternative periodic ring satisfying the properties:

- (i) for any x, y in R , $m(m+1)[x, y] = 0$ implies $[x, y] = 0$ and
- (ii) $[(yx)^m x^m - x^m (xy)^m, x] = 0$, for all x, y in $R \setminus N(R)$. Further, suppose that $N(R)$ is commutative then R is commutative.

Proof: Since R is periodic and $N(R)$ is commutative, Lemma 3 yields that the commutator ideal $C(R)$ in nil that is $C(R) \subseteq N(R)$ and $N(R)$ forms an ideal of R . But $N(R)$ is commutative and also $(N(R))^2 \subseteq Z(R)$.

Corresponding author: Y. S. Kalyan Chakravarthy, E-mail: yskchakri@gmail.com

First we show that the idempotents of R are central.

Let $e^2 = e \in R$ and $r \in R$.

By replacing x by e and y by $e + er - ere$ in (ii), we get

$$((e + er - ere)e)^m e^m - e^m ((e + er - ere)e)^m \in Z(R).$$

This implies that $ere - er \in Z(R)$. Thus $ere - er = e(ere - er) = (ere - er)e = 0$.

So, $ere - er = 0$. Therefore, $ere = er$. Similarly, $ere = re$.

Thus $er = re$, for all r in R .

We now prove the theorem for R with identity 1.

Suppose that a in $N(R)$ and R in $R \setminus N(R)$. By using (ii), we get

$$[b^m(1+a)^m - (1+a)^{m+1}b^m(1+a)^{-1}, 1+a] = 0, \quad (1)$$

for all a in $N(R)$ and b in $R \setminus N(R)$. This implies that

$$\{b^m(1+a)^m - (1+a)^{m+1}b^m(1+a)^{-1}\}(1+a) = (1+a)\{b^m(1+a)^m - (1+a)^{m+1}b^m(1+a)^{-1}\}.$$

$$\text{So, } \{b^m(1+a)^{m+1} - (1+a)^{m+1}b^m\} = (1+a)\{b^m(1+a)^m - (1+a)^{m+1}b^m(1+a)^{-1}\}.$$

Using the binomial expansion and the condition $(N(R))^2 \subseteq Z(R)$, we get

$$\begin{aligned} b^m(1 + (m+1)a + \dots + (m+1)a^m + a^{m+1}) - (1 + (m+1)a + \dots + (m+1)a^m + a^{m+1})b^m \\ = (1+a)\{b^m(1+a)^m - (1+a)^{m+1}b(1+a)^{-1}\}, \end{aligned}$$

$$\text{So, } (m+1)(b^m a - ab^m) = (1+a)\{b^m(1+a)^m - (1+a)^{m+1}b(1+a)^{-1}\}. \quad (2)$$

Since $N(R)$ is a commutative ideal, $(m+1)(b^m a - ab^m) = b^m a - ab^m$, and also by (2),

$$\text{We have } (1+a)(m+1)(b^m a - ab^m) = (1+a)\{b^m(1+a)^m - (1+a)^{m+1}b(1+a)^{-1}\}.$$

Since a in $N(R)$, $1+a$ in $U(R)$ and by (1), we get

$$(m+1)(b^m a - ab^m) = (1+a)\{b^m(1+a)^m - (1+a)^{m+1}b(1+a)^{-1}\} \in Z(R).$$

This implies that, $(m+1)[b^m, a] \in Z(R)$. Since R is $m(m+1)$ -torsion free, we get

$$[b^m, a] \in Z(R), \text{ for all } a \text{ in } N(R), b \text{ in } R \setminus N(R). \quad (3)$$

Now since $N(R)$ is commutative, (2) implies that

$$[b^m, a] \in Z(R), \text{ for all } a \text{ in } N(R), b \text{ in } R. \quad (4)$$

Now let $x_1, x_2, \dots, x_n \in R$. Then $R \setminus C(R)$ is commutative. So, by Lemma 3, we get

$$(x_1 \cdot x_2 \cdot \dots \cdot x_n)^m - x_1^m \cdot x_2^m \cdot \dots \cdot x_n^m \in C(R) \subseteq N(R).$$

Therefore $N(R)$ is commutative yields that

$$[(x_1 \dots x_n)^m, a] = [x_1^m \dots x_n^m, a], \text{ for all } a \text{ in } N(R). \quad (5)$$

Combining (4) and (5), we get

$$[x_1^m \dots x_n^m, a] \in Z(R), \text{ for all } a \text{ in } N(R), x_1, x_2, \dots, x_n \in R \text{ and } n \geq 1. \quad (6)$$

Let S be the subring generated by the m -th powers of the elements of R . Then by (6), we have

$$[x, a] \in Z(S), \text{ for all } a \in N(S), x \in S. \quad (7)$$

Here $Z(S)$ and $N(S)$ represent the center of S and the set of nilpotent elements of S respectively. Combining the fact that S is periodic, $N(S)$ is commutative, and (7), Lemma 1 shows that S is commutative. Hence $[x^m, y^m] = 0$, for all x, y in R . This implies that R satisfies $[x^m, y^m] = 0$, for all x, y in R . But R also satisfies (i) and (ii).

By Lemma 3, we get the required result. It follows that for every nonzero idempotent e , eR is commutative and hence $e[x, y] = 0$, for all x, y in R . Thus if a in R is potent with $a^n = a$, $n > 1$ then $a^{n-1}[a, b] = 0 = [a, b]$, for all b in R . Since every element in a periodic ring is the sum of a potent element and nilpotent element, this gives $N(R) \subseteq Z(R)$ and R is commutative by well-known theorem of Herstein[3].

Example: 1 Let $R = \left\{ \begin{pmatrix} \alpha & \beta & \gamma \\ 0 & \alpha^2 & 0 \\ 0 & 0 & \alpha \end{pmatrix} : \alpha, \beta, \gamma \in GF(3) \right\}$. For $m = 5$, R satisfies all the hypothesis of Theorem 1 except that R is $m(m+1)$ -torsion free. But R is not commutative. This shows that the condition $m(m+1)$ -torsion free is essential in Theorem 1.

Example: 2 Let $R = \left\{ \begin{pmatrix} \alpha & \beta & \gamma \\ 0 & \alpha^2 & 0 \\ 0 & 0 & \alpha \end{pmatrix} : \alpha, \beta, \gamma \in GF(3) \right\}$. The ring R satisfies all the hypothesis of Theorem 1 except the hypothesis that $N(R)$ is commutative. This shows that commutativity of $N(R)$ is essential in Theorem 1.

REFERENCES

- [1] Abu-Khuzam, H., "A commutativity theorem for periodic rings", Math. Japon., 32(1987), pp.1-3.
- [2] Abu-Khuzam, H. and Yaqub, A., "Commutativity of rings satisfying some polynomial constraints", Acta Math. Hungar., 67(1995), pp.207-217.
- [3] Herstein, I.N., "A note on rings with central nilpotent elements", Proc. Amer. Math. Soc., 5(1954), pp.620.
- [4] Khan, M.A., "Classification of rings satisfying some constraints on subsets", Archivum Mathematicum, Tomus 43(2000), pp.19-29.

Source of Support: Nil, Conflict of interest: None Declared

[Copy right © 2014 This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]