



FIXED POINT THEOREM IN Menger SPACE

Kavita Rohit (Shrivastava)*

Department of Mathematics and Statistics,
Dr. Harisingh Gour Vishwavidyalaya, Sagar - (M.P.), India.

(Received on: 26-05-14; Revised & Accepted on: 09-06-14)

ABSTRACT

In this paper, we have accomplished the task of generalizing the fixed point theorems due to Som [3], Taskovic [4] Mukherjee [1] in the context of semi-compatibility and weak compatibility in Menger space and also it has been applied to prove common fixed point theorems for three, four and sequence of mappings. A coincidence point theorem is also established for a multi-valued mapping satisfying a generalized condition of Hausdorff distance function in (T, f, x) -orbitally complete Menger space. This theorem is generalized result of Tiwari and Shrivastava [5] and Singh [2].

BASIC PRELIMINARIES

Following notations and definitions will be used in this paper.

$CL(X) = \{A : A \text{ is non-empty closed subset of } X\}$.

Definition 1: Let T be a multi-valued mapping on a Menger space (X, F, Δ) and $x_0 \in X$. A sequence $\{x_n\}$ in X said to be an **orbit of T at x_0** denoted by $o(T, x_0)$ if $x_{n-1} \in T^n(x_0)$, i.e., $x_n \in Tx_{n-1}, \forall n \in \mathbb{N}$.

If T is a self mapping then the sequence $\{x_n\}, x_n = T^{n-1}(x_0), \forall n \in \mathbb{N}$, is the **orbit of T at x_0** .

Definition 2: A Menger space $(X, F\Delta)$ is said to be **T -orbitally complete** iff every cauchy sequence of the form $(x_{n_i} : x_{n_i} \in Tx_{n_{i-1}})$ converges in X .

Definition 3: Let S and T be a single-valued mappings and multi-valued mapping respectively on a Menger space (X, F, Δ) . A Menger space (X, F, Δ) is said to be **(T, S, X) -orbitally complete** iff every cauchy sequence of the form $(Sx_{n_i} : Sx_{n_i} \in Tx_{n_{i-1}})$ converges in X .

Definition 4: A multi-valued mapping T in Menger space (X, F, Δ) is said to be **asymptotically regular at x_0** , if for each sequence $\{x_n\}$ in $X, x_n \in Tx_{n-1}$ and $F_{x_n, x_{n+1}}(t) \rightarrow 1$ as $n \rightarrow \infty$, for all $t > 0$

Definition 5: Let S and T be a single-valued mappings and multi-valued mapping respectively on a Menger space (X, F, Δ) . A point x in X is said to be a **coincidence point** of S and T if $Sx \in Tx$.

Definition 6: Self mappings A and S of a menger space $(X, F\Delta)$ are called **semi-compatible** if $F_{AS_n, S_u}(\epsilon) \rightarrow 1$, for all $\epsilon > 0$, whenever $\{x_n\}$ is a sequence in X such that $Ax_n, Sx_n \rightarrow u$, for some u in X .

Proposition 7: Let A and S be self mappings on a Menger space $(X, E\Delta)$ with $\Delta(a, a) \geq a$, for all $a \in [0, 1]$. If S is continuous then (A, S) is semi-compatible iff (A, S) is compatible.

Corresponding author: Kavita Rohit (Shrivastava)

Department of Mathematics and Statistics, Dr. Harisingh Gour Vishwavidyalaya,
Sagar - (M.P.), India. E-mail: Kavita.rohit@rediffmail.com

Proof: Consider a sequence $\{x_n\}$ in X such that $\{Ax_n\} \rightarrow u$ and $\{Sx_n\} \rightarrow u$, As S is continuous, we have $Sx_n \rightarrow Su$.

Suppose that (A, S) is semi-compatible, then for given (ϵ, λ) , we have a positive integer $N_0(\epsilon, \lambda)$ such that

$$F_{AS_n, Su} \left(\frac{\epsilon}{2} \right) \geq 1 - \lambda \quad \text{and} \quad F_{SA_n, Su} \left(\frac{\epsilon}{2} \right) \geq 1 - \lambda, \quad \forall n > N_0.$$

Now

$$\begin{aligned} F_{ASx_n, SAx_n}(\epsilon) &\geq \Delta \left(F_{ASx_n, Su} \left(\frac{\epsilon}{2} \right), F_{SAx_n, Su} \left(\frac{\epsilon}{2} \right) \right) \\ &\geq \Delta(1-\lambda, 1-\lambda), \quad \forall n \geq N_0 \\ &\geq 1-\lambda, \quad \forall n \geq N_0. \end{aligned}$$

Hence the pair (A, S) is compatible.

Conversely, let the pair (A, S) be compatible. Then for given (ϵ, λ) , we have a positive integer $N_0(\epsilon, \lambda)$ such that

$$F_{ASx_n, SAx_n} \left(\frac{\epsilon}{2} \right) \geq 1 - \lambda, \quad F_{SAx_n, Su} \left(\frac{\epsilon}{2} \right) \geq 1 - \lambda, \quad \forall n \geq N_0.$$

Now,

$$\begin{aligned} F_{ASx_n, Su}(\epsilon) &\geq \Delta \left(F_{ASx_n, SAx_n} \left(\frac{\epsilon}{2} \right), F_{SAx_n, Su} \left(\frac{\epsilon}{2} \right) \right) \\ &\geq \Delta(1-\lambda, 1-\lambda) \\ &\geq 1-\lambda; \quad \forall n \geq N_0. \end{aligned}$$

Hence $ASx_n \rightarrow Su$, i.e. (A, S) is semi-compatible.

Definition 8: Two self mappings A and S of a menger space (X, F, Δ) are said to be **reciprocally continuous** if

$$\lim_{n \rightarrow \infty} ASx_n = At \quad \text{and} \quad \lim_{n \rightarrow \infty} SAx_n = St,$$

whenever $\{x_n\}$ is a sequence in X such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Sx_n = t; \quad \text{for some } t \in X$$

If A and S are both continuous, then they are obviously reciprocally continuous but the converse is not true.

MAIN RESULTS

The following theorem is given by R. Tiwari and S.K. Shrivastava [5]

Theorem 9: Let T be a multi-valued mapping from a metric space X to $CL(X)$. If there exists $f: X \rightarrow X$ such that $TX \subseteq fX$, for each $x, y \in X$, and

$$H(Tx, Ty) \leq \phi \left(\alpha d(fx, fy) + \beta [D(fx, Tx) + D(fx, Ty)] + \gamma [D(fx, fy) + D(fy, Tx)] \right. \\ \left. \sigma \max \left\{ d(fx, fy), \frac{1}{2} [D(fx, Tx) + D(fy, Ty)], \frac{1}{2} [D(fx, Ty) + D(fy, Tx)] \right\} \right)$$

where $\max\{\alpha + 2\gamma + \sigma, \beta + \gamma + \sigma\} \leq 1$, $\alpha, \beta, \gamma \geq 0$, $0 < \sigma \leq 1$, $\phi(t) < qt$ for each $t > 0$ for some fixed $0 < q < 1$, $\phi \in \Psi$ and there exists an $x_0 \in X$ such that T is asymptotically regular at x_0 , and X is (T, f, x) -orbitally complete, then T and f have a coincidence point.

Taking the clue from above theorem 9, we prove the following theorem

Theorem 10: Let (X, F, Δ) be a Menger space, where $\Delta(a, b) = \min\{a, b\}$, for all $a, b \in [0, 1]$ and T be a multi-valued mapping from X to $CL(X)$. If there exists mapping $S: X \rightarrow X$ such that

(a) $TX \subseteq SX$, for each $x, y \in X$, and

$$(b) \quad F_{Tx,Ty}(\phi t) \geq \min \left(F_{Sx,Sy}(t), F_{Sx,Tx}(t), F_{Sy,Ty}(t), F_{Sx,Ty}(t), F_{Sy,Tx}(t), \right. \\ \left. \max \left\{ F_{Sx,Sy}(t), F_{Sx,Tx}(\alpha t), F_{Sy,Ty}((2-\alpha)t), F_{Sy,Ty}(\beta t), F_{Sy,Tx}((2-\beta)t) \right\} \right)$$

for all $t > 0, \alpha, \beta \in (0, 1)$,

(c) $\phi(t) < qt, \forall t > 0, 0 < q < 1, \phi \in \Phi$,

(d) There exists an $x_0 \in X$ such that T is asymptotically regular at x_0 ,

(e) (T, S, x) - orbitally complete.

Then T and S have a coincidence point.

Proof: Choose $x_0 \in X$ satisfying (a). We shall construct two sequences $\{x_n\}$ and $\{y_n\}$ as follows:

Since $TX \subset SX$, Choose $y_1 = Sx_1 \in Tx_0$. If $Tx_0 = Tx_1$, Choose $y_2 = Sx_2 \in Tx_1$ such that $y_1 = y_2$, If $Tx_0 \neq Tx_1$, from the definition of Hausdorff distance one can choose $y_2 = Sx_2 \in Tx_1$ such that

$$F_{y_1, y_2}(t) \geq F_{Tx_0, Tx_1}(t)$$

In general, Choose $y_{n+2} = Sx_{n+2} \in Tx_{n+1}$, such that $y_{n+1} = y_{n+2}$ if $Tx_n = Tx_{n+1}$ and $F_{y_{n+1}, y_{n+2}}(t) \geq F_{Tx_n, Tx_{n+1}}(t)$

Otherwise.

We wise to show that $\{y_n\}$ is cauchy. For this it is sufficient to show that $\{y_{2n}\}$ is cauchy.

Suppose on the contrary that $\{y_{2n}\}$ is not cauchy. Then there is an $\epsilon > 0$ such that for each integer $2k, k = 0, 1, 2, \dots$ there exists even integers $2nk$ and $2mk$ with $2k < 2nk < 2mk$ such that

$$F_{y_{2nk}, y_{2mk}}(\epsilon) < 1-\lambda. \tag{1}$$

Let for each integer $2k, 2mk$ be the least positive integer exceeding $2nk$ satisfying (1). Then,

$$F_{y_{2nk}, y_{2mk}}(\epsilon) \geq 1-\lambda \tag{2}$$

$$F_{y_{2nk}, y_{2mk}}(\epsilon) < 1-\lambda.$$

As such, for each even integer $2k$, we have

$$1-\lambda > F_{y_{2nk}, y_{2mk}}(\epsilon) \geq F_{y_{2nk}, y_{2mk}}(\epsilon) \geq \Delta \left(F_{y_{2nk}, y_{2mk-2}} \left(\frac{\epsilon}{3} \right), F_{y_{2mk-2}, y_{2mk-1}} \left(\frac{\epsilon}{3} \right), F_{y_{2mk-1}, y_{2mk}} \left(\frac{\epsilon}{3} \right) \right)$$

So by (2) and $k \rightarrow \infty$, we get

$$\lim_{n \rightarrow \infty} F_{y_{2nk}, y_{2mk}}(\epsilon) = 1-\lambda. \tag{3}$$

Now, using (3) in the triangle inequality

$$F_{y_{2nk}, y_{2mk-1}}(\epsilon) \geq \Delta \left(F_{y_{2nk}, y_{2mk}} \left(\frac{\epsilon}{2} \right), F_{y_{2mk}, y_{2mk-1}} \left(\frac{\epsilon}{2} \right) \right)$$

and

$$F_{y_{2nk+1}, y_{2mk-1}}(\epsilon) \geq \Delta \left(F_{y_{2nk+1}, y_{2nk}} \left(\frac{\epsilon}{3} \right), F_{y_{2nk}, y_{2mk}} \left(\frac{\epsilon}{3} \right), F_{y_{2mk}, y_{2mk-1}} \left(\frac{\epsilon}{3} \right) \right)$$

Taking $k \rightarrow \infty$

$$F_{y_{2nk+1}, y_{2mk-1}}(\epsilon) \geq \Delta [1-\lambda, 1] = 1-\lambda \text{ and} \tag{4}$$

$$F_{y_{2nk+1}, y_{2mk-1}}(\epsilon) \geq \Delta [1-\lambda, 1, 1] = 1-\lambda. \tag{5}$$

Then,

$$\begin{aligned}
 F_{y_{2nk}, y_{2mk}}(\phi t) &\geq \Delta \left(F_{y_{2nk}, y_{2nk+1}} \left(\frac{\phi t}{2} \right), F_{y_{2nk+1}, y_{2mk}} \left(\frac{\phi t}{2} \right) \right) \\
 &\geq \Delta \left(F_{y_{2nk}, y_{2nk+1}} \left(\frac{\phi t}{2} \right), F_{Tx_{2nk+1}, Tx_{2mk}} \left(\frac{\phi t}{2} \right) \right) \\
 &\geq \Delta \left(F_{y_{2nk}, y_{2nk+1}} \left(\frac{\phi t}{2} \right), \min \left(\begin{aligned} &F_{Sx_{2nk+1}, Sx_{2mk}}(2t), F_{Sx_{2nk+1}, Tx_{2nk+1}}(2t), F_{Sx_{2mk}, Tx_{2mk}}(2t), \\ &F_{Sx_{2nk+1}, Tx_{2mk}}(2t), F_{Sx_{2mk}, Tx_{2nk+1}}(2t), \\ &\max \left(F_{Sx_{2nk+1}, Sx_{2mk}}(2t), F_{Sx_{2nk+1}, Tx_{2nk+1}}(2(1-r)t), F_{Sx_{2nk+1}, Tx_{2nk+1}}(2(1+r)t), \right. \\ &\left. F_{Sx_{2nk+1}, Tx_{2mk}}(2(1-q)t), F_{Sx_{2mk}, Tx_{2nk+1}}(2(1+q)t) \right) \end{aligned} \right) \right)
 \end{aligned}$$

Putting $\beta = 1-q, \alpha = 1-r, q, r \in (0, 1)$

$$\geq \Delta \left(F_{y_{2nk}, y_{2nk+1}} \left(\frac{\phi t}{2} \right), \min \left(\begin{aligned} &F_{y_{2nk+1}, y_{2mk}}(2t), F_{y_{2nk+1}, y_{2nk+2}}(2t), F_{y_{2mk}, y_{2mk+1}}(2t), \\ &F_{y_{2nk+1}, y_{2mk+1}}(2t), F_{y_{2mk}, y_{2nk+2}}(2t), \\ &\max \left(F_{y_{2nk+1}, y_{2mk}}(2t), F_{y_{2nk+1}, y_{2nk+2}}(2(1-r)t), F_{y_{2nk+1}, y_{2nk+1}}(2(1+r)t), \right. \\ &\left. F_{y_{2nk+1}, y_{2mk+1}}(2(1-q)t), F_{y_{2mk}, y_{2nk+1}}(2(1+q)t) \right) \end{aligned} \right) \right)$$

Since ϕ is upper semi-continuous, taking the limit as $k \rightarrow \infty$

$$1-\lambda \geq \Delta(1, (1-\lambda), 1, 1, (1-\lambda), (1-\lambda), \max\{(1-\lambda), 1, 1, (1-\lambda), (1-\lambda)\})$$

$$1-\lambda \geq 1-\lambda,$$

which is a contraction.

Thus $\{y_n\}$ is a Cauchy sequence. Since SX is (T, S, x_0) -orbitally complete, $\{y_n\}$ converges to a point u in X . Hence there exists a point z in SX such that $u = Sz$. Then,

$$\begin{aligned}
 F_{S_z, T_z}(\phi t) &\geq \Delta \left(F_{S_z, S_{x_{n+1}}} \left(\frac{\phi t}{2} \right), F_{S_{x_{2n+1}}, T_z} \left(\frac{\phi t}{2} \right) \right) \\
 &\geq \Delta \left(F_{S_z, S_{x_{n+1}}} \left(\frac{\phi t}{2} \right), F_{Tx_n, T_z} \left(\frac{\phi t}{2} \right) \right) \\
 &\geq \Delta \left(F_{S_z, S_{x_{n+1}}} \left(\frac{\phi t}{2} \right), \min \left(\begin{aligned} &F_{S_{x_n}, S_z}(2t), F_{S_{x_n}, Tx_n}(2t), F_{S_z, T_z}(2t), \\ &F_{S_{x_n}, T_z}(2t), F_{S_z, Tx_n}(2t), \\ &\max \left(F_{S_{x_n}, S_z}(2t), F_{S_{x_n}, Tx_n}(2(1-r)t), F_{S_z, T_z}(2(1+r)t), \right. \\ &\left. F_{S_{x_n}, T_z}(2(1-q)t), F_{S_z, Tx_n}(2(1+q)t) \right) \end{aligned} \right) \right)
 \end{aligned}$$

Taking limit $n \rightarrow \infty$, we have

$$F_{S_z, T_z}(\phi t) \geq \Delta \left(1, \min \left(\begin{array}{l} 1, F_{S_z, T_z}(2t), F_{S_z, T_z}(2t), \\ F_{S_z, T_z}(2t), F_{S_z, T_z}(2t), \\ \max \left(\begin{array}{l} 1, F_{S_z, T_z}(2(1-r)t), F_{S_z, T_z}(2(1+r)t), \\ F_{S_z, T_z}(2(1-q)t), F_{S_z, T_z}(2(1+q)t) \end{array} \right) \end{array} \right) \right) \\ \geq \Delta \left(1, \min \left(1, F_{S_z, T_z}(2t), 1 \right) \right)$$

$$F_{S_z, T_z}(\phi t) \geq F_{S_z, T_z}(2t)$$

$$F_{S_z, T_z}(t) \geq F_{S_z, T_z}(\phi^{-1} t)$$

Hence $Sz \in Tz$.

z is a coincidence point of S and T .

APPLICATIONS

In this section we study the existence of fixed point for multi-valued and self-mappings in a metric space (X, d) using the results in main result.

Theorem 11: Let (X, d) be a complete metric space and $T: (X, d) \rightarrow (CL(X), d_H)$. If there exists a mapping $S: (X, d) \rightarrow (X, d)$ such that

- (a) $TX \subseteq SX$, for each $x, y \in X$, and
- (b) $d_H(Tx, Ty) \leq \phi \max \{d(Sx, Sy), d_H(Sx, Tx), d_H(Sy, Ty), d_H(Sx, Ty), d_H(Sy, Tx)\}, \min \{d(Sx, Sy), \frac{1}{2} [d_H(Sy, Ty) + d_H(Sx, Tx)], \frac{1}{2} [d_H(Sx, Ty) + d_H(Sy, Tx)]\}$,

where $\phi(t) < qt$ for each $t > 0, 0 < q > 1, \phi \in \Phi$ and

- (c) there exists an $x_0 \in X$ such that T is asymptotically regular at x_0 ,
- (d) X is (T, S, X) - orbitally complete.

Then T and S have a coincidence point.

Proof: If we define $F: X \times X \rightarrow D^+$ by $F_{A, B}(t) = H(t - d_H(A, B))$, where $A, B \in CL(X)$, then the space (X, F, \min) with a t -norm $\Delta = \min$ is a Menger space and the topology induced by the metric d coincides with the topology τ . And for any $Tx, Ty \in CL(X)$, we have

$$\begin{aligned} F_{Tx, Ty}(\phi t) &\geq H[\phi t - d_H(Tx, Ty)] \\ &\geq H[\phi t - \max \{d(Sx, Sy), d_H(Sx, Tx), d_H(Sx, Ty), \\ &\quad d_H(Sx, Ty), d_H(Sx, Tx)\}, \min \{d(Sx, Sy), \\ &\quad \frac{1}{2} [d_H(Sx, Tx) + d_H(Sx, Ty)], \frac{1}{2} [d_H(Sx, Ty) + d_H(Sy, Tx)]\}] \\ &= H[t - \max \{d_1, d_2, d_3, d_4, d_5, \min \{d_6, \frac{1}{2} (d_7 + d_8), \frac{1}{2} (d_9 + d_{10})\}\}] \end{aligned}$$

where, $d_1 = d(Sx, Sy), d_2 = d_H(Sx, Tx), d_3 = d_H(Sx, Ty), d_4 = d_H(Sx, Ty),$

$d_5 = d_H(Sx, Tx), d_6 = d(Sx, Sy), d_7 = d_H(Sx, Tx), d_8 = d_H(Sx, Ty),$

$$\begin{aligned}
 d_9 &= d_H(Sx, Ty), d_{10} = d_H(Sy, Tx) \\
 &= H[\min\{(t - d_1), (t - d_2), (t - d_3), (t - d_4), (t - d_5), \\
 &\quad \max\{(t - d_6), (t - \frac{1}{2}(d_7 - d_8)), (t - \frac{1}{2}(d_9 + d_{10}))\}\} \\
 &= \min\{H(t - d_1), H(t - d_2), H(t - d_3), H(t - d_4), H(t - d_5), \\
 &\quad \max\{H(t - d_6), H(\alpha t - d_7), H((2 - \alpha)t - d_8), H(\beta t - d_9), \\
 &\quad H((2 - \beta)t - d_{10})\}\} \text{ for some } \alpha, \beta \in (0, 2) \\
 &= \min\{F_{Sx, Sy}(t), F_{Sx, Tx}(t), F_{Sy, Ty}(t), F_{Sx, Ty}(t), F_{Sy, Tx}(t), \\
 &\quad \max\{F_{Sx, Sy}(t), F_{Sx, Tx}(\alpha t), F_{Sy, Ty}((2 - \alpha)t), F_{Sx, Ty}(\beta t), F_{Sy, Tx}((2 - \beta)t)\}
 \end{aligned}$$

Thus Theorem 10 follows from Theorem 11 immediately.

Hence there exists a coincident point.

REFERENCES

- [1] Mukherjee, R.N.: *Indian J. pure Appl. Math.* 12(8) (1981), 930.
- [2] Singh, Y.R.: Studies on fixed point, common fixed points and coincidence, *Doctoral Thesis, Manipur University (Kanchipur)*, (2002).
- [3] Som, T.: Few common fixed points for comparative mappings, *Bull. Cal. Math. Soc.*, 95(4) (2003), 307-312.
- [4] Taskovik, M.R.: Some results in fixed point theory, *Publ. L's. Inst. Math. (Beograd) N.S.*, 20 (1976), 2311-242.
- [5] Tiwari, R. and Shrivastava, S.K.: Fixed point theorems and coincidence point, *South East Asian J. Math. & Math. Sc.* 5(1) (2006), 91-96.

Source of Support: Nil, Conflict of interest: None Declared

[Copy right © 2014 This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]