



The Algorithm for Inverse Matrices of Symmetric r-Circulant Matrices over Skew Field

Song Zhenzhen* and Wang Junqing

Department of Mathematics, Tianjin Polytechnic University, Tianjin 300387, PR China.

(Received On: 25-03-15; Revised & Accepted On: 13-04-15)

ABSTRACT

This paper mainly introduces elementary column transformation method of inverse matrix of symmetric r-circulant matrix over skew field by using the elementary column transformation of polynomial matrix.

Keywords: skew field; symmetric r-circulant matrix; elementary column transformation.

1. INTRODUCTION

Symmetric r-circulant matrix is an important special matrix. Therefore, the algorithms for inverse matrix of the matrix over skew field cause research of a lot of mathematicians. In this paper, using the elementary column transformation of polynomial matrix is given elementary column transformation method of inverse matrix of symmetric r-circulant matrix. And the algorithm has smaller amount of computations.

2. PREPARATION KNOWLEDGE

In this paper, let K be a skew field and x be unknown, and its primitive formula

$$f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n \text{ where } a_0, a_1, \dots, a_n \in K$$

is defined to be unary polynomial over skew field K , simply indicating $f(x) = \sum_{t=0}^n a_t x^t$. $K[x]$ is a ring, and it is

defined to be monadic polynomial ring over skew field K . $K^{m \times n}$ is denoted $m \times n$ -factorial-matrix. $d(x) = (f(x), g(x))$ of leading coefficient being 1 are respectively denoted the greatest left common divisor $f(x)$ and $g(x)$.

Definition 2.1: Let $r \in K$, matrix

$$A = \begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_{n-2} & a_{n-1} \\ ra_{n-1} & a_0 & a_1 & \dots & a_{n-3} & a_{n-2} \\ ra_{n-2} & ra_{n-1} & a_0 & \dots & a_{n-4} & a_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ ra_1 & ra_2 & ra_3 & \dots & ra_{n-1} & a_0 \end{pmatrix}$$

is called r-circulant matrix. Referred to as $A = K_r(a_0, a_2, \dots, a_{n-1}) \in KM_r$.

Definition 2.2: Let $r \in K$, matrix

$$A = \begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_{n-2} & a_{n-1} \\ a_1 & a_2 & a_3 & \dots & a_{n-1} & ra_0 \\ a_2 & a_3 & a_4 & \dots & ra_0 & ra_1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{n-1} & ra_0 & ra_1 & \dots & ra_{n-3} & ra_{n-2} \end{pmatrix}$$

is called symmetric r-circulant matrix. Referred to as $A = SK_r(a_0, a_2, \dots, a_n) \in SKM_r$.

Definition 2.3: Introducing n order identity matrix $J_n = (e_n, e_{n-1}, \dots, e_1)$ and e_i being n order unit column vector of the first i component for 1 and the other component for 0, then AJ_n is called r -circulant matrix.

3. ALGORITHM

Lemma 3.1: Set $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r$, $B = SK_r(a_{n-1}, a_{n-2}, \dots, a_0) \in SKM_r$, then $BP = A$ and $B = AP$, $P = SK_1(0, \dots, 0, 1) \in SKM_1$.

Lemma 3.2: Set $A \in K^{m \times n}$, $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r \Leftrightarrow A = f(J)$, $f(x) = \sum_{i=0}^{n-1} a_i x^i$.

Lemma 3.3: Set $A \in K^{m \times n}$, $A = SK_r(a_0, a_1, \dots, a_{n-1}) \in SKM_r \Leftrightarrow A = \left(\sum_{i=1}^n a_{n-i} J^{i-1} \right) P$.

Lemma 3.4: Set $A = SK_r(a_0, a_1, \dots, a_{n-1}) \in SKM_r$, $B = K_r(a_{n-1}, a_{n-2}, \dots, a_0) \in KM_r (r \neq 0)$, then A is non singular $\Leftrightarrow B$ is non singular.

Lemma 3.5: Set $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r (r \neq 0)$ be non singular, then $A^{-1} \in KM_r$.

Lemma 3.6: Set $A = SK_r(a_0, a_1, \dots, a_{n-1}) \in SKM_r (r \neq 0)$ be non singular, then $A^{-1} \in SKM_r$.

Lemma 3.7: Set $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r (r \neq 0)$, then A is non singular $\Leftrightarrow (f(x), g(x)) = 1$.

Lemma 3.8: Set $A = SK_r(a_0, a_1, \dots, a_{n-1}) \in SKM_r (r \neq 0)$, then A is non singular $\Leftrightarrow (f(x), g(x)) = 1$.

Theorem 3.1: Set $f(x), g(x)$ be nonzero polynomial of $K[x]$, If $C = (f(x), g(x))$, by a series of the elementary column transformation of C , we have $(d(x), 0)$. Then $(f(x), g(x)) = d(x)$, $d(x) = u(x)f(x) + v(x)g(x)$.

Theorem 3.2: Set $f(x), g(x)$ be nonzero polynomial of $K[x]$. If $C = \begin{pmatrix} f(x) & g(x) \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$, by a series of the elementary

column transformation of C , we have $\begin{pmatrix} d(x) & 0 \\ u(x) & s(x) \\ v(x) & t(x) \end{pmatrix}$. Then $(f(x), g(x)) = d(x)$, $d(x) = u(x)f(x) + v(x)g(x)$.

Corollary 3.1: Set $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r (r \neq 0)$ and non singular, then we have polynomial $u(x) = \sum_{i=0}^{n-1} b_i x^i$, and $A^{-1} = u(J)$.

Proof: Because $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r (r \neq 0)$ and non singular, according to lemma 3.7 we have $(f(x), g(x)) = 1$.

By a series of the elementary column transformation for polynomial matrix $C = \begin{pmatrix} f(x) & g(x) \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$, we get

$$\begin{pmatrix} 1 & 0 \\ u(x) & s(x) \\ v(x) & t(x) \end{pmatrix} \cdot \text{And } (f(x), g(x)) \begin{pmatrix} u(x) & s(x) \\ v(x) & t(x) \end{pmatrix} = (1, 0),$$

$A^{-1} = u(J) = \sum_{i=0}^{n-1} b_i J^i$, $u(x) = \sum_{i=0}^{n-1} b_i x^i$. So we have $u(x)f(x) + v(x)g(x) = 1$, if $x = J$, then $u(J)f(J) + v(J)g(J) = E$.

Corollary 3.2: Set $B = SK_r(a_0, a_1, \dots, a_{n-1}) \in SKM_r$, $A = K_r(a_{n-1}, a_{n-2}, \dots, a_0) \in KM_r (r \neq 0)$, and $B^{-1} = P(u(J))$.

Proof: According to lemma 3.1 $B = AP$ and $P^{-1} = P$, so $B^{-1} = (AP)^{-1} = P^{-1}A^{-1} = PA^{-1} = P(u(J))$.

According to corollary 3.1 and corollary 3.2 we get elementary transformation algorithm. Follow the steps outlined below :

Step-1: According to $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r (r \neq 0)$, we get $f(x), g(x)$.

Step-2: By a series of the elementary column transformation for $C = \begin{pmatrix} f(x) & g(x) \\ 1 & 0 \end{pmatrix}$, we get $\begin{pmatrix} d(x) & u(x) \\ 0 & s(x) \end{pmatrix}$.

Step-3: If $d(x) = 1$, then $A = K_r(a_0, a_1, \dots, a_{n-1}) \in KM_r (r \neq 0)$ and non singular. We get $A^{-1} = u(J)$.

Step-4: We get $B = SK_r(a_0, a_1, \dots, a_{n-1}) \in SKM_r$ and non singular and $A^{-1} = u(J)$, so we get $B^{-1} = P(u(J))$.

REFERENCE

1. Wajin Zhuang, Introduction to Matrix Theory over Skew Field [M], Beijing: Science press, 2006.
2. Weiwei Xu. The Inverse Matrix of the Special Binary Symmetry Cyclic Matrix [J], Journal of Yunnan University, 2006, 28 (S1): 45-49.
3. Jingliang Chen, Xianghui Chen. Special Matrices [M], Beijing: Qinghua University Press, 2001.
4. Quan Lu, Zhong Xu, Fang Chen, Zhijie Yuan. A Fast Algorithm for the Triangular Factorization of Symmetric r-circulant Matrix and Inverse Matrix[J], Mathematics in Practice and Theory, 2006, 36(5): 212-217.
5. Jiaqing jiang. On a Fast and Convenient Algorithm for the Inverse of Two Types of Cyclic Matrix [J], Journal of Qufu Normal University, 2011, 37(4): 45-48.

Corresponding author: Song Zhenzhen
Department of Mathematics, Tianjin Polytechnic University, Tianjin 300387, PR China.

Source of Support: Nil, Conflict of interest: None Declared

[Copy right © 2015, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]