



COMMON FIXED POINT THEOREMS
FOR WEAKLY COMPATIBLE MAPPINGS IN G-METRIC SPACES

SMITA NAIR*, SHALU SAXENA

Department of Mathematics, Sri Sathya Sai College for Women, Bhopal – (M.P.), India.

(Received On: 17-02-16; Revised & Accepted On: 10-03-16)

ABSTRACT

In this paper we prove common fixed theorem using the concept of weakly compatible mappings. The results can be considered as an extension of theorems to G-metric space.

Key Words: Weakly compatible mapping, Complete G-metric space.

INTRODUCTION

The study of common fixed points of mappings satisfying various contractive conditions has been the centre of rigorous research activity. The concept of commuting maps was replaced by weakly commuting maps introduced by Sessa [4]. The concept of weakly commuting maps was further generalized by Jungck who introduced the concept of compatible mappings [1, 2, 3] and then weakly compatible mappings. Common fixed point theorems have been proved using these concepts in metric space, 2 metric space. Sims [5, 6] have shown that most of the results related to D-metric spaces are invalid. Therefore they introduced the concept of G-metric space. The purpose of this paper is to obtain common fixed point theorem for mappings satisfying condition of weak compatibility

Theorem: Let (X, G) be a complete G-metric Space. Let f and g be self mappings of X satisfying the following conditions

- (i) $f(X) \subseteq g(X)$, $f(X)$ is closed subset of $g(X)$
- (ii) f and g are compatible mappings
- (iii) $G(fx, fy, fz) \leq \alpha G(gx, gy, gz) + \beta G(gx, fx, fx) + \gamma G(gy, fy, fy) + \delta G(gz, fz, fz) + \eta \max \{G(gx, gy, gz), G(gx, fx, fx), G(gx, fy, fy), G(gy, fy, fy), G(gy, fx, fx), G(gy, fz, fz), G(gz, fz, fz), G(gz, fx, fx), G(gz, fy, fy)\}$ for all $x, y, z, \in X$, where $\alpha + \beta + \gamma + \delta + \eta \leq 1/2$

Then f and g have a unique common fixed point in X .

Proof: Let x_0 be an arbitrary point in X . By (1) we can choose a point x_1 in X such that $fx_0 = gx_1$, In general we can choose x_{n+1} such that

$$y_n = fx_n = gx_{n+1}, n = 0, 1, 2, \dots$$

From (3) we can write

$$G(fx_n, fx_{n+1}, fx_{n+2}) \leq \alpha G(gx_n, gx_{n+1}, gx_{n+2}) + \beta G(gx_n, fx_n, fx_n) + \gamma G(gx_{n+1}, fx_{n+1}, fx_{n+1}) + \delta G(gx_{n+2}, fx_{n+2}, fx_{n+2}) + \eta \max \{G(gx_n, gx_{n+1}, gx_{n+2}), G(gx_n, fx_n, fx_n), G(gx_n, fx_{n+1}, fx_{n+1}), G(gx_{n+1}, fx_{n+1}, fx_{n+1}), G(gx_{n+1}, fx_n, fx_n), G(gx_{n+1}, fx_{n+2}, fx_{n+2}), G(gx_{n+2}, fx_{n+2}, fx_{n+2}), G(gx_{n+2}, fx_n, fx_n), G(gx_{n+2}, fx_{n+1}, fx_{n+1})\}$$

Or

$$G(y_n, y_{n+1}, y_{n+2}) \leq \alpha G(y_{n-1}, y_n, y_{n+1}) + \beta G(y_{n-1}, y_n, y_n) + \gamma G(y_n, y_{n+1}, y_{n+1}) + \delta G(y_{n+1}, y_{n+2}, y_{n+2}) + \eta \max \{G(y_{n-1}, y_n, y_{n+1}), G(y_{n-1}, y_n, y_n), G(y_{n-1}, y_{n+1}, y_{n+1}), G(y_n, y_{n+1}, y_{n+1}), G(y_n, y_n, y_n), G(y_n, y_{n+1}, y_{n+1}), G(y_{n+1}, y_{n+2}, y_{n+2}), G(y_{n+1}, y_n, y_n), G(y_{n+1}, y_{n+1}, y_{n+1})\}$$

$$\leq \alpha G(y_{n-1}, y_n, y_{n+1}) + 2\beta G(y_{n-1}, y_n, y_{n+1}) + 2\gamma G(y_n, y_{n+1}, y_{n+2}) + 2\delta G(y_n, y_{n+1}, y_{n+2}) + \eta \max \{G(y_{n-1}, y_n, y_{n+1}), 2G(y_{n-1}, y_n, y_{n+1}), 2G(y_{n-1}, y_n, y_{n+1}), 2G(y_n, y_{n+1}, y_{n+2}), 2G(y_n, y_{n+1}, y_{n+2}), 2G(y_n, y_{n+1}, y_{n+2}), 2G(y_n, y_{n+1}, y_{n+2})\}$$

Corresponding Author: Smita Nair*

Department of Mathematics, Sri Sathya Sai College for Women, Bhopal – (M.P.), India.

$$\begin{aligned} &\leq \alpha G(y_{n-1}, y_n, y_{n+1}) + 2\beta G(y_{n-1}, y_n, y_{n+1}) + 2\gamma G(y_n, y_{n+1}, y_{n+2}) + 2\delta G(y_n, y_{n+1}, y_{n+2}) \\ &\quad + \eta \max \{ 2 G(y_{n-1}, y_n, y_{n+1}), 2 G(y_n, y_{n+1}, y_{n+2}) \} \\ &\leq (\alpha + 2\beta) G(y_{n-1}, y_n, y_{n+1}) + (2\gamma + 2\delta) G(y_n, y_{n+1}, y_{n+2}) + \eta \max \{ 2 G(y_{n-1}, y_n, y_{n+1}), \\ &\quad 2 G(y_n, y_{n+1}, y_{n+1}) \} \end{aligned}$$

We have following cases

Case -I: $\max = 2 G(y_{n-1}, y_n, y_{n+1})$

$$G(y_n, y_{n+1}, y_{n+2}) \leq (2\eta + \alpha + 2\beta) G(y_{n-1}, y_n, y_{n+1}) + (2\gamma + 2\delta) G(y_n, y_{n+1}, y_{n+2})$$

$$G(y_n, y_{n+1}, y_{n+2}) \leq \left[\frac{2\eta + \alpha + 2\beta}{1 - 2\gamma - 2\delta} \right] G(y_{n-1}, y_n, y_{n+1})$$

$$G(y_n, y_{n+1}, y_{n+2}) \leq G(y_{n-1}, y_n, y_{n+1}) \text{ as } \alpha + \beta + \delta + \gamma + \eta \leq 1/2$$

Case- II: $\max = 2 G(y_n, y_{n+1}, y_{n+2})$

$$G(y_n, y_{n+1}, y_{n+2}) \leq (\alpha + 2\beta) G(y_{n-1}, y_n, y_{n+1}) + (2\gamma + 2\delta + 2\eta) G(y_n, y_{n+1}, y_{n+2})$$

$$G(y_n, y_{n+1}, y_{n+2}) \leq \left[\frac{\alpha + 2\beta}{1 - 2\gamma - 2\delta - 2\eta} \right] G(y_{n-1}, y_n, y_{n+1})$$

$$G(y_n, y_{n+1}, y_{n+2}) \leq G(y_{n-1}, y_n, y_{n+1}) \text{ as } \alpha + \beta + \delta + \gamma + \eta \leq 1/2$$

Similarly now we find

$$\begin{aligned} G(fx_{n+1}, fx_{n+2}, fx_{n+3}) &\leq \alpha G(gx_{n+1}, gx_{n+2}, gx_{n+3}) + \beta G(gx_{n+1}, fx_{n+1}, fx_{n+1}) + \gamma G(gx_{n+2}, fx_{n+2}, fx_{n+2}) \\ &\quad + \delta G(gx_{n+3}, fx_{n+3}, fx_{n+3}) + \eta \max \{ G(gx_{n+1}, gx_{n+2}, gx_{n+3}), G(gx_{n+1}, fx_{n+1}, fx_{n+1}), \\ &\quad G(gx_{n+1}, fx_{n+2}, fx_{n+2}), G(gx_{n+2}, fx_{n+2}, fx_{n+2}), G(gx_{n+2}, fx_{n+1}, fx_{n+1}), G(gx_{n+2}, fx_{n+3}, fx_{n+3}), \\ &\quad G(gx_{n+3}, fx_{n+3}, fx_{n+3}), G(gx_{n+3}, fx_{n+1}, fx_{n+1}), G(gx_{n+3}, fx_{n+2}, fx_{n+2}) \} \end{aligned}$$

Or

$$\begin{aligned} G(y_{n+1}, y_{n+2}, y_{n+3}) &\leq \alpha G(y_n, y_{n+1}, y_{n+2}) + \beta G(y_n, y_{n+1}, y_{n+1}) + \gamma G(y_{n+1}, y_{n+2}, y_{n+2}) + \delta G(y_{n+2}, y_{n+3}, y_{n+3}) \\ &\quad + \eta \max \{ G(y_n, y_{n+1}, y_{n+2}), G(y_n, y_{n+1}, y_{n+1}), G(y_n, y_{n+2}, y_{n+2}), G(y_{n+1}, y_{n+2}, y_{n+2}), \\ &\quad G(y_{n+1}, y_{n+1}, y_{n+1}), G(y_{n+1}, y_{n+3}, y_{n+3}), G(y_{n+2}, y_{n+3}, y_{n+3}), G(y_{n+2}, y_{n+1}, y_{n+1}), G(y_{n+2}, y_{n+2}, y_{n+2}) \} \end{aligned}$$

Or

$$\begin{aligned} G(y_{n+1}, y_{n+2}, y_{n+3}) &\leq \alpha G(y_n, y_{n+1}, y_{n+2}) + 2\beta G(y_n, y_{n+1}, y_{n+2}) + 2\gamma G(y_{n+1}, y_{n+2}, y_{n+3}) + 2\delta G(y_{n+1}, y_{n+2}, y_{n+3}) \\ &\quad + \eta \max \{ G(y_n, y_{n+1}, y_{n+2}), 2G(y_n, y_{n+1}, y_{n+2}), 2 G(y_n, y_{n+1}, y_{n+2}), 2G(y_{n+1}, y_{n+2}, y_{n+3}), \\ &\quad 2G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+1}, y_{n+2}, y_{n+3}) \} \end{aligned}$$

Or

$$\begin{aligned} &\leq \alpha G(y_n, y_{n+1}, y_{n+2}) + 2\beta G(y_n, y_{n+1}, y_{n+2}) + 2\gamma G(y_{n+1}, y_{n+2}, y_{n+3}) + 2\delta G(y_{n+1}, y_{n+2}, y_{n+3}) \\ &\quad + \eta \max \{ 2 G(y_n, y_{n+1}, y_{n+2}), 2G(y_{n+1}, y_{n+2}, y_{n+3}), \end{aligned}$$

$$\begin{aligned} &\leq (\alpha + 2\beta) G(y_n, y_{n+1}, y_{n+2}) + (2\gamma + 2\delta) G(y_{n+1}, y_{n+2}, y_{n+3}) + \eta \max \{ 2G(y_n, y_{n+1}, y_{n+2}), \\ &\quad 2G(y_{n+1}, y_{n+2}, y_{n+3}) \} \end{aligned}$$

We have following cases:

Case- I: $\max = 2 G(y_n, y_{n+1}, y_{n+2})$

$$G(y_{n+1}, y_{n+2}, y_{n+3}) \leq (\alpha + 2\beta + 2\eta) G(y_n, y_{n+1}, y_{n+2}) + (2\gamma + 2\delta) G(y_{n+1}, y_{n+2}, y_{n+3})$$

$$G(y_{n+1}, y_{n+2}, y_{n+3}) \leq \left(\frac{\alpha + 2\beta + 2\eta}{1 - 2\gamma - 2\delta} \right) G(y_n, y_{n+1}, y_{n+2})$$

$$G(y_{n+1}, y_{n+2}, y_{n+3}) \leq G(y_n, y_{n+1}, y_{n+2}) \text{ as } \alpha + \beta + \gamma + \delta + \eta \leq 1/2$$

Case- II: $\max = 2 G(y_{n+1}, y_{n+2}, y_{n+3}),$

$$G(y_{n+1}, y_{n+2}, y_{n+3}) \leq (\alpha + 2\beta) G(y_n, y_{n+1}, y_{n+2}) + (2\gamma + 2\delta + 2\eta) G(y_{n+1}, y_{n+2}, y_{n+3})$$

$$G(y_{n+1}, y_{n+2}, y_{n+3}) \leq G(y_n, y_{n+1}, y_{n+2}) \text{ as } \alpha + \beta + \gamma + \delta + \eta \leq 1/2$$

Similarly now we find

$$G(fx_{n+2}, fx_{n+3}, fx_{n+4}) \leq \alpha G(gx_{n+2}, gx_{n+3}, gx_{n+4}) + \beta G(gx_{n+2}, fx_{n+2}, fx_{n+2}) + \gamma G(gx_{n+3}, fx_{n+3}, fx_{n+3}) + \delta G(gx_{n+4}, fx_{n+4}, fx_{n+4}) + \eta \max \{G(gx_{n+2}, gx_{n+3}, gx_{n+4}), G(gx_{n+2}, fx_{n+2}, fx_{n+2}), G(gx_{n+2}, fx_{n+3}, fx_{n+3}), G(gx_{n+3}, fx_{n+2}, fx_{n+2}), G(gx_{n+3}, fx_{n+4}, fx_{n+4}), G(gx_{n+4}, fx_{n+2}, fx_{n+2}), G(gx_{n+4}, fx_{n+3}, fx_{n+3})\}$$

Or

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq \alpha G(y_{n+1}, y_{n+2}, y_{n+3}) + \beta G(y_{n+1}, y_{n+2}, y_{n+2}) + \gamma G(y_{n+2}, y_{n+3}, y_{n+3}) + \delta G(y_{n+3}, y_{n+4}, y_{n+4}) + \eta \max \{G(y_{n+1}, y_{n+2}, y_{n+3}), G(y_{n+1}, y_{n+2}, y_{n+2}), G(y_{n+1}, y_{n+3}, y_{n+3}), G(y_{n+2}, y_{n+2}, y_{n+2}), G(y_{n+2}, y_{n+4}, y_{n+4}), G(y_{n+3}, y_{n+2}, y_{n+2}), G(y_{n+3}, y_{n+3}, y_{n+3})\}$$

Or

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq \alpha G(y_{n+1}, y_{n+2}, y_{n+3}) + 2\beta G(y_{n+1}, y_{n+2}, y_{n+3}) + 2\gamma G(y_{n+2}, y_{n+3}, y_{n+4}) + 2\delta G(y_{n+2}, y_{n+3}, y_{n+4}) + \eta \max \{G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+2}, y_{n+3}, y_{n+4}), 2G(y_{n+2}, y_{n+3}, y_{n+4})\}$$

Or

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq \alpha G(y_{n+1}, y_{n+2}, y_{n+3}) + 2\beta G(y_{n+1}, y_{n+2}, y_{n+3}) + 2\gamma G(y_{n+2}, y_{n+3}, y_{n+4}) + 2\delta G(y_{n+2}, y_{n+3}, y_{n+4}) + \eta \max \{2 G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+2}, y_{n+3}, y_{n+4})\} \\ \leq (\alpha+2\beta) G(y_{n+1}, y_{n+2}, y_{n+3}) + (2\gamma+2\delta) G(y_{n+2}, y_{n+3}, y_{n+4}) + \eta \max \{2G(y_{n+1}, y_{n+2}, y_{n+3}), 2G(y_{n+2}, y_{n+3}, y_{n+4})\}$$

We have following two cases:

Case- I: $\max = 2 G(y_{n+1}, y_{n+2}, y_{n+3}), G(y_{n+2}, y_{n+3}, y_{n+4}) \leq (\alpha+2\beta+2\eta) G(y_{n+1}, y_{n+2}, y_{n+3}) + (2\gamma + 2\delta) G(y_{n+2}, y_{n+3}, y_{n+4})$

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq \left(\frac{\alpha + 2\beta + 2\eta}{1 - 2\gamma - 2\delta} \right) G(y_{n+1}, y_{n+2}, y_{n+3})$$

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq G(y_{n+1}, y_{n+2}, y_{n+3}) \text{ as } \alpha + \beta + \gamma + \delta + \eta \leq 1/2$$

Case-II: $\max = 2 G(y_{n+2}, y_{n+3}, y_{n+4}),$

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq (\alpha + 2\beta) G(y_{n+1}, y_{n+2}, y_{n+3}) + (2\gamma + 2\delta + 2\eta) G(y_{n+2}, y_{n+3}, y_{n+4})$$

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq \left(\frac{\alpha + 2\beta}{1 - 2\gamma - 2\delta - 2\eta} \right) G(y_{n+1}, y_{n+2}, y_{n+3})$$

$$G(y_{n+2}, y_{n+3}, y_{n+4}) \leq G(y_{n+1}, y_{n+2}, y_{n+3}) \text{ as } \alpha + \beta + \gamma + \delta + \eta \leq 1/2$$

$$G(y_n, y_n, y_m) \leq G(y_n, y_n, y_{n+1}) + G(y_{n+1}, y_{n+1}, y_{n+2}) + \dots + G(y_{m-1}, y_{m-1}, y_m) \\ \leq p^n G(y_0, y_1, y_2) + p^{n+1} G(y_0, y_1, y_2) + \dots + p^{m-1} G(y_0, y_1, y_2) \\ \leq (p^n + p^{n+1} + \dots + p^{m-1}) G(y_0, y_1, y_2) \\ \leq \left[\frac{p^n}{1 - p} \right] G(y_0, y_1, y_2) \rightarrow 0 \text{ as } n \rightarrow \infty, p = \alpha + \beta + \gamma + \eta$$

Subsequence $\{fx_n\}, \{gx_{n+1}\}$ converge to y . As $f(X)$ is closed subset of $g(X)$ Then there exist $u \in X$ such that $gu = y$

Now we shall prove that $fu = y$. So on the contrary Let $fu \neq y$.

Then consider on.

$$G(fu, fx_n, fx_n) \leq \alpha G(gu, gx_n, gx_n) + \beta G(gu, fu, fu) + \gamma G(gx_n, fx_n, fx_n) + \delta G(gx_n, fx_n, fx_n) + \eta \max \{G(gu, gx_n, gx_n), G(gu, fu, gu), G(gu, fx_n, fx_n), G(gx_n, fx_n, fx_n), G(gx_n, fu, fu), G(gx_n, fx_n, fx_n), G(gx_n, fx_n, fx_n), G(gx_n, fu, fu), G(gx_n, fx_n, fx_n)\} \\ \leq \alpha G(y, y, y) + \beta G(y, fu, y) + \gamma G(y, y, y) + \delta G(y, y, y) + \eta \max \{G(y, y, y), G(y, fu, y), G(y, y, y), G(y, y, y), G(y, fu, fu), G(y, y, y)\} \\ = \beta G(y, fu, fu) + \eta \max \{G(fu, y, y), G(y, fu, fu)\} \\ = \beta G(fu, y, y) + 2\eta G(fu, y, y) \\ = (\beta + 2\eta) G(fu, y, y)$$

Hence we get

$$(1 - \beta - 2\eta) G(fu, y, y) \leq 0$$

$$\Rightarrow fu = y$$

$\Rightarrow fu = y = gu$. As (f, g) are weakly compatible.

$$\Rightarrow gfu = fg u$$

$$\Rightarrow g y = fy$$

Now we shall prove that $fy = y$. So on the contrary Let $fy \neq y$. So we can write

$$\begin{aligned} G(fgx_n, fx_n, fx_n) &\leq \alpha G(ggx_n, gx_n, gx_n) + \beta G(ggx_n, fgx_n, fgx_n) + \gamma G(gx_n, fx_n, fx_n) + \delta G(gx_n, fx_n, fx_n) \\ &\quad + \eta \max\{G(ggx_n, gx_n, gx_n), G(ggx_n, fgx_n, fgx_n), G(ggx_n, fx_n, fx_n), G(gx_n, fx_n, fx_n), G(gx_n, fgx_n, fgx_n), \\ &\quad G(gx_n, fx_n, fx_n), G(gx_n, fx_n, fx_n), G(gx_n, fgx_n, fgx_n), G(gx_n, fx_n, fx_n)\} \\ &\leq \alpha G(gy, y, y) + \beta G(gy, fy, fy) + \gamma G(y, y, y) + \delta G(y, y, y) + \eta \max\{G(gy, y, y), \\ &\quad G(gy, fy, fy), G(gy, y, y), G(y, y, y), G(y, fy, fy), G(y, y, y), G(y, y, y), G(y, fy, fy), G(y, y, y)\} \\ &\leq \alpha G(fy, y, y) + \eta \max\{G(fy, y, y), G(fy, fy, fy), G(fy, y, y), G(y, fy, fy), G(y, fy, fy)\} \\ &\leq \alpha G(fy, y, y) + 2\eta G(fy, y, y) \\ &= (\alpha + 2\eta) G(fy, y, y) \end{aligned}$$

Or

$$(1 - \alpha - 2\eta) G(fy, y, y) \leq 0$$

$$\Rightarrow fy = y = gy$$

Uniqueness:-To prove this part we assume $y_1 \neq y$ is another fixed point of f and g .

Then consider on

$$G(y_1, y, y) = G(fy_1, fy, fy)$$

$$\begin{aligned} &\leq \alpha G(gy_1, gy, gy) + \beta G(gy, fy, fy) + \gamma G(gy, fy, fy) + \delta G(gy, fy, fy) + \eta \max\{G(gy_1, gy, gy), \\ &\quad G(gy, fy_1, fy_1), G(gy_1, fy, fy), G(gy, fy, fy), G(gy, fy_1, fy_1), G(gy, fy, fy), G(gy, fy, fy) G(gy, fy_1, fy_1), \\ &\quad G(gy, fy, fy)\} \end{aligned}$$

Or

$$G(y_1, y, y) \leq \alpha G(y_1, y, y) + \beta G(y_1, y_1, y_1) + \gamma G(y, y, y) + \delta G(y, y, y) + \eta \max\{G(y_1, y, y), G(y, y_1, y) G(y_1, y, y), G(y, y, y), G(y, y_1, y), G(y, y, y), G(y, y, y), G(y, y_1, y), G(y, y, y)\}$$

Or

$$\begin{aligned} G(y_1, y, y) &\leq \alpha G(y_1, y, y) + \eta \max\{G(y_1, y, y), G(y, y_1, y_1)\} \\ &\leq \alpha G(y_1, y, y) + \eta \max\{G(y_1, y, y), G(y, y_1, y) + G(y_1, y, y)\} \\ &= \alpha G(y_1, y, y) + \eta \max\{G(y_1, y, y), 2 G(y_1, y, y)\} \\ &= (\alpha + 2\eta) G(y_1, y, y) \end{aligned}$$

Or

$$(1 - \alpha - 2\eta) G(y_1, y, y) \leq 0$$

$$\Rightarrow G(y_1, y, y) = 0$$

$$\Rightarrow y_1 = y$$

Theorem 2: Let A, B, C, S, R and T be self mappings of a complete G -metric space (X, G) and

- (1) $A(X) \subseteq T(X), B(X) \subseteq S(X), C(X) \subseteq R(X)$ and $A(X)$ or $B(X)$ or $C(X)$ is a closed subset of X .
- (2) $G(Ax, By, Cz) \leq a G(By, Ax, Rx) + b G(Cz, By, Ty) + c G(Ax, Cz, Sz) + d \max\{G(Cz, By, Ty) + G(Ty, Ax, Rx) + G(Ax, Cz, Sz), G(Sz, By, Ty) + G(By, Ax, Rx), G(Rx, Cz, Sz)\}$ where $a, b, c, d \geq 0$ and $a + b + 2c + 3d < 1$.
- (3) The pairs $(A, R), (B, T)$ and (C, S) are weakly compatible pairs. Then the mappings A, B, C, S, T and R have a common fixed point in X .

Proof: Let $x_0 \in X$ be n arbitrary point By (1) there exist $x_1, x_2, x_3 \in X$ such that $Ax_0 = Tx_1 = y_0, Bx_1 = Sx_2 = y_1$ and $Cx_2 = Rx_3 = y_2$ Inductively we get a sequence $\{y_n\}$ in X such that
 $Ax_n = Tx_{n+1} = y_n, Bx_{n+1} = Sx_{n+2} = y_{n+1}, Cx_{n+2} = Rx_{n+3} = y_{n+2},$ for $n = 0, 1, 2, 3 \dots \{y_n\}$

Now we shall prove that the sequence is a Cauchy sequence. Let

$$d_n = G(y_n, y_{n+1}, y_{n+2})$$

Then we have

$$d_n = G(y_n, y_{n+1}, y_{n+2})$$

$$= G(Ax_n, Bx_{n+1}, Cx_{n+2})$$

$$\leq a G(Bx_{n+1}, Ax_n, Rx_n) + b G(Cx_{n+2}, Bx_{n+1}, Tx_{n+1}) + c G(Ax_n, Cx_{n+1}, Sx_{n+1}) + d \max\{G(Cx_{n+2}, Bx_{n+1}, Tx_{n+1}) + G(Tx_{n+1}, Ax_n, Rx_n), G(Ax_n, Cx_{n+2}, Sx_{n+2}), G(Sx_{n+2}, Bx_{n+1}, Tx_{n+1}), G(Bx_{n+1}, Ax_n, Rx_n), G(Rx_n, Cx_{n+2}, Sx_{n+2})\}$$

$$\leq a G(y_{n+1}, y_n, y_{n-1}) + b G(y_{n+2}, y_{n+1}, y_n) + c G(y_n, y_{n+1}, y_n) + d \max\{G(y_{n+2}, y_{n+1}, y_n) + G(y_n, y_n, y_{n-1}) + G(y_n, y_{n+2}, y_{n+1}), G(y_{n+1}, y_{n+1}, y_n) + G(y_{n+1}, y_n, y_{n-1}), G(y_{n-1}, y_{n+2}, y_{n+1})\}$$

$$\leq a d_{n-1} + b d_n + c d_n + d \max\{d_n + d_{n-1} + d_n, d_n + d_{n-1}, d_n + d_{n-1}\}$$

Clearly max will be last term

$$\text{i.e. } d_n + d_{n-1}$$

$$d_n \leq a d_{n-1} + b d_n + c d_n + d (d_n + d_{n-1})$$

$$d_n \leq (a + d) d_{n-1} + (b + c + d) d_n$$

$$(1 - b - c - d) d_n \leq (a + d) d_{n-1}$$

$$d_n \leq \left(\frac{a + b}{1 - b - c - d} \right) d_{n-1}$$

So we have

$$d_n \leq d_{n-1}$$

Next we find.

$$d_{n+1} = G(y_{n+1}, y_{n+2}, y_{n+3})$$

$$= G(Ax_{n+1}, Bx_{n+2}, Cx_{n+3})$$

$$\leq a G(Bx_{n+2}, Ax_{n+1}, Rx_{n+1}) + b G(Cx_{n+3}, Bx_{n+2}, Tx_{n+2}) + c G(Ax_{n+1}, Cx_{n+3}, Sx_{n+3}) + d \max\{G(Cx_{n+3}, Bx_{n+2}, Tx_{n+2}), G(Tx_{n+2}, Ax_{n+1}, Rx_{n+1}), G(Ax_{n+1}, Cx_{n+3}, Sx_{n+3}), G(Sx_{n+3}, Bx_{n+2}, Tx_{n+2}), G(Bx_{n+2}, Ax_{n+2}, Rx_{n+1}), G(Rx_{n+1}, Cx_{n+3}, Sx_{n+3})\}$$

$$\leq a G(y_{n+2}, y_{n+1}, y_n) + b G(y_{n+3}, y_{n+2}, y_{n+1}) + c G(y_{n+1}, y_{n+3}, y_{n+2}) + d \max\{G(y_{n+3}, y_{n+2}, y_{n+1}) G(y_{n+1}, y_{n+1}, y_n), G(y_{n+1}, y_{n+3}, y_{n+2}), G(y_{n+2}, y_{n+2}, y_{n+1}), G(y_{n+2}, y_{n+1}, y_n), G(y_n, y_{n+3}, y_{n+2})\}$$

Thus for every $n \in \mathbb{N}$ we have $d_n \leq d_{n-1}$. Further we can say $d_n \leq p d_{n-1}$ where $p = a+b+2c +3d$. Thus we have

$$d_n = G(y_n, y_{n+1}, y_{n+2}) \leq p G(y_{n-1}, y_n, y_{n+1}) \leq \dots \leq p^n G(y_0, y_1, y_2)$$

And as $G(x, x, y) \leq G(x, y, z)$ we have $G(y_n, y_n, y_{n+1}) \leq p^n G(y_0, y_1, y_2)$

So,

$$G(y_n, y_n, y_m) \leq G(y_n, y_n, y_{n+1}) + G(y_{n+1}, y_{n+1}, y_{n+2}) + \dots + G(y_{m-1}, y_{m-1}, y_m)$$

$$\leq p^n G(y_0, y_1, y_2) + p^{n+1} G(y_0, y_1, y_2) + \dots + p^{m-1} G(y_0, y_1, y_2)$$

$$\leq (p^n + p^{n+1} + \dots + p^{m-1}) G(y_0, y_1, y_2)$$

$$\leq \left[\frac{p^n}{1 - p} \right] G(y_0, y_1, y_2) \rightarrow 0 \text{ as } n \rightarrow \infty$$

So, the sequence $\{y_n\}$ is Cauchy sequence. in X , let it converge to a point y in X i.e. $\lim_{n \rightarrow \infty} y_n = y$. So the subsequences $\{A_{x_n}\}$, $\{T_{x_{n+1}}\}$, $\{B_{x_{n+1}}\}$, $\{S_{x_{n+2}}\}$, $\{C_{x_{n+2}}\}$, $\{R_{x_{n+3}}\}$ all converge to y . As $C(x)$ is Closed Subset of $R(X)$ these exist $u \in X$ such that $Ru = y$. Now we shall prove that $Au = y$

Consider

$$\begin{aligned} G(Au, B_{x_{n+1}}, C_{x_{n+2}}) &\leq a G(B_{x_{n+1}}, Au, Ru) + b G(C_{x_{n+2}}, B_{x_{n+1}}, T_{x_{n+1}}) + c G(Au, C_{x_{n+2}}, S_{x_{n+2}}) \\ &\quad + d \max\{G(C_{x_{n+2}}, B_{x_{n+1}}, T_{x_{n+1}}), G(T_{x_{n+1}}, Au, Ru), G(Au, C_{x_{n+2}}, S_{x_{n+2}}), G(S_{x_{n+2}}, B_{x_{n+1}}, T_{x_{n+1}}), \\ &\quad G(B_{x_{n+1}}, Au, Ru), G(Ru, C_{x_{n+2}}, S_{x_{n+2}})\} \\ &\leq a G(y, Au, Ru) + b G(y, y, y) + c G(Au, y, y) + d \max\{G(y, y, y), G(y, Au, Ru), \\ &\quad G(Au, y, y), G(y, y, y), G(y, Au, Ru), G(Ru, y, y), G(y, Au, Ru), G(Ru, y, y)\} \\ &\leq a G(y, Au, Ru) + c G(Au, y, y) + d \max\{G(Au, Ru, y), G(Au, y, y), G(Ru, y, y), \text{as } Ru = y \\ &\leq a G(y, Au, y) + c G(Au, y, y) + d \max\{G(Au, y, y), G(Au, y, y), G(y, y, y)\} \end{aligned}$$

Or

$$G(Au, y, y) \leq (a + c + d) G(Au, y, y)$$

$$\Rightarrow (1 - a - c - d) G(Au, y, y) \leq 0$$

$$\Rightarrow G(Au, y, y) = 0$$

$$\Rightarrow Au = y \text{ as } Ru = y, Au = y$$

$$\Rightarrow Au = Ru = y$$

Now we will use the definition of weak compatibility of (R, A)

$$\Rightarrow ARu = RAu$$

$$\Rightarrow Ay = Ry$$

Now we shall prove $Ay = y$ and on the contrary Let $Ay \neq y$ then we consider on

$$\begin{aligned} G(Ay, B_{x_{n+1}}, C_{x_{n+2}}) &\leq a G(B_{x_{n+1}}, Ay, Ry) + b G(C_{x_{n+2}}, B_{x_{n+1}}, T_{x_{n+1}}) + c d (Ay, C_{x_{n+2}}, S_{x_{n+2}}) \\ &\quad + d \max\{G(C_{x_{n+2}}, B_{x_{n+1}}, T_{x_{n+1}}), G(T_{x_{n+1}}, Ay, Ry), G(Ay, C_{x_{n+2}}, S_{x_{n+2}}), G(S_{x_{n+2}}, B_{x_{n+1}}, T_{x_{n+1}}), \\ &\quad G(B_{x_{n+1}}, Ay, Ry), G(Ry, C_{x_{n+2}}, S_{x_{n+2}})\} \\ &\leq a G(y, Ay, Ry) + b G(y, y, y) + c d (Ay, y, y) + d \max\{G(y, y, y), G(y, Ay, Ay), G(Ay, y, y), \\ &\quad G(y, y, y), G(y, Ay, Ay), G(Ay, y, y)\} \\ &\leq a G(y, y, Ay) + c G(y, y, Ay) + d G(Ay, y, y) \end{aligned}$$

$$\Rightarrow G(Ay, y, y) \leq (a + c + d) G(Ay, y, y)$$

$$\Rightarrow Ay = y = Ry.$$

Thus y is common fixed point of A and R . As $y = Ay \subseteq T(X) \exists w$ such that $Tv = y$. we shall now prove $Bv = y$

$$G(y, Bv, C_{x_{n+2}}) = G(Ay, Bv, C_{x_{n+2}})$$

$$\begin{aligned} &\leq a G(Bv, Ay, Ry) + b G(C_{x_{n+2}}, Bv, Tv) + c G(Ay, C_{x_{n+2}}, S_{x_{n+2}}) + d \max\{G(C_{x_{n+2}}, Bv, Tv), G(Tv, Ay, Ry), \\ &\quad G(Ay, C_{x_{n+2}}, S_{x_{n+2}}), G(S_{x_{n+2}}, Bv, Tv), G(Bv, Ay, Ry), G(Ry, C_{x_{n+2}}, S_{x_{n+2}})\} \\ &\leq a G(Bv, y, y) + b G(y, Bv, Tv) + c G(y, y, y) + d \max\{G(y, Bv, Tv), G(Tv, Ay, Ay), G(y, y, y), \\ &\quad G(y, Bv, Tv), G(Bv, y, y), G(Ay, y, y)\} \\ &\leq a G(Bv, y, y) + b G(Bv, y, y) + d (Bv, y, y) \end{aligned}$$

$$\Rightarrow (1 - a - b - d) G(Bv, y, y) \leq 0$$

$$\Rightarrow Bv = y$$

Hence

$$Bv = y = Tv$$

By weak compatibility $BTv = TBv$ hence $By = Ty$

Now we prove $By = y$. So on the contrary Let $By \neq y$

$$\begin{aligned} G(Ay, By, Cx_{n+2}) &\leq a G(By, Ay, Ry) + bG(Cx_{n+2}, By, Ty) + c G(Ay, Cx_{n+2}, Sx_{n+2}) + d \max\{G(Cx_{n+2}, By, Ty), \\ &G(Ty, Ay, Ry), G(Ay, Cx_{n+2}, Sx_{n+2}), G(Sx_{n+2}, By, Ty), G(By, Ay, Ry), G(Ry, Cx_{n+2}, Sx_{n+2})\} \\ &\leq aG(Ty, Ay, Ry) + bG(y, By, y) + G(y, y, y) + d \max\{G(y, By, By), G(y, y, y), G(y, y, y), G(y, y, y), \\ &G(By, y, y), G(y, y, y)\} \end{aligned}$$

$$G(y, By, y) \leq b G(y, By, y) + d G(y, By, y)$$

$$\Rightarrow (1 - b - d) G(y, By, y) \leq 0$$

Hence $By = y = Ty$. As $y = By \in S(X)$ there exist $w \in X$ such that $y = sw$. We now prove that $Cw = y$.

$$\begin{aligned} G(y, y, Cw) &= G(Ay, By, Cw) \\ &\leq a G(By, Ay, Ry) + b G(Cw, By, Ty) + c G(Ay, Cw, Sw) + d \max\{G(Cw, By, Ty), G(Ty, Ay, Ry), \\ &G(Ay, Cw, Sw), G(Sw, By, Ty), G(By, Ay, Ry), G(Ry, Cw, Sw)\} \\ &\leq a G(y, y, y) + b G(Cw, y, y) + c G(y, Cw, y) + d \max\{G(Cw, y, y), G(y, y, y), G(y, Cw, y), G(y, y, y), \\ &G(y, y, y), G(Cw, y, y)\} \\ &\leq (b + c + b) G(cw, y, y) \end{aligned}$$

Or

$$(1 - b - c - d) G(cw, y, y) \leq 0$$

$$\Rightarrow Cw = y$$

$$G(y, y, Cy) = G(Ay, By, Cy)$$

$$\begin{aligned} &\leq a G(By, Ay, Ry) + b G(Cy, By, Ty) + c G(Ay, Cy, Sy) + d \max\{G(Cy, By, Ty), G(Ty, Ay, Ry), \\ &G(Ay, Cy, Sy), G(Sy, By, Ty), G(By, Ay, Ry), G(Ry, Cy, Sy)\} \\ &\leq a G(y, y, y) + bG(y, y, y) + c G(y, Cy, y) + d \max\{G(Cy, y, y), G(y, y, y), G(y, Cy, y), G(y, y, y), \\ &G(y, y, y), G(y, Cy, y)\} \\ &\leq (b + c + d) G(y, y, y) \end{aligned}$$

Or

$$G(Cy, y, y) (1 - b - c - d) \leq 0$$

$$G(Cy, y, y) = 0$$

$Cy = y$ as $Sy = y \Rightarrow Sy = Cy = y$ i.e. y is a common fixed point of S and C . Thus y will be the common fixed point of A, B, C, S, T and R

i.e

$$Ay = Sy = By = Ty = Cy = Ry = y$$

Now we will prove the uniqueness. Let y' is another fixed point of A, B, C, S, T and R .

$$\begin{aligned} G(y, y, y') &= G(Ay, By, Cy) \leq a G(By, Ay, Ry) + b G(Cy', By, Ty) + c G(Ay, Cy', Sy) + d \max\{G(Cy', By, Ty) \\ &+ G(Ty, Ay, Ry) + G(Ay, Cy', Sy), G(Sy, By, Ty) + G(By, Ay, Ry), G(Ry, Cy', Sy)\} \\ &\leq a G(y, y, y) + b G(y', y, y) + c G(y, y', y') + d \max\{G(y', y, y) + G(y, y, y) + G(y, y', y'), G(y', y, y) \\ &+ G(y, y, y), G(y, y, y')\} \\ &\leq b G(y', y, y) + 2 c G(y', y, y) + d \max\{3 G(y', y, y), G(y', y, y), 2 G(y', y, y)\} \\ &\leq (b + 2 c) G(y', y, y) + 3 d G(y', y, y) \end{aligned}$$

Or

$$G(y, y, y') \leq (b + 2c + 3d) G(y', y, y)$$

$$(1 - b - 2c - 3d) G(y, y, y) \leq 0$$

$$\text{As } a + b + 2c + 3d < 1$$

$$\text{We get } G(y, y, y') \leq 0$$

$$\Rightarrow y = y'$$

Hence y is the unique common fixed point of A, B, C, S, T and R .

REFERENCES

1. Jungck, Compatible mappings and common fixed points, *Int. J. Math. Sci.*, 9 (4) (1986), 771-779.
2. G. Jungck, Common fixed points for non continuous nonself maps on non-metric spaces, *far East J. Math. Sci.*, 4 (1996), 199-215.
3. G. Jungck, B.E. Rhoades, fixed points for set valued functions without continuity, *Indian J. Pure Appl. Math.* 29 (1998), 227-238.
4. S. Sessa, On a weak commutativity condition of mappings in fixed point consideration, *Publ. Inst. Math. Soc.* 32 (1982), 149-153.
5. Z. Mustafa, B. Sims, Some remarks concerning D-metric spaces, in *Proceedings of the International Conferences on Fixed Point Theory and Applications*, pp. 189-198, Valencia, Spain, July 2003.
6. Z. Mustafa, B. Sims, A new approach to generalized metric spaces, *J. Nonlinear Convex Anal.* 7 (2006), 289-297.

Source of Support: Nil, Conflict of interest: None Declared

[Copy right © 2016, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]