



GENERAL TOPOLOGICAL INDICES OF CIRCUMCORONENE SERIES OF BENZENOID

V. R. KULLI\*

Department of Mathematics, Gulbarga University, Gulbarga 585106, India.

(Received On: 04-04-17; Revised & Accepted On: 05-05-17)

ABSTRACT

In this paper, we determine generalized version of the first Zagreb index, general connectivity index, general sum connectivity index, general reformulated index and other topological indices for circumcoronene series of benzenoid.

**Keywords:** Zagreb indices, connectivity index, sum connectivity index, reformulated index, K-edge index, circumcoronene series of benzenoid.

**Mathematics Subject Classification:** 05C05.

1. INTRODUCTION

The graphs considered here are finite, undirected without loops and multiple edges. Let  $G$  be a connected graph. The degree  $d_G(v)$  of a vertex  $v$  is the number of vertices adjacent to  $v$ . The edge connecting the vertices  $u$  and  $v$  will be denoted by  $uv$ . Let  $d_G(e)$  denote the degree of an edge  $e$  in  $G$ , which is defined by  $d_G(e) = d_G(u) + d_G(v) - 2$  with  $e = uv$ . For all further notation and terminology we refer to reader to [1].

A molecular graph is a simple graph related to the structure of a chemical compound. Each vertex of this graph represents an atom of the molecule and its edges to the bonds between atoms. A topological index is a numerical parameter mathematically derived from the graph structure. These indices are useful for establishing correlation between the structure of a molecular compound and its physico-chemical properties.

The first and second Zagreb indices of a graph  $G$  are defined as

$$M_1(G) = \sum_{u \in V(G)} d_G(u)^2 \quad \text{or} \quad M_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]$$

$$M_2(G) = \sum_{uv \in E(G)} d_G(u)d_G(v).$$

These indices were introduced by Gutman *et al.* in [2].

Another vertex degree based topological index was defined in [2] and it was studied by Furtula *et al.* in [3].

The forgotten topological index or F-index is defined as

$$F(G) = \sum_{u \in V(G)} d_G(u)^3.$$

The generalized version of the first Zagreb index [4] of a graph  $G$  is defined as

$$M_1^{a+1}(G) = \sum_{u \in V(G)} d_G(u)^{a+1} = \sum_{uv \in E(G)} [d_G(u)^a + d_G(v)^a], \text{ where } a \in \mathbb{R}. \tag{1}$$

**Corresponding Author: V. R. Kulli\***

**Department of Mathematics, Gulbarga University, Gulbarga 585106, India.**

The modified first and second Zagreb indices [5] are respectively defined as

$${}^m M_1(G) = \sum_{u \in V(G)} \frac{1}{d_G(u)^2}, \quad {}^m M_2(G) = \sum_{uv \in E(G)} \frac{1}{d_G(u)d_G(v)}.$$

The first hyper-Zagreb index of a graph  $G$  is defined as

$$HM_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2.$$

This index was introduced by Shirdel *et al.* in [6].

The sum connectivity index of a graph  $G$  is defined as

$$X(G) = \sum_{uv \in E(G)} \sqrt{\frac{1}{d_G(u) + d_G(v)}}$$

This index was introduced by Zhou and Trinajstić in [7].

The general sum connectivity index was introduced by Zhou and Trinajstić in [8] and it is defined as

$$M_1^a(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^a. \tag{2}$$

This index was also studied, for example, in [9].

In [10], the second hyper Zagreb index of a graph  $G$  is defined as

$$HM_2(G) = \sum_{uv \in E(G)} [d_G(u)d_G(v)]^2.$$

The Randić index or product connectivity index of a graph  $G$  is defined as

$$\chi(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_G(u)d_G(v)}}.$$

This topological index was proposed by Randić in [11] and was studied, for example, in [12, 13].

The general product connectivity index [9, 14] is defined as

$$M_2^a(G) = \sum_{uv \in E(G)} [d_G(u)d_G(v)]^a \tag{3}$$

The reformulated first Zagreb index of a graph  $G$  is defined as

$$EM_1(G) = \sum_{e \in E(G)} d_G(e)^2.$$

This index was introduced by Miličević *et al.* in [15].

Recently in [16], Kulli introduced the  $K$ -edge index of a graph  $G$  and it is defined as


$$K_e(G) = \sum_{e \in E(G)} d(e)^3.$$

This index was also studied in [17].

The general reformulated Zageb index [18] of a graph  $G$  is defined as

$$EM_1^a(G) = \sum_{e \in E(G)} d(e)^a \tag{4}$$

where  $a$  is a real number.



**Figure-1:** The graph of circumcoronene series of benzenoid  $H_k, k \geq 1$ .

In this paper, we compute several topological indices for circumcoronene series of benzenoid  $H_k, k \geq 1$ .

The benzene molecule is a molecule in chemistry and nanosciences and it is useful to synthesize aromatic compounds. The circumcoronene series of benzenoid is a benzenoid family and it is symbolized by  $H_k, k \geq 1$  which consists copies of benzene  $C_6$  on circumference, see [19].

By algebraic method, we obtain  $|V(H_k)| = 6k^2$  and  $|E(H_k)| = 9k^2 - 3k$ . From Figure 1, it is easy to see that there are two partitions of the vertex set of  $H_k$  as follows:

$$V_2 = \{u \in V(H_k) \mid d_{H_k}(u) = 2\}, |V_2| = 6k.$$

$$V_3 = \{u \in V(H_k) \mid d_{H_k}(u) = 3\}, |V_3| = 6k(k - 1).$$

Also by algebraic method, we obtain three edge partitions of  $H_k$  based on the sum of degrees of the end vertices as follows:

$$E_4 = E_4^* = \{uv \in E(H_k) \mid d_{H_k}(u) = d_{H_k}(v) = 2\}, |E_4| = |E_4^*| = 6.$$

$$E_5 = E_6^* = \{uv \in E(H_k) \mid d_{H_k}(u) = 2, d_{H_k}(v) = 3\}, |E_5| = |E_6^*| = 12(k - 1)$$

$$E_6 = E_9^* = \{uv \in E(H_k) \mid d_{H_k}(u) = d_{H_k}(v) = 3\}, |E_6| = |E_9^*| = 9k^2 - 15k + 6.$$

The edge degree partition of  $H_k$  is given in Table 1.

$d_{H_k}(u), d_{H_k}(v) \setminus e = uv \in E(H_k)$	(2, 2)	(2, 3)	(3,3)
$d_{H_k}(e)$	2	3	4
Number of edges	6	$12(k - 1)$	$9k^2 - 15k + 6$

**Table-1:** Edge degree partition of  $H_k$

We compute the generalized version of the first Zagreb index of  $H_k$ .

**Theorem 1:** The generalized version of the first Zagreb index of  $H_k$  is given by

$$M_1^{a+1}(H_k) = (3^{a+1})6k^2 + (2^{a+1} - 3^{a+1})6k. \tag{5}$$

**Proof:** Let  $G = H_k$ . From equation (1) and by cardinalities of the vertex partition of  $H_k$ , we have

$$M_1^{a+1}(H_k) = \sum_{u \in V(H_k)} d(u)^{a+1} = \sum_{u \in V_2} d(u)^{a+1} + \sum_{u \in V_3} d(u)^{a+1}$$

$$= 2^{a+1}6k + 3^{a+1}6k(k - 1) = (3^{a+1})6k^2 + (2^{a+1} - 3^{a+1})6k.$$

We obtain the following corollaries by using Theorem 1.

**Corollary 1.1:** The first Zagreb index of  $H_k$  is given by  $M_1(H_k) = 54k^2 - 30k$ .

**Proof:** Put  $a = 1$  in equation (5), we obtain the desired result.

**Corollary 1.2:** The  $F$ -index of  $H_k$  is given by  $F(H_k) = 162k^2 - 114k$ .

**Proof:** Put  $a = 2$  in equation (5), we obtain the desired result.

**Corollary 1.3:** The modified first Zagreb index of  $H_k$  is given by  ${}^m M_1(H_k) = \frac{2}{3}k^2 + \frac{5}{36}k$ .

**Proof:** Put  $a = -3$  in equation (5), we obtain the desired result.

In the next result, we determine the general sum connectivity index of  $H_k$ ,  $k \geq 1$ .

**Theorem 2:** The general sum connectivity index of  $H_k$  is given by

$$M_1^a(H_k) = 9 \times 6^a k^2 + (4 \times 5^a - 5 \times 6^a)3k + (4^a - 2 \times 5^a + 6^a)6 \tag{6}$$

**Proof:** From equation (2) and cardinalities of the edge partitions of  $H_k$ , we have

$$\begin{aligned} M_1^a(H_k) &= \sum_{uv \in E(H_k)} [d_{H_k}(u) + d_{H_k}(v)]^a \\ &= \sum_{uv \in E_4} [d_{H_k}(u) + d_{H_k}(v)]^a + \sum_{uv \in E_5} [d_{H_k}(u) + d_{H_k}(v)]^a + \sum_{uv \in E_6} [d_{H_k}(u) + d_{H_k}(v)]^a \\ &= 4^a \times 6 + 5^a \times 12(k-1) + 6^a \times (9k^2 - 12k + 6) \\ &= 9 \times 6^a k^2 + (4 \times 5^a - 5 \times 6^a)3k + (4^a - 2 \times 5^a + 6^a)6 \end{aligned}$$

We obtain the following corollaries by using Theorem 2.

**Corollary 2.1:** The first Zagreb index of  $H_k$  is given by  $M_1(H_k) = 54k^2 - 30k$ .

**Proof:** Put  $a = 1$  in equation (6), we get the desired result.

**Corollary 2.2:** The first hyper-Zagreb index of  $H_k$  is given by  $HM_1(H_k) = 324k^2 - 240k + 12$ .

**Proof:** Put  $a = 2$  in equation (6), we get the desired result.

**Corollary 2.3:** The sum connectivity index of  $H_k$  is given by  $X(H_k) = \frac{1}{10} (15k^2 - k + 1)$ .

**Proof:** Put  $a = -\frac{1}{2}$  in equation (6), we get the desired result.

We now compute the general product connectivity index of  $H_k$ ,  $k \geq 1$ .

**Theorem 3:** The general product connectivity index of  $H_k$  is given by

$$M_2^a(H_k) = 9^{a+1}k^2 + (12 \times 6^a - 15 \times 9^a)k + (6 \times 4^a - 12 \times 6^a + 6 \times 9^a). \tag{7}$$

**Proof:** From equation (3) and by cardinalities of the edge partition of  $H_k$  based on the product degrees of the end vertices, we have

$$\begin{aligned} M_2^a(H_k) &= \sum_{uv \in E(H_k)} [d_{H_k}(u)d_{H_k}(v)]^a \\ &= \sum_{uv \in E_4} [d_{H_k}(u)d_{H_k}(v)]^a + \sum_{uv \in E_5} [d_{H_k}(u)d_{H_k}(v)]^a + \sum_{uv \in E_6} [d_{H_k}(u)d_{H_k}(v)]^a \\ &= 4^a \times 6 + 6^a \times 12(k-1) + 9^a \times (9k^2 - 15k + 6) \\ &= 9^{a+1}k^2 + (12 \times 6^a - 15 \times 9^a)k + (6 \times 4^a - 12 \times 6^a + 6 \times 9^a). \end{aligned}$$

We obtain the following corollaries by Theorem 3.

**Corollary 3.1:** The second Zagreb index of  $H_k$  is given by  $M_2(H_k) = 81k^2 - 63k + 6$ .

**Proof:** Put  $a = 1$  in equation (7), we get the desired result.

**Corollary 3.2:** The second hyper Zagreb index of  $H_k$  is given by

$$HM_2(H_k) = 729k^2 - 783k + 150.$$

**Proof:** Put  $a = 2$  in equation (7), we get the desired result.

**Corollary 3.3:** The modified second Zagreb index of  $H_k$  is given by  ${}^m M_2(H_k) = k^2 + \frac{1}{3}k + \frac{1}{6}$ .

**Proof:** Put  $a = -1$  in equation (7), we get the desired result.

**Corollary 3.4:** The Randić connectivity index of  $H_k$  is given by  $\chi(H_k) = 3k^2 + (2\sqrt{6} - 5)(k - 1)$

**Proof:** Put  $a = -\frac{1}{2}$  in equation (7), we get the desired result.

In the following theorem, we compute the general first reformulated Zagreb index of  $H_k$ ,  $k \geq 1$ .

**Theorem 4:** The general first reformulated Zagreb index of  $H_k$  is given by

$$EM_1^a(H_k) = 9 \times 4^a k^2 + (12 \times 3^a - 15 \times 4^a)k + (6 \times 2^a - 12 \times 3^a + 6 \times 4^a). \quad (8)$$

**Proof:** From equation (4) and by the edge degree partition of  $H_k$ , we have

$$\begin{aligned} EM_1^a(H_k) &= \sum_{e \in E(G)} d_G(e)^a = \sum_{e \in E_4} d_G(e)^a + \sum_{e \in E_5} d_G(e)^a + \sum_{e \in E_6} d_G(e)^a \\ &= 6 \times 2^a + 12(k-1) \times 3^a + (9k^2 - 15k + 6) \times 4^a \\ &= 9 \times 4^a k^2 + (12 \times 3^a - 15 \times 4^a)k + (6 \times 2^a - 12 \times 3^a + 6 \times 4^a). \end{aligned}$$

We obtain the following results by using Theorem 4.

**Corollary 4.1:** The first reformulated Zagreb index of  $H_k$  is given by  $EM_1(H_k) = 144k^2 - 132k + 12$ .

**Proof:** Put  $a = 2$  in equation (8), we get the desired result.

**Corollary 4.2:** The  $K$ -edge index of  $H_k$  is given by  $K_e(H_k) = 576k^2 - 636k + 108$ .

**Proof:** Put  $a = 3$  in equation (8), we get the desired result.

## REFERENCES

1. V.R.Kulli, *College Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
2. I. Gutman and N. Trinajstić, Graph theory and molecular orbitals. Total  $\pi$ -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* 17, (1972) 535-538.
3. B. Furtula, I. Gutman, A forgotten topological index, *J. Math. Chem.* 53 (2015), 1184-1190.
4. X.Li and J. Zheng, A unified approach to the extremal trees for different indices, *MATCH Commun. Math. Comput. Chem.* 54 (2005) 195-208.
5. S. Nikolić, G. Kovačević, A. Miličević and N. Trinajstić, The Zagreb indices 30 years after, *Croatica Chemica Acta CCACAA* 76(2), (2003) 113-124.
6. G.H.Shirdel, H.Rezapour and A.M.Sayadi, The hyper-Zagreb index of graph operations, *Iranian J.Math.Chem.* 4(2) (2013) 213-220.
7. B. Zhou and N.Trinajstić, On a novel connectivity index, *J. Math. Chem.* 46 (2009) 1252-1270.
8. B. Zhou and N.Trinajstić, , *J. Math. Chem.* 47 (2010) 210.
9. V.R.Kulli, The first and second  $ka$  indices and coindices of graphs, *Internal Journal of Mathematical Archive*, 7(5) (2016) 71-77.
10. M.R.Farahani, M.R.R.Kanna and R.Pradeep Kumar, On the hyper-Zagreb indices of nanostructures, *Asian Academic Research J. Multidisciplinary*, 3(1), (2016).

11. M. Randić, On characterization of molecular branching, *Journal of the American Chemical Society*, 97(23) (1975) 6609-6615.
12. V.R.Kulli, B.Chaluvaraju and H.S.Boregowda, Some degree based connectivity indices of Kulli cycle windmill graphs, *South Asian Journal of Mathematics*, 6(6) (2016) 263-268.
13. V.R.Kulli, B.Chaluvaraju and H.S.Boregowda, Computation of connectivity indices of path windmill graphs, submitted.
14. L. Clark and I. Gutman, *J. Math. Chem.* 43 (2008) 32.
15. A. Miličević, S.Nikolić and N. Trinajstić, On reformulated Zagreb indices, *Mol. Divers.* 8 (2004) 393-399.
16. V.R. Kulli, On  $K$  - edge index and coindex of graphs, *International Journal of Fuzzy Mathematical Archive*, 10(2) (2016) 111-116.
17. V.R.Kulli,  $K$ -edge index of some nanostructures, *Journal of Computer and Mathematical Sciences*, 7(7) (2016) 373-378.
18. V.R.Kulli, Computation of general topological indices for titania nanotubes, *International Journal of Mathematical Archive*, 7(12) (2016) 33-38.
19. S.Klavzar, A bird's eye view of the cut method and a survey of its applications in chemical graph theory, *MATCH Commun. Math. Comput. Chem.* 60 (2008) 255-274.

**Source of Support: Nil, Conflict of interest: None Declared**

**[Copy right © 2017, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**