

INVERSE SUM LEAP INDEX  
AND HARMONIC LEAP INDEX OF CERTAIN WINDMILL GRAPHS

V. R. KULLI\*

Department of Mathematics, Gulbarga University, Gulbarga 585 106, India.

(Received On: 06-02-19; Revised & Accepted On: 16-02-19)

---

ABSTRACT

We propose two new topological indices the inverse sum leap index and harmonic leap index of a graph. Also we compute the inverse sum leap index and harmonic leap index of Kulli cycle windmill graphs, Kulli path windmill graphs, Dutch windmill graphs and French windmill graphs.

**Keywords:** Inverse sum leap index, harmonic leap index, windmill graph.

**Mathematics Subject Classification:** 05C05, 05C07, 05C12, 05C35.

---

1. INTRODUCTION

Let  $G$  be a finite, simple, connected graph with vertex set  $V(G)$  and edge set  $E(G)$ . The degree  $d(v)$  of a vertex  $v$  is the number of edges incident to  $v$ . We refer [1] for other undefined notations and terminologies.

The distance  $d(u, v)$  between any two vertices  $u$  and  $v$  is equal to the length of a shortest path connecting them. For a positive integer  $k$ , the open  $k$ -neighborhood  $N_k(v)$  of a vertex  $v$  in  $G$  is defined as  $N_k(v) = \{u \in V(G) : d(u, v) = k\}$ . The  $k$ -distance degree  $d_k(v)$  of  $v$  in  $G$  is defined as the number of  $k$  neighbors of  $v$  in a graph  $G$ .

In [2], Vukičević et al. observed that many topological indices are defined simply as the sum of individual bond contributions. They have proposed a class of discrete Adriatic indices to study whether there other possibly significant topological indices of this form. One of these discrete Adriatic indices is the inverse sum indeg index, which is defined as

$$ISI(G) = \sum_{uv \in E(G)} \frac{d(u)d(v)}{d(u) + d(v)}$$

This index was also studied in [3, 4, 5].

Motivated by the definition of inverse sum indeg index, we propose the inverse sum leap index and harmonic leap index of a graph as follows:

The inverse sum leap index of a graph  $G$  is defined as

$$ISL(G) = \sum_{uv \in E(G)} \frac{d_2(u)d_2(v)}{d_2(u) + d_2(v)} \tag{1}$$

The harmonic leap index of a graph  $G$  is defined as

$$HL(G) = \sum_{uv \in E(G)} \frac{2}{d_2(u) + d_2(v)}$$

The general harmonic leap index of a graph  $G$  is defined as

$$HL^a(G) = \sum_{uv \in E(G)} \left( \frac{2}{d_2(u) + d_2(v)} \right)^a \tag{2}$$

---


Corresponding Author: V. R. Kulli\*  
Department of Mathematics, Gulbarga University, Gulbarga 585 106, India.

Recently, some leap indices were introduced and studied such as minus leap and square leap indices [6], leap hyper Zagreb indices [7], sum connectivity leap and geometric-arithmetic leap index and ABC leap index [8], F-leap indices [9], product connectivity leap index and ABC leap index [10], augmented leap index [11]. The harmonic index was studied in [12, 13, 14].

In this paper, the inverse sum leap index, harmonic leap index and general harmonic leap index of four types of windmill graphs are computed. For windmill graphs, see [15, 16, 17, 18].

**2. RESULTS FOR KULLI PATH WINDMILL GRAPHS**

The Kulli path windmill graph  $P_{n+1}^m$  is the graph obtained by taking  $m$  copies of the graph  $K_1 + P_n$  with a vertex  $K_1$  in common, see Figure 1.



**Figure-1:** Kulli path windmill graph  $P_{n+1}^m$

Let  $G = P_{n+1}^m$  be a Kulli path windmill graph with  $mn + 1$  vertices and  $2mn - m$  edges,  $m \geq 2, n \geq 5$ . Then  $G$  has four types of the 2-distance degree of edges as given in Table 1.

| $d_2(u), d_2(v) \setminus uv \in E(G)$ | $(0, mn - 2)$ | $(0, mn - 3)$ | $(mn - 2, mn - 3)$ | $(mn - 3, mn - 3)$ |
|--|---------------|---------------|--------------------|--------------------|
| Number of edges                        | $2m$          | $mn - 2m$     | $2m$               | $mn - 3m$          |

**Table-1:** 2-distance degree edge partition of  $P_{n+1}^m$

**Theorem 1:** The inverse sum leap index of a Kulli path windmill graph  $P_{n+1}^m$  is

$$ISL(P_{n+1}^m) = m(mn - 3) \left[ \frac{2mn - 4}{2mn - 5} + \frac{n - 3}{2} \right]$$

**Proof:** Let  $G = P_{n+1}^m$ . By using equation (1), we have

$$ISL(P_{n+1}^m) = \sum_{uv \in E(G)} \frac{d_2(u)d_2(v)}{d_2(u) + d_2(v)}$$

Then by using Table 1, we obtain

$$\begin{aligned} ISL(P_{n+1}^m) &= \left( \frac{0 \times (mn - 2)}{0 + mn - 2} \right) 2m + \left( \frac{0 \times (mn - 3)}{0 + mn - 3} \right) (mn - 2m) \\ &\quad + \left( \frac{(mn - 2) \times (mn - 3)}{mn - 2 + mn - 3} \right) 2m + \left( \frac{(mn - 3) \times (mn - 3)}{mn - 3 + mn - 3} \right) (mn - 3m) \\ &= m(mn - 3) \left[ \frac{2mn - 4}{2mn - 5} + \frac{n - 3}{2} \right]. \end{aligned}$$

**Theorem 2:** The general harmonic leap index of a Kulli path windmill graph  $P_{n+1}^m$  is

$$HL^a(P_{n+1}^m) = \left( \frac{2}{mn - 2} \right)^a 2m + \left( \frac{2}{mn - 3} \right)^a m(n - 2) + \left( \frac{2}{2mn - 5} \right)^a 2m + \left( \frac{1}{mn - 3} \right)^a m(n - 3) \tag{3}$$

**Proof:** Let  $G = P_{n+1}^m$ . From equation (2), we have

$$HL(P_{n+1}^m) = \sum_{uv \in E(G)} \left( \frac{2}{d_2(u) + d_2(v)} \right)^a.$$

Then by using Table 1, we derive

$$HL^a(P_{n+1}^m) = \left( \frac{2}{mn-2} \right)^a 2m + \left( \frac{2}{mn-3} \right)^a m(n-2) + \left( \frac{2}{2mn-5} \right)^a 2m + \left( \frac{1}{mn-3} \right)^a m(n-3)$$


**Corollary 2.1:** The harmonic leap index of  $P_{n+1}^m$  is

$$HL(P_{n+1}^m) = \left( \frac{2}{mn-2} \right) 2m + \left( \frac{2}{mn-3} \right) m(n-2) + \left( \frac{2}{2mn-5} \right) 2m + \left( \frac{1}{mn-3} \right) m(n-3)$$

**Proof:** Put  $a = 1$  in equation (3), we get the desired result.

### 3. RESULTS FOR KULLI CYCLE WINDMILL GRAPHS

The Kulli cycle windmill graph  $C_{n+1}^m$  is the graph obtained by taking  $m$  copies of the graph  $K_1 + C_n$  for  $n \geq 3$  with a vertex  $K_1$  is common, see Figure 2.



**Figure-2:** Kulli cycle windmill graph  $C_{n+1}^m$

Let  $G = C_{n+1}^m$  be a Kulli cycle windmill graph with  $mn + 1$  vertices and  $2mn$  edges,  $m \geq 2, n \geq 5$ . Then  $G$  has two types of the 2-distance degree of edges as given in Table 2.

| $d_2(u), d_2(v) \setminus uv \in E(G)$ | $(0, mn - 2)$ | $(mn - 2, mn - 2)$ |
|--|---------------|--------------------|
| Number of edges                        | $mn$          | $mn$               |

**Table-2:** 2-distance degree edge partition of  $C_{n+1}^m$

**Theorem 3:** The inverse sum leap index of a Kulli cycle windmill graph  $C_{n+1}^m$  is

$$ISL(C_{n+1}^m) = \frac{1}{2} mn(mn - 2).$$

**Proof:** Let  $G = C_{n+1}^m$ . From equation (1), we have

$$ISL(C_{n+1}^m) = \sum_{uv \in E(G)} \frac{d_2(u)d_2(v)}{d_2(u) + d_2(v)}$$

By using Table 2, we deduce

$$\begin{aligned} ISL(C_{n+1}^m) &= \left( \frac{0 \times (mn - 2)}{0 + mn - 2} \right) mn + \left( \frac{(mn - 2) \times (mn - 2)}{mn - 2 + mn - 2} \right) mn \\ &= \frac{1}{2} mn(mn - 2). \end{aligned}$$

**Theorem 4:** The general harmonic leap index of a Kulli cycle windmill graph  $C_{n+1}^m$  is

$$HL^a(C_{n+1}^m) = \left(\frac{2}{mn-2}\right)^a mn + \left(\frac{1}{mn-2}\right)^a mn. \tag{4}$$

**Proof:** Let  $G = C_{n+1}^m$ . By using equation (2), we get

$$HL^a(C_{n+1}^m) = \sum_{uv \in E(G)} \left(\frac{2}{d_2(u) + d_2(v)}\right)^a$$

From Table 2, we deduce


$$\begin{aligned} HL(C_{n+1}^m) &= \left(\frac{2}{0+mn-2}\right)^a mn + \left(\frac{2}{mn-2+mn-2}\right)^a mn. \\ &= \left(\frac{2}{mn-2}\right)^a mn + \left(\frac{1}{mn-2}\right)^a mn. \end{aligned}$$

**Corollary 4.1:** The harmonic leap index of  $C_{n+1}$  is  $\frac{3mn}{mn-2}$ .

**Proof:** Put  $a = 1$  is equation (4); we get the desired result.

#### 4. RESULTS FOR FRENCH WINDMILL GRAPHS

The French windmill graph  $F_{n+1}^m$  is the graph obtained by taking  $m \geq 2$  copies of the graph  $K_n$ ,  $n \geq 2$  with a vertex in common, see Figure 3.



**Figure-3:** French windmill graph  $F_{n+1}^m$

Let  $G = F_{n+1}^m$  be a French windmill graph which has  $1 + m(n - 1)$  vertices and  $\frac{1}{2}mn(n - 1)$  edges,  $m \geq 2, n \geq 2$ . Then there are two types of the 2-distance degree of edges as given in Table 3.

| $d_2(u), d_2(v) \setminus uv \in E(G)$ | $(0, (n - 1)(m - 1))$ | $((n - 1)(m - 1), (n - 1)(m - 1))$ |
|--|-----------------------|------------------------------------|
| Number of edges                        | $m(n - 1)$            | $\frac{1}{2}m(n - 1)(n - 2)$       |

Table 3. 2-distance degree edge partition of  $F_{n+1}^m$

**Theorem 5:** The inverse sum leap index of a French windmill graph  $F_{n+1}^m$  is

$$ISL(F_{n+1}^m) = \frac{1}{4}m(m - 1)(n - 1)^2(n - 2).$$

**Proof:** Let  $G = F_{n+1}^m$ . From equation (1), we obtain

$$ISL(F_{n+1}^m) = \sum_{uv \in E(G)} \frac{d_2(u)d_2(v)}{d_2(u) + d_2(v)}$$

By using Table 3, we deduce

$$IS(F_{n+1}^m) = \left( \frac{0 \times (n-1)(m-1)}{0 + (n-1)(m-1)} \right) m(m-1) + \left( \frac{(n-1)(m-1) \times (n-1)(m-1)}{(n-1)(m-1) + (n-1)(m-1)} \right) \frac{1}{2} m(n-1)(m-2) = \frac{1}{4} m(m-1)(n-1)^2 (n-2).$$

**Theorem 6:** The general harmonic leap index of a French windmill graph  $F_{n+1}^m$  is

$$HL^a(F_{n+1}^m) = \left( \frac{2}{(n-1)(m-1)} \right)^a m(n-1) + \left( \frac{1}{(n-1)(m-1)} \right)^a \frac{1}{2} m(n-1)(n-2). \tag{5}$$

**Proof:** Let  $G = F_{n+1}^m$ . From equation (2), we have

$$HL^a(F_{n+1}^m) = \sum_{uv \in E(G)} \left( \frac{2}{d_2(u) + d_2(v)} \right)^a$$

By using Table 3, we derive


$$HL^a(F_{n+1}^m) = \left( \frac{2}{0 + (n-1)(m-1)} \right)^a m(n-1) + \left( \frac{2}{(n-1)(m-1) + (n-1)(m-1)} \right)^a \frac{1}{2} m(n-1)(n-2) = \left( \frac{2}{(n-1)(m-1)} \right)^a m(n-1) + \left( \frac{1}{(n-1)(m-1)} \right)^a \frac{1}{2} m(n-1)(n-2).$$

**Corollary 6.1:** The harmonic leap index of  $F_{n+1}^m$  is  $\frac{m(n+2)}{2(m-1)}$ .

**Proof:** Put  $a = 1$  in equation (5), we get the desired result.

**5. RESULTS FOR DUTCH WINDMILL GRAPHS**

The Dutch windmill graph  $D_n^m$   $m \geq 2, n \geq 5$  is the graph obtained by taking  $m$  copies of the cycle  $C_n$  with a vertex in common, see Figure 4.



**Figure-4:** Dutch windmill graph  $D_n^m$

Let  $G = D_n^m$  be a Dutch windmill graph with  $1 + mn - m$  vertices and  $mn$  edges,  $m \geq 2, n \geq 5$ . Then  $G$  has three types of 2-distance degree of edges as given in Table 4.

|  |            |           |          |
|--|------------|-----------|----------|
| $d_2(u), d_2(v) \setminus uv \in E(G)$ | $(2m, 2m)$ | $(2m, 2)$ | $(2, 2)$ |
| Number of edges                        | $2m$       | $2m$      | $m(n-4)$ |

**Table-4:** 2-distance degree edge partition of  $D_n^m$

**Theorem 7:** The inverse sum leap index of a Dutch windmill graph  $D_n^m$  is

$$ISL(D_n^m) = m^2 \left( \frac{m+3}{m+1} \right) + m(n-4).$$

**Proof:** Let  $G = D_n^m$ . From equation (1), we have

$$ISL(D_n^m) = \sum_{uv \in E(G)} \frac{d_2(u)d_2(v)}{d_2(u)+d_2(v)}.$$

By using Table 4, we deduce

$$\begin{aligned} ISL(D_n^m) &= \left( \frac{2m \times 2m}{2m+2m} \right) 2m + \left( \frac{2m \times 2}{2m+2} \right) 2m + \left( \frac{2 \times 2}{2+2} \right) m(n-4) \\ &= m^2 \left( \frac{m+3}{m+1} \right) + m(n-4). \end{aligned}$$

**Theorem 8:** The general harmonic leap index of a Dutch windmill graph  $D_n^m$  is

$$HL^a(D_n^m) = \left( \frac{1}{2m} \right)^a 2m + \left( \frac{1}{m+1} \right)^a 2m + \left( \frac{1}{2} \right)^a 2m(n-4) \tag{6}$$

**Proof:** Let  $G = D_n^m$ . By using equation (2), we get

$$HL^a(D_n^m) = \sum_{uv \in E(G)} \left( \frac{2}{d_2(u)+d_2(v)} \right)^a$$

By using Table 4, we derive

$$\begin{aligned} HL^a(D_n^m) &= \left( \frac{2}{2m+2m} \right)^a 2m + \left( \frac{2}{2m+2} \right)^a 2m + \left( \frac{2}{2+2} \right)^a m(n-4) \\ &= \left( \frac{1}{2m} \right)^a 2m + \left( \frac{1}{m+1} \right)^a 2m + \left( \frac{1}{2} \right)^a m(n-4). \end{aligned}$$

**Corollary 8.1:** The harmonic leap index of  $D_n^m$  is

$$\frac{1}{2}mn - 2m + \frac{2m}{m+1} + 1.$$

**Proof:** Put  $a = 1$  in equation (6), we obtain the desired result.

**REFERENCES**

1. V.R.Kulli, *College Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
2. D. Vukičević and M. Gašperov, Bond additive modeling 1. Adriatic indices, *Croat. Chem. Acta*, 83(2010) 243-260.
3. V.R. Kulli, General topological indices of tetrameric 1, 3-adamantane, *International Journal of Current Research in Science and Technology*, 3(8) (2017) 26-33.
4. V.R. Kulli, Direct and inverse sum Bhanhatti indices, *International Research Journal of Pure Algebra* 7(9) (2017) 823-827.
5. J. Sedlar, D. Stevanović and A. Vasilyev, On the inverse sum indeg index, *Discrete Applied Mathematics*, 184 (2015) 202.
6. V.R. Kulli, Minus leap and square leap indices and their polynomials of some special graphs, *International Research Journal of Pure Algebra*, 8(11) (2018) 54-60.
7. V.R.Kulli, Leap hyper-Zagreb indices and their polynomials of certain graphs, *International Journal of Current Research in Life Sciences*, 7(10) (2018) 2783-2791.
8. V.R. Kulli, Sum connectivity leap index and geometric-arithmetic leap index of certain windmill graphs, *Journal of Global Research in Mathematical Archives*, 6(1) (2019).
9. V.R. Kulli, On F-leap indices and F-leap polynomials of some graphs, *International Journal of Mathematical Archive*, 9(12) (2018) 41-49.
10. V.R. Kulli, Product connectivity leap index and ABC leap index of helm graphs, *Annals of Pure and Applied Mathematics*, 18(2) (2018) 189-193.

11. V.R. Kulli, On augmented leap index and its polynomial of some wheel graphs, submitted.
12. V.R. Kulli, Computation of some topological indices of certain networks, *International Journal of Mathematical Archive*, 8(2) (2017) 99-106.
13. V.R.Kulli, Some topological indices of certain nanotubes, *Journal of Computer and Mathematical Sciences*, 8(1), (2017) 1-7.
14. L.Zhong, The harmonic index for graphs, *Appl. Math. Lett.*, 25(3) (2012) 561-566.
15. V.R. Kulli, B. Chalugaraju and H.S. Boregowda, Computation of connectivity indices of Kulli path windmill graphs, *TWMS J. Appl. Eng. Math.* 6(1) (2016) 1-8.
16. V.R.Kulli, B.Chalugaraju and H.S.Boregowda, Some degree based connectivity indices of Kulli cycle windmill graphs, *South Asian Journal of Mathematics*, 6(6) (2016) 263-268.
17. V.R.Kulli, B.Chalugaraju and H.S. Boregowda,  $K$ -Banhatti and  $K$  hyper-Banhatti indices of windmill graphs, *South East Asian J. of Math. and Math. Sci*, 13(1) (2017) 11-18.
18. J. Gallian, Graph Labeling, *Electronic J. Combin.* (2017) 1-58.

**Source of Support: Nil, Conflict of interest: None Declared**

**[Copy right © 2019, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]**