

GERAGHTY CONTRACTION PRINCIPLE FOR DIGITAL IMAGES

P. H. KRISHNA*¹ AND D. A. TATAJEE

*Department of Mathematics,
Centurion University of Technology and Management, Andhra Pradesh, India.

Department of ECE,
Visakha Technical Campus, Visakhapatnam, Andhra Pradesh, India.

(Received On: 08-08-19; Revised & Accepted On: 13-09-19)

ABSTRACT

In this paper, we prove Geraghty contraction principle for digital images. Our results extend the Banach fixed point theorem for digital images. Which is an extension of the Banach contraction principle. Example is provided to illustrate our results.

Keywords: Fixed points; Banach Contraction Principle; Geraghty contraction; Digital Image; Digital Contraction.

AMS (2010) Mathematics Subject Classification: 47H10, 54H25, 54E35, 68U10.

1. INTRODUCTION AND PRELIMINARIES

Fixed point theory deals many fields of mathematics such as Functional Analysis and general topology. Digital topology is area related to features of 2D and 3D digital images using objects with topological properties.

The Banach[1] contraction mapping theorem is well known in fixed point theory. In 1973, [14] Geraghty introduced the extension to the Banach contraction principle.

$$S = \{\beta : [0, \infty) \rightarrow [0, 1) / \beta(t_n) \rightarrow 1 \Rightarrow t_n \rightarrow 0\}.$$

Definition [14] Let (X, d) be a metric space. A self map $f : X \rightarrow X$ is said to be a *Geraghty contraction* if there exists $\beta \in S$ such that

$$d(f(x), f(y)) \leq \beta(d(x, y))d(x, y) \text{ for all } x, y \in X.$$

Theorem [14] Let (X, d) be a complete metric space. Let $f : X \rightarrow X$ be a *Geraghty contraction*. Then for any choice of initial point $x_0 \in X$, the iteration $\{x_n\}$ defined by $x_n = f(x_{n-1})$ for $n = 1, 2, 3, \dots$ converges to the unique fixed point z of f in X

Let X be a subset of Z^n for a positive integer n where Z^n is the set of lattice points in the n - Dimensional Euclidean Space and k be represent an adjacency relation for the members of X . A digital image consists (X, k) .

Definition 1.1: [5] Let k, m be positive integers, $1 \leq l \leq m$ and two distinct points $(p = p_1, p_2, p_3, \dots, p_m)$, $(q = q_1, q_2, q_3, \dots, q_m) \in Z^n$ p and q are k_l - adjacent if there are at most l indices i such that $|p_i - q_i| = 1$, and for all other indices j such that $|p_j - q_j| \neq 1, p_j = q_j$.

The following are the consequences above definition.

Two points p and q in Z are 2- adjacent if $|p - q| = 1$.




Figure-1.2: adjacent

Corresponding Author: P. H. Krishna*¹, *Department of Mathematics,
Centurion University of Technology and Management, Andhra Pradesh, India.

Two points p and q in Z^2 are 8- adjacent if they are distinct and differ by at most 1 in each coordinate.

Two points p and q in Z^2 are 4- adjacent if they are 8 - adjacent and differ in exactly one coordinate.




Figure- 1.3:

Two points p and q in Z^3 are 26- adjacent if they are distinct and differ by at most 1 in each coordinate.

Two points p and q in Z^2 are 18- adjacent if they are 26- adjacent and differ at most two coordinates.

Two points p and q in Z^2 are 6- adjacent if they are 18- adjacent and differ at most two coordinates. [5] A k - neighbor of $p \in Z^n$ is a point of Z^n that is k - adjacent to p where $k \in \{2,4,6,8,18,26\}$ and $n \in 1,2,3$.

The set $N_k(p) = \{q | q \text{ is } k\text{-adjacent to } p\}$ is called the k - neighborhood of p .

A digital interval is defined by $[a, b]_Z = \{z \in Z | a \leq z \leq b\}$, where $a, b \in Z$ and $a < b$.

A digital image $X \subset Z^n$ is k - connected [7] if and only if for every pair of different points $x, y \in X$, there is a set $\{x_0, x_1, x_2, \dots, x_r\}$ of points of a digital image X such that $x = x_0, y = x_r$ and x_i and x_{i+1} are k - neighbors where $i = 0, 1, 2, \dots, r-1$.

Definition [7]: Let $(X, k_0) \subset Z^{n_0}, (Y, k_1) \subset Z^{n_1}$ be digital images and $f: X \rightarrow Y$ be a function.

If for every k_0 - connected subset U of X , $f(U)$ is a k_1 - connected subset of Y , then f is said to be (k_0, k_1) - continuous.

f is (k_0, k_1) - continuous if and only if for every k_0 - adjacent points $\{x_0, x_1\}$ of X , either $f(x_0) = f(x_1)$ or $f(x_0)$ and $f(x_1)$ are a k_1 - adjacent in Y .

f is (k_0, k_1) - continuous, bijective and f^{-1} is (k_1, k_0) -continuous, then f is called (k_0, k_1) - isomorphism and denoted by $X \cong (k_0, k_1)^Y$.

A $(2-k)$ - continuous function $f: [0, m]_Z \rightarrow X$ such that $f(0) = x$, and $f(m) = y$ is called a digital k - path from x to y in a digital image X . In a digital image (X, k) , for every two points, if there is a k -path, then X is called k -path connected.

A simple closed k -curve of $m \geq 4$ points in a digital image X is a sequence $\{f(0), (1), f(2), \dots, f(m-1)\}$ of images of the k -path $f: [0, m-1]_Z \rightarrow X$ such that $f(i)$ and $f(j)$ are k -adjacent if and only if $j = i \pm 1 \text{ mod } m$.

A point $x \in X$ is called k -corner if x is k - adjacent to two and only two points $y, z \in X$ such that y and z are k -adjacent to each other.

If y, z are not k -corners and if x is the only point k -adjacent to both y, z then we say that the k - corner is simple. X is called a generalized simple closed k -curve if what is obtained by removing all simple k -corners of X is a simple closed k -curve.

For a k -connected digital image (X, k) in Z^n , there is a following statement

$$|X|^x = N_{3^n-1}^x(x) \cap X$$

$$k \in \{2n(n \geq 1), 3^n - 1(n \geq 2), 3^n - \sum_{t=0}^{r-2} C_t^n 2^{n-t} - 1(2 \leq r \leq n-1, n \geq 3)\},$$

where $C_t^n = \frac{n!}{(n-t)!t!}$

Definition: [18] Let (X, k) be a digital image in Z^n , $n \geq 3$ and $\bar{X} = Z^n - X$. Then X is called a closed k - surface if it satisfies the following.

If $(k, \bar{k}) \in \{(k, 2n), (2n, 3^n - 1)\}$, where the \bar{k} - adjacency is taken from (2.1) with $k \neq 3^n - 2^n - 1$ and \bar{k} is the adjacency on X , then

- For each point $x \in X$, $|X|^x$ has exactly one k -component k -adjacent to x ;
- $|\bar{X}|^x$ has exactly two \bar{k} - adjacent to xx ; we denote by C^{xx} and D^{xx} these two components; and
- For any point $y \in N_k(x) \cap X$, $N_{\bar{k}}(y) \cap C^{xx} \neq \emptyset$ and $N_{\bar{k}}(y) \cap D^{xx} \neq \emptyset$, where $N_k(x)$ means the k - neighbors of x .

Further, if a closed k -surface X does not have a simple k -point, then X is called simple.

If $(k, \bar{k}) = (3^n - 2^n - 1, 2n)$ then X is connected, for each point $x \in X$, $|X|^x$ is a generalized simple closed k - curve. If the image $|X|^x$ is a simple closed k -curve and the closed k -surface X is called simple.

Let (X, k) be a digital image and its subset be (A, k) . (X, A) is called a digital image pair with k -adjacency and when A is a singleton set $\{x_0\}$, then (X, x_0) is called a point digital image .

Banach Contraction Principle for Digital Images.

Let (X, k) be a digital image and $f: (X, k) \rightarrow (X, k)$ be any (k, k) – continuous function. We say the digital image (X, k) has the fixed point property. If for every (\bar{k}, k) – continuous map: $f: X \rightarrow X$ there exists $x \in X$ such that $f(x) = x$.

The fixed point property is preserved by any digital isomorphism. It is a topological invariant.

Let (X, d, k) be denote the digital metric space with k -adjacency where d is usual Euclidean metric for Z^n .

Definition: A sequence $\{x_n\}$ of points of digital metric space (X, d, k) is said to be a Cauchy sequence if for all $\epsilon > 0$, there exists $\alpha \in N$ such that for all $n, m > \alpha$ then $d(x_n, x_m) < \epsilon$.

Definition: A sequence $\{x_n\}$ of points of digital metric space (X, d, k) converges to a limit $a \in X$ if for all $\epsilon > 0$, there exists $\alpha \in N$ such that for all $n > \alpha$ then $d(x_n, a) < \epsilon$.

Definition: A digital metric space (X, d, k) is said to be a complete digital metric space if any Cauchy sequence $\{x_n\}$ of points (X, d, k) converges to a point a of (X, d, k) .

Definition: Let (X, k) be any digital image. A function $f: (X, k) \rightarrow (X, k)$ is called right- continuous if $f(a) = \lim_{x \rightarrow a^+} f(x)$ where $a \in X$.

Definition: [12] Let (X, d, k) be any digital metric space and $f: (X, d, k) \rightarrow (X, d, k)$ be a self digital map. If there exists $\lambda \in (0, 1)$ such that for all $x, y \in X$, $d(f(x), f(y)) \leq \lambda d(x, y)$, then f is called a digital contraction map.

Proposition: Every digital contraction map is digitally continuous.

Theorem [12]: Let (X, d, k) be a complete digital metric space which has a usual Euclidean metric in Z^n . Let $f: X \rightarrow X$ be a digital contraction map. Then f has a unique fixed point, i.e., there exists a unique $u \in X$ such that $f(u) = u$.

In 2015 Ozgur Ege, Ismet Karaca generalized Banach contraction Principle as follows.

Theorem: [12] Let (X, d, k) be a complete digital metric space which has a usual Euclidean metric d in Z^n and let $f: X \rightarrow X$ be a digital self map. Assume that there exists a right- continuous real function $\gamma: [0, u] \rightarrow [0, u]$ where u is sufficiently large real number such that $\gamma(a) < a$ if $a > 0$, and let f satisfies

$$d(f(x_1), f(x_2)) \leq \gamma(d(x_1, x_2)) \text{ for all } x_1, x_2 \in (X, d, k).$$

Then f has a unique fixed point $u \in (X, d, k)$ and the sequence $f^n(x)$ converges to u for every $x \in X$.

Now we define digital Geraghty contraction map.

Definition: Let (X, d, k) be any digital metric space and $f: (X, d, k) \rightarrow (X, d, k)$ be a self digital map is said to be a digital Geraghty contraction map if there exists $\beta \in S$ such that $d(f(x), f(y)) \leq \beta(d(x, y))d(x, y)$ for all $x, y \in X$.

Here we observe that every digital contraction map is a digital Geraghty contraction map but its converse need not be true.

Now we prove Geraghty contraction theorem in digital metric spaces.

Theorem: Let (X, d, k) be a complete digital metric space with Euclidean metric d in Z^n and let $f: X \rightarrow X$ be a digital Geraghty contraction map. Then f has a unique fixed point $u \in (X, d, k)$.

Proof: Let $x_0 \in (X, d, k)$, we define the sequence $x_n = f(x_{n-1})$ for each $n \geq 1$.

If $x_n = x_{n+1}$ for some n then x_n is a fixed point of f .

Without loss of generality, we assume that if $x_n \neq x_{n+1}$ for each n .

$$\text{We have } d(x_{n+1}, x_n) = d(f(x_n), f(x_{n-1})) \leq \beta(d(x_n, x_{n-1}))d(x_n, x_{n-1}) \quad (3.1)$$

Since $\beta \in S$,

$d(x_{n+1}, x_n) < d(x_n, x_{n-1})$, which follows that $\{d(x_{n+1}, x_n)\}$ is a decreasing sequence of non-negative reals and so $\lim_{n \rightarrow \infty} (d(x_{n+1}, x_n))$ exists and it is r (say).

Now we show that $r = 0$.

If $r > 0$ then from (3.1) we have

$$\begin{aligned} d(x_{n+1}, x_n) &= d(f(x_n), f(x_{n-1})) \leq \beta(d(x_n, x_{n-1}))d(x_n, x_{n-1}) \\ \frac{d(x_{n+1}, x_n)}{d(x_n, x_{n-1})} &\leq \beta(d(x_n, x_{n-1})) < 1 \text{ for each } n \geq 1. \end{aligned}$$

On letting $n \rightarrow \infty$, we get

$$1 = \lim_{n \rightarrow \infty} \left(\frac{d(x_{n+1}, x_n)}{d(x_n, x_{n-1})} \right) \leq \lim_{n \rightarrow \infty} \beta(d(x_n, x_{n-1})) \leq 1.$$

So that $\beta(d(x_n, x_{n-1})) \rightarrow 1$ as $n \rightarrow \infty$, that implies $\lim_{n \rightarrow \infty} d(x_{n+1}, x_n) = 0$.

Hence $r = 0$.

Let $\{x_n\}$ be a sequence in X such that $d(x_n, x_{n+1}) \rightarrow 0$ as $n \rightarrow \infty$.

Suppose that $\{x_n\}$ not a Cauchy sequence. Then there exists $\epsilon > 0$ integers

$$m(k) \text{ and } n(k) \text{ with } m(k) > n(k) > k \text{ such that } d(x_{m(k)}, x_{n(k)}) \geq \epsilon \quad (3.2)$$

we choose $m(k)$, the least positive integer satisfying $d(x_{m(k)}, x_{n(k)}) \geq \epsilon$, then we have $m(k) > n(k) > k$ with

$$\begin{aligned} d(x_{m(k)}, x_{n(k)}) &\geq \epsilon, d(x_{m(k)-1}, x_{n(k)}) < \epsilon \\ \epsilon &\leq d(x_{m(k)}, x_{n(k)}) \leq d(x_{m(k)}, x_{m(k)-1}) + d(x_{m(k)-1}, x_{n(k)}) \\ &< d(x_{m(k)}, x_{m(k)-1}) + \epsilon. \end{aligned}$$

Since $d(x_{n(k)}, x_{n(k)+1}) \rightarrow 0$ as $k \rightarrow \infty$, we have $\epsilon \leq d(x_{m(k)}, x_{n(k)}) < \epsilon$.

This implies $d(x_{m(k)}, x_{n(k)}) \rightarrow \epsilon$ as $k \rightarrow \infty$.

$$\begin{aligned} \epsilon &\leq d(x_{m(k)}, x_{n(k)}) \leq d(x_{m(k)}, x_{m(k)-1}) + d(x_{m(k)-1}, x_{n(k)+1}) + d(x_{n(k)+1}, x_{n(k)}) \\ &< d(x_{m(k)}, x_{m(k)-1}) + \epsilon + d(x_{n(k)+1}, x_{n(k)}). \end{aligned}$$

Since $d(x_{n(k)}, x_{n(k)+1}) \rightarrow 0$ as $k \rightarrow \infty$, we have $\epsilon \leq d(x_{m(k)}, x_{n(k)}) < \epsilon$.

This implies $d(x_{m(k)-1}, x_{n(k)+1}) \rightarrow \epsilon$ as $k \rightarrow \infty$.

$$\begin{aligned} \epsilon &\leq d(x_{m(k)}, x_{n(k)}) \leq d(x_{m(k)}, x_{m(k)+1}) + d(x_{m(k)+1}, x_{n(k)-1}) + d(x_{n(k)-1}, x_{n(k)}) \\ &< d(x_{m(k)}, x_{m(k)+1}) + \epsilon + d(x_{n(k)+1}, x_{n(k)}). \end{aligned}$$

Since $d(x_{m(k)}, x_{m(k)+1}) \rightarrow 0$ as $k \rightarrow \infty$, we have $\epsilon \leq d(x_{m(k)}, x_{n(k)}) < \epsilon$.

This implies $d(x_{m(k)+1}, x_{n(k)-1}) \rightarrow \epsilon$ as $k \rightarrow \infty$.

$$\begin{aligned} d(x_{m(k)+1}, x_{n(k)-1}) &= d(f(x_{m(k)}), f(x_{n(k)-1})) \leq \beta(d(x_{m(k)}, x_{n(k)-1}))d(x_{m(k)}, x_{n(k)-1}) \\ \frac{d(x_{m(k)+1}, x_{n(k)-1})}{d(x_{m(k)}, x_{n(k)-1})} &\leq \beta(d(x_{m(k)}, x_{n(k)-1})) < 1 \end{aligned}$$

On letting $k \rightarrow \infty$, we get

$$1 = \frac{\epsilon}{\epsilon} \leq \lim_{k \rightarrow \infty} \beta(d(x_{m(k)}, x_{n(k)-1})) \leq 1.$$

So that $\beta \left(d(x_{m(k)}, x_{n(k)-1}) \right) \rightarrow 1$ as $k \rightarrow \infty$,

Since $\beta \in S, d(x_{m(k)}, x_{n(k)-1}) \rightarrow 0$ as $k \rightarrow \infty$.

Hence it follows that $\epsilon = 0$, a contradiction.

Therefore $\{x_n\}$ is a Cauchy sequence in X , and since (X, d, k) is a complete digital metric space, $f^n(x)$ converges in (X, d, k) .

Geraghty contraction principle can be used to image compression of a digital image.

Example: Let F_0 be a digital image, we make a copy of F_0 and glue it on the lower left vertex.

We create a copy of F_0 and glue it on lower right vertex, so that we have a new digital image of F_1 as follows.

As in the above same procedure of F_1 , we have a new digital image F_2 which is identical to F_1 .

As a result, F_2 is the fixed point in this process.

Let T be a function which have F_1 to $T(F_1)$.

Here we have F_2 is a fixed point of this function, by continuing this process, we get an infinite sequence of sets $\{F_n\}$, then the sequence $\{F_n\}$ converges to F_2 . It cannot be distinguished F_6 from F_2 . As a result, the computer programme use F_6 instead of F_2 to be better resolution. At the same time, the programme could use F_2 in place of F_6 to determine easily some properties of digital image.

CONCLUSION

In this paper we give the digital version of Geraghty contract principle. This will be useful for digital topology and fixed point theory and to understand the better structure of digital images.

REFERENCES

1. S. Banach, Sur les operations dans les ensembles abstraits et leurs applications aux equations integrales, Fund. Math., 3 (1922), 133-181.
2. G. Bertrand, Simple points, topological numbers and geodesic neighborhoods in cubic grids, Pattern Recognition Letters, 15 (1994), 1003-1011.
3. G. Bertrand, R. Malgouyres, Some topological properties of discrete surfaces, J. Math. Imaging Vis., 20 (1999), 207-221.
4. L. Boxer, Digitally continuous functions, Pattern Recognition Letters, 15 (1994), 833-839. [5] L. Boxer, A classical construction for the digital fundamental group, J. Math. Imaging Vis., 10 (1999), 51-62.
5. L. Boxer, Properties of digital homotopy, J. Math. Imaging Vis., 22 (2005), 19-26.
6. L. Boxer, Digital products, wedges and covering spaces, J. Math. Imaging Vis., 25 (2006), 159-171.
7. L. Boxer, Continuous maps on digital simple closed curves, Appl. Math., 1 (2010), 377-386.
8. O. Ege, I. Karaca, Fundamental properties of simplicial homology groups for digital images, American Journal of Computer Technology and Application, 1 (2013), 25-42.
9. O. Ege, I. Karaca, Lefschetz Fixed Point Theorem for Digital Images, Fixed Point Theory Appl., 2013 (2013), 13 pages.
10. O. Ege, I. Karaca, Applications of the Lefschetz Number to Digital Images, Bull. Belg. Math. Soc. Simon Stevin, 21 (2014), 823-839.
11. Ozgur Ege, Ismet Karaca, Banach fixed point theorem for digital images, Journal of Non linear Science and Applications, 8(2015), 237-245.
12. G. V. R. Babu, K.K. M. Sarma and P. H. Krishna. Fixed points of ψ – weak Geraghty contractions in partially ordered metric spaces. Journal of Advanced research in pure mathematics. 6(4), 2014, 9 - 23.
13. M. A. Geraghty. On contractive maps. Proc. of Amer. Math. Soc., 1973, 40:604-608.
14. S. E. Han, An extended digital $(k_0; k_1)$ -continuity, J. Appl. Math. Comput, 16 (2004), 445-452.
15. S. E. Han, Minimal simple closed 18-surfaces and a topological preservation of 3D surfaces, Inform. Sci., 176(2006), 120{134. 2.4O. Ege, I. Karaca, J. Nonlinear Sci. Appl. 8 (2015), 237-245.
16. S. E. Han, Connected sum of digital closed surfaces, Inform. Sci., 176 (2006), 332-348.
17. G. T. Herman, Oriented surfaces in digital spaces, CVGIP: Graphical Models and Image Processing, 55 (1993), 381-396.
18. M. Jleli, B. Samet, A new generalization of the Banach contraction principle, J. Inequal. Appl., 2014 (2014), 8 pages.

19. I. Karaca, O. Ege, Some results on simplicial homology groups of 2D digital images, *Int. J. Inform. Computer Sci.*, 1 (2012), 198-203.
20. T. Y. Kong, A digital fundamental group, *Computers and Graphics*, 13 (1989), 159-166.
21. R. Malgouyres, G. Bertrand, A new local property of strong n-surfaces, *Pattern Recognition Letters*, 20 (1999), 417-428.
22. A. Rosenfeld, Digital topology, *Amer. Math. Monthly*, 86 (1979), 76-87.
23. W. Shatanawi, H. K. Nashine, A generalization of Banach's contraction principle for nonlinear contraction in a partial metric space, *J. Nonlinear Sci. Appl.*, 5 (2012), 37-43.

Source of Support: Nil, Conflict of interest: None Declared

[Copy right © 2019, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]