

COMMON FIXED POINT THEOREMS OF  
GERAGHTY TYPE CONTRACTION MAPS IN RECTANGULAR METRIC SPACES

P. HARIKRISHNA\*<sup>1</sup>, R. SANTHI KUMAR<sup>2</sup>

<sup>1</sup>Department of Mathematics,  
Centurion University of Technology and Management, Andhra Pradesh, India.

<sup>2</sup>Department of Mathematics,  
AITAM College of Engineering, Tekkali, Andhra Pradesh, India.

(Received On: 30-06-2020; Revised & Accepted On: 11-07-2020)

---

ABSTRACT

In this paper, we define generalized Geraghty contraction maps in rectangular metric spaces with an altering distance function for a pair of maps, and prove the existence of common fixed points. Our results extend some of the known results and provided example in support of main theorem.

**Keywords:** Common fixed point; rectangular metric spaces; Geraghty contraction; weakly compatible.

**AMS (2010) Mathematics Subject Classification:** 47H10, 54H25.

---

1. INTRODUCTION AND PRELIMINARIES

Fixed point theory is very important tool in Non linear analysis. Banach contraction principle is one of the fundamental results in fixed point theory. There are several generalizations of metric spaces. In 1973, Geraghty [10] extended the Banach contraction theorem by replacing the contraction by a function with specific properties and proved the existence of fixed points, In 1984 Khan, Swalwh and Sessa [14] studied fixed points with altering distance functions. In 2000 Branciari [4] generalized metric spaces, in which triangular inequality is replaced by quadrilateral inequality which is known as rectangular metric spaces. In such extensions some of the authors are focused on rectangular metric spaces and proved the existence of fixed and common fixed points. We refer [2, 4, 6, 13, and 15].

**Definition 1.1:** [4] Let  $X$  be a nonempty set. A function  $d: X \times X \rightarrow [0, \infty)$  satisfy the following conditions for all  $x, y \in X$  and all distinct  $u, v \in X$  each of them different from  $x$  and  $y$

- (i)  $d(x, y) = 0$  if and only if  $x = y$
- (ii)  $d(x, y) = d(y, x)$ , and
- (iii)  $d(x, y) \leq d(x, u) + d(u, v) + d(v, y)$ . (rectangular inequality)

Then the function  $d$  is called a rectangular (generalized) metric and the pair  $(X, d)$  is called a rectangular (generalized) metric space (in short RMS).

**Definition 1.2:** [4] Let  $(X, d)$  be a rectangular metric space (in short RMS) and  $\{x_n\}$  be a sequence in  $X$ .

- (i)  $\{x_n\}$  is called (RMS) convergent to  $x$  in  $X$  if and only if  $d(x_n, x) \rightarrow 0$  as  $n \rightarrow \infty$ .
- (ii)  $\{x_n\}$  is called (RMS) Cauchy sequence if and only if for every  $\epsilon > 0$  there exists positive integer  $N(\epsilon)$  such that  $d(x_n, x_m) < \epsilon$  for all  $m > n \geq N(\epsilon)$ .

A rectangular metric space  $(X, d)$  is called complete if every (RMS) Cauchy sequence is a (g.m.s) convergent.

**Definition 1.3:** ([14]) A function  $\psi: R^+ \rightarrow R^+$ ,  $R^+ = [0, \infty)$  is said to be an *altering distance function* if the following conditions hold:

- (i)  $\Psi$  is continuous,
- (ii)  $\psi$  is non-decreasing, and
- (iii)  $\psi(t) = 0$  if and only if  $t = 0$ .

---

**Corresponding Author:** P. HariKrishna\*<sup>1</sup>,

<sup>1</sup>Department of Mathematics,

Centurion University of Technology and Management, Andhra Pradesh, India.

In 1973, [10] Geraghty extended Banach contraction theorem by replacing the contraction constant by a function with specific properties.

$$S = \{ \beta: [0, \infty) \rightarrow [0, 1) / \beta(t_n) \rightarrow 1 \Rightarrow t_n \rightarrow 0 \}.$$

**Definition 1.4:** [10, 16] Let  $(X, d)$  be a metric space. A self map  $f: X \rightarrow X$  is said to be a Geraghty contraction if there exists  $\beta \in S$  such that  $d(f(x), f(y)) \leq \beta(d(x, y))d(x, y)$  for all  $x, y \in X$ .

**Theorem 1.5:** [10] Let  $(X, d)$  be a complete metric space. Let  $f: X \rightarrow X$  be a Geraghty contraction.

Then for any choice of initial point  $x_0 \in X$ , the iteration  $\{x_n\}$  defined by  $x_n = f(x_{n-1})$  for  $n = 1, 2, 3, \dots$  converges to the unique fixed point  $z$  of  $f$  in  $X$ .

In 2019 P.H. Krishna [15] *et.al* proved the following fixed theorem in rectangular maps with admissible maps of Geraghty type contraction condition.

**Definition 1.6:** [15] Let  $(X, d)$  be a rectangular metric space and let  $T: X \rightarrow X$  be a self map. If there exists  $\beta \in S$  such that  $d(Tx, Ty) \leq \beta(\varphi(M(x, y)))\varphi(M(x, y))$

Where  $M(x, y) = \max\{d(x, y), d(x, Tx), d(y, Ty), \frac{1}{1+d(x, y)}[d(x, Tx)d(y, Ty)], \frac{1}{1+d(Tx, Ty)}[d(x, Tx)d(y, Ty)]\}$  for all  $x, y \in X$  then we call  $T$  is a  $\varphi_M$ -generalized Geraghty contraction in rectangular metric spaces.

**Theorem 1.7:** [15] Let  $(X, d)$  be a Hausdorff and complete rectangular metric space.

Let  $T: X \rightarrow X$  be an  $\alpha$ -admissible mapping with respect to  $\eta$ . Assume that there exists an altering distance function  $\varphi$  such that  $x, y \in X, \alpha(x, y) \geq \eta(x, y)$ , implies  $d(x, y) \leq \beta(\varphi(M(x, y)))\varphi(M(x, y))$  (2.1.1)

where  $M(x, y) = \max\{d(x, y), d(x, Tx), d(y, Ty), \frac{1}{1+d(x, y)}[d(x, Tx)d(y, Ty)], \frac{1}{1+d(Tx, Ty)}[d(x, Tx)d(y, Ty)]\}$

Also, suppose that the following assertions are hold;

- (i) there exists  $x_0 \in X$  such that  $\alpha(x_0, Tx_0) \geq \eta(x_0, Tx_0)$
- (ii) for all  $x, y \in X, \alpha(x, y) \geq \eta(x, y)$  and  $\alpha(y, z) \geq \eta(y, z)$  implies  $\alpha(x, z) \geq \eta(x, z)$
- (iii)  $T$  is continuous.

Then  $T$  has a periodic point  $a \in X$  and  $\alpha(x, Ta) \geq \eta(a, Ta)$  holds for each periodic point then  $T$  has a fixed point.

**Definition 1.8:** [12] Let  $f$  and  $g$  be self-mappings of a nonempty set  $X$ .

A point  $x$  in  $X$  is said to be common fixed point of  $f$  and  $g$  if  $x = fx = gx$ .

A point  $x$  in  $X$  is said to be coincidence point of  $f$  and  $g$  if  $fx = gx$ . And if  $u = fx = gx$ , then  $u$  is said to be a point of coincidence of  $f$  and  $g$ .

The mappings  $f, g: X \rightarrow X$  are said to be weakly compatible if they commute at their coincidence points. i.e.,  $fgx = gfx$  whenever  $fx = gx$ .

**Definition 1.9:** [12] Two self mappings  $S$  and  $T$  of a rectangular metric space  $(X, d)$  are said to be compatible if  $d(S(Tx_n), T(Sx_n)) = 0$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_{n \rightarrow \infty} S(x_n) = \lim_{n \rightarrow \infty} T(x_n) = u$  for some  $u \in X$ .

**Definition 1.10:** Suppose that  $(X, d, k)$  be a rectangular metric spaces and  $S, T: X \rightarrow X$  be a map defined on  $X$ . Then  $S$  and  $T$  are said to be weakly commutative iff  $d(S(T(x)), T(S(x))) \leq d(S(x), T(x))$  for all  $x$  in  $X$ . 2013, Muhammad Arsal *et.al* [13] proved common fixed point theorem on Hausdorff rectangular metric spaces.

**Theorem 1.11:** [13] Let  $(X, d)$  be a Hausdorff rectangular metric space and let  $F, g: X \rightarrow X$  be self mappings such that  $Fx \subset g(X)$ . Assume that  $(gX, d)$  is a complete rectangular metric space. Suppose that the following conditions hold.

$$\psi(d(F(x), F(y))) \leq (\psi(M(g(x), g(y))) - \Phi(M(g(x), g(y))))$$
 for all  $x, y$  in  $X$  and  $\psi, \Phi$  in  $\Psi$ ,

where  $\psi$  is non decreasing and  $M(g(x), g(y)) = \max\{d(g(x), g(y)), d(g(x), f(x)), d(g(y), F(y))\}$ . Then  $F$  and  $g$  have a unique coincident point in  $X$ . Moreover, if  $F$  and  $g$  are weakly compatible, then  $F$  and  $g$  have a unique common fixed point.

The following Lemma is useful to prove the Cauchy's sequence.

**Lemma 1.12:** [7] Let  $(X, d)$  be metric space. Let  $\{x_n\}$  be a sequence in  $X$  such that  $d(x_{n+1}, x_n) \rightarrow 0$  as  $n \rightarrow \infty$ . If  $\{x_n\}$  is not a Cauchy sequence then there exist an  $\epsilon > 0$  and sequences of positive integers  $\{m(k)\}$  and  $\{n(k)\}$  with  $n(k) > m(k) > k$  and  $d(x_{m(k)}, x_{n(k)}) \geq \epsilon$ . For each  $k > 0$ , corresponding to  $m(k)$ , we can choose  $n(k)$  to be the smallest integer such that  $d(x_{m(k)}, x_{n(k)}) \geq \epsilon$  and  $d(x_{m(k)}, x_{n(k)-1}) < \epsilon$ . It can be shown that the following identities are satisfied.

- (i)  $\lim_{k \rightarrow \infty} d(x_{n(k)}, x_{m(k)}) = \epsilon$
- (ii)  $\lim_{k \rightarrow \infty} d(x_{n(k)-1}, x_{m(k)+1}) = \epsilon$ ,
- (iii)  $\lim_{k \rightarrow \infty} d(x_{n(k)-1}, x_{m(k)}) = \epsilon$ , and (iv)  $\lim_{k \rightarrow \infty} d(x_{n(k)}, x_{m(k)+1}) = \epsilon$ .

Now, we prove the existence of common fixed points of generalized Geraghty contraction maps with rectangular metric spaces for a pair of maps.

### MAIN RESULTS

Now we define Geraghty contraction for a pair of maps in Hausdorff rectangular metric spaces.

**Definition 2.1:** Let  $(X, d)$  be a Hausdorff rectangular metric space. Let  $T$  and  $S$  be self maps on  $X$ . If there exists  $\beta \in S$  such that

$$d(S(x), S(y)) \leq \beta(d(T(x), T(y))(d(T(x), T(y))) \tag{2.1.1}$$

for all  $x, y \in X$ , then we say that  $(T, S)$  is a pair of Geraghty contraction maps in Hausdorff rectangular metric spaces.

**Definition 2.2:** Let  $(X, d)$  be a Hausdorff rectangular metric space. Let  $T$  and  $S$  be self maps on  $X$ . If there exists  $\beta \in S$  such that,

$$d(S(x), S(y)) \leq \beta(m(T(x), T(y)))(m(T(x), T(y))) \tag{2.2.1}$$

Where  $m(T(x), T(y)) = \max\{d(Tx, Ty), d(Tx, Sx), d(Ty, Sy), [\frac{d(Tx, Sy)+d(Ty, Sx)}{2}]\}$  for all  $x, y \in X$ , then we say that  $(T, S)$  is a pair of Generalized Geraghty contraction maps in Hausdorff rectangular metric spaces.

**Definition 2.3:** Let  $(X, d)$  be a Hausdorff rectangular metric space. Let  $T$  and  $S$  be self maps on  $X$ . If there exists  $\beta \in S, \psi \in \psi$  such that

$$\psi(d(S(x), S(y))) \leq \beta(\psi(m(T(x), T(y)))\psi(m(T(x), T(y)))) \tag{2.3.1}$$

Where  $m(T(x), T(y)) = \max\{d(Tx, Ty), d(Tx, Sx), d(Ty, Sy), [\frac{d(Tx, Sy)+d(Ty, Sx)}{2}]\}$  for all  $x, y \in X$ , then we say that  $(T, S)$  is a pair of  $\psi$ -Generalized Geraghty contraction maps in Hausdorff rectangular metric spaces.

Now we prove the existence of common fixed points of Geraghty contraction maps in pair of maps in  $\psi$ -Generalized Geraghty contraction maps in Hausdorff rectangular metric spaces.

**Theorem 2.4:** Let  $(X, d)$  be a Hausdorff rectangular metric space and let  $S, T: X \rightarrow X$  be selfmaps such that  $S(X) \subset T(X)$ . Assume that  $(Tx, d)$  be a complete rectangular metric space. Suppose that  $(T, S)$  is a pair of  $\psi$ -Generalized Geraghty contraction maps. Then  $S$  and  $T$  have a unique coincidence point in  $X$ . Moreover, if  $S$  and  $T$  are weakly compatible, then  $S$  and  $T$  have a unique common fixed point.

**Proof:** Let  $x_0 \in (X, d)$ ,

Since  $S(X) \subset T(X)$ , we define the sequence  $Tx_n = S(x_{n-1})$  for each  $n \geq 1$ .

If  $Tx_{n+1} = Tx_{n+2}$  for some  $n$ , then  $Tx_{n+1} = Sx_{n+1}$  and hence  $x_{n+1}$  is a coincident point of  $T$  and  $S$ . Without loss of generality, we assume that if  $Tx_{n+1} \neq Tx_{n+2}$  for each  $n$ , then we have  $d(Tx_{n+2}, Tx_{n+1}) > 0$

We consider

$$\Psi(d(Tx_{n+2}, Tx_{n+1})) = \psi(d(Sx_{n+1}, Sx_n)) \leq \beta(\psi(m(Tx_{n+1}, Tx_n)))\psi(m(Tx_{n+1}, Tx_n)) \tag{2.4.1}$$

$$\begin{aligned} m(Tx_{n+1}, Tx_n) &= \max\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Sx_{n+1}), d(Tx_n, Sx_n), [\frac{d(Tx_{n+1}, Sx_n)+d(Tx_n, Sx_{n+1})}{2}]\} \\ &= \max\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Tx_{n+2}), d(Tx_n, Tx_{n+1}), [\frac{d(Tx_{n+1}, Tx_{n+1}) + d(Tx_n, Tx_{n+2})}{2}]\} \\ &= \max\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Tx_{n+2}), \frac{d(Tx_n, Tx_{n+2})}{2}\} \\ &\leq \max\left\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Tx_{n+2}), \frac{d(Tx_n, Tx_{n+1}) + d(Tx_{n+1}, Tx_{n+2})}{2}\right\} \\ &\leq \max\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Tx_{n+2})\}. \end{aligned}$$

If  $\max\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Tx_{n+2})\} = d(Tx_{n+1}, Tx_{n+2})$

Since  $\beta \in S$  then 2.4.1 implies

$$\Psi(d(Tx_{n+2}, Tx_{n+1})) < \Psi(d(Tx_{n+2}, Tx_{n+1})), \text{ Which is a contradiction.}$$

Therefore  $\max\{d(Tx_{n+1}, Tx_n), d(Tx_{n+1}, Tx_{n+2})\} = d(Tx_{n+1}, Tx_n)$

Now  $\Psi(d(Tx_{n+2}, Tx_{n+1})) = \psi(d(Sx_{n+1}, Sx_n)) \leq \beta(\psi(m(Tx_{n+1}, Tx_n)))\psi(d(Tx_{n+1}, Tx_n))$

Since  $\beta \in S$ ,

$$\psi(d(Tx_{n+2}, Tx_{n+1})) < \psi(d(Tx_{n+1}, Tx_n)) \quad (2.4.2)$$

Which follows that  $\{\psi(d(Tx_{n+2}, Tx_{n+1}))\}$  is a decreasing sequence of non-negative reals and since  $\psi$  is continuous, it follows that  $\{d(Tx_{n+2}, Tx_{n+1})\}$  is also a decreasing sequence of non-negative reals so that  $\lim_{n \rightarrow \infty} d(Tx_{n+2}, Tx_{n+1})$  exists and it is  $r$  (say).

Now we show that  $r = 0$ .

If  $r > 0$  then from (2.4.1) we have

$$\begin{aligned} \psi(d(Tx_{n+2}, Tx_{n+1})) &= \psi(d(Sx_{n+1}, Sx_n)) \leq \beta(\psi(m(Tx_{n+1}, Tx_n)))\psi(m(Tx_{n+1}, Tx_n)) \\ \frac{\psi(d(Tx_{n+2}, Tx_{n+1}))}{\psi(d(Tx_{n+1}, Tx_n))} &\leq \beta(\psi(m(Tx_{n+1}, Tx_n))) < 1 \text{ for each } n \geq 1. \end{aligned}$$

On letting  $n \rightarrow \infty$ , we get

$$1 = \lim_{n \rightarrow \infty} \frac{\psi(d(Tx_{n+2}, Tx_{n+1}))}{\psi(d(Tx_{n+1}, Tx_n))} \leq \lim_{n \rightarrow \infty} \beta(\psi(m(Tx_{n+1}, Tx_n))) \leq 1.$$

So that  $\beta(\psi(m(Tx_{n+1}, Tx_n))) \rightarrow 1$  as  $n \rightarrow \infty$ , that implies

$$\lim_{n \rightarrow \infty} \psi(d(Tx_{n+1}, Tx_n)) = 0. \quad (2.4.3)$$

Hence  $r = 0$ .

Let  $\{x_n\}$  be a sequence in  $X$  such that  $d(Tx_{n+1}, Tx_n) \rightarrow 0$  as  $n \rightarrow \infty$ .

Suppose that  $\{Tx_n\}$  not a Cauchy sequence. Then there exists  $\epsilon > 0$  integers  $m(k)$  and  $n(k)$  with  $m(k) > n(k) > k$  such that  $d(Tx_{m(k)}, Tx_{n(k)}) \geq \epsilon$  (2.4.4)

We choose  $m(k)$ , the least positive integer satisfying  $d(Tx_{m(k)}, Tx_{n(k)}) \geq \epsilon$ , then

$$\begin{aligned} \text{we have } m(k) > n(k) > k \text{ with } d(Tx_{m(k)}, Tx_{n(k)}) &\geq \epsilon, d(Tx_{m(k)-1}, Tx_{n(k)}) < \epsilon \\ \epsilon &\leq d(Tx_{m(k)}, Tx_{n(k)}) \leq d(Tx_{m(k)}, Tx_{m(k)-1}) + d(Tx_{m(k)-1}, Tx_{n(k)}) \\ &< d(Tx_{m(k)}, Tx_{m(k)-1}) + \epsilon. \end{aligned}$$

Since  $d(Tx_{m(k)}, Tx_{n(k)+1}) \rightarrow 0$  as  $k \rightarrow \infty$ , we have  $\epsilon \leq d(Tx_{m(k)}, Tx_{n(k)}) < \epsilon$ .

This implies  $d(Tx_{m(k)}, Tx_{n(k)}) \rightarrow \epsilon$  as  $k \rightarrow \infty$ .

$$\begin{aligned} \epsilon &\leq d(Tx_{m(k)}, Tx_{n(k)}) \leq d(Tx_{m(k)}, Tx_{m(k)-1}) + d(Tx_{m(k)-1}, Tx_{n(k)+1}) + d(Tx_{n(k)+1}, Tx_{n(k)}) \\ &< d(Tx_{m(k)}, Tx_{m(k)-1}) + \epsilon + d(Tx_{n(k)+1}, Tx_{n(k)}). \end{aligned}$$

Since  $d(Tx_{n(k)}, Tx_{n(k)+1}) \rightarrow 0$  as  $k \rightarrow \infty$ , we have  $\epsilon \leq d(Tx_{m(k)}, Tx_{n(k)}) < \epsilon$

This implies  $d(Tx_{m(k)-1}, Tx_{n(k)+1}) \rightarrow \epsilon$  as  $k \rightarrow \infty$ .

$$\begin{aligned} \epsilon &\leq d(Tx_{m(k)}, Tx_{n(k)}) \leq d(Tx_{m(k)}, Tx_{m(k)+1}) + d(Tx_{m(k)+1}, Tx_{n(k)-1}) + d(Tx_{n(k)-1}, Tx_{n(k)}) \\ &< d(Tx_{m(k)}, Tx_{m(k)+1}) + \epsilon + d(Tx_{n(k)+1}, Tx_{n(k)}). \end{aligned}$$

Since  $d(Tx_{m(k)}, Tx_{m(k)+1}) \rightarrow 0$  as  $k \rightarrow \infty$ , we have  $\epsilon \leq d(Tx_{m(k)}, Tx_{n(k)}) < \epsilon$

This implies  $d(Tx_{m(k)+1}, Tx_{n(k)-1}) \rightarrow \epsilon$  as  $k \rightarrow \infty$ .

Now, we consider

$$\begin{aligned} \psi(d(Tx_{m(k)+1}, Tx_{n(k)})) &= \psi(d(S(x_{m(k)}), S(x_{n(k)-1}))) \\ &\leq \beta \left( \psi \left( m(Tx_{m(k)}, Tx_{n(k)-1}) \right) \right) \psi \left( m(Tx_{m(k)}, Tx_{n(k)-1}) \right) \end{aligned} \quad (2.4.5)$$

$$\begin{aligned} m(Tx_{m(k)}, Tx_{n(k)-1}) &= \max\{d(Tx_{m(k)}, Tx_{n(k)}), (Tx_{m(k)}, Sx_{m(k)}), d(Tx_{n(k)}, Sx_{n(k)}), \\ &\quad 1/2[d(Tx_{m(k)}, Sx_{n(k)}) + d(Tx_{n(k)}, Sx_{m(k)})]\} \\ &= \max\{d(Tx_{m(k)}, Tx_{n(k)}), (Tx_{m(k)}, Tx_{m(k)+1}), d(Tx_{n(k)}, Tx_{n(k)+1}), \\ &\quad 1/2[d(Tx_{m(k)}, Tx_{n(k)+1}) + d(Tx_{n(k)}, Tx_{m(k)+1})]\} \end{aligned}$$

On letting  $k \rightarrow \infty$ , and by Lemma (1.12), it follows that

$$\lim_{k \rightarrow \infty} m(Tx_{m(k)}, Tx_{n(k)-1}) = \max\{\epsilon, 0, 0, 1/2[\epsilon + \epsilon]\} = \epsilon.$$

$$\frac{d(Tx_{m(k)+1}, Tx_{n(k)})}{d(Tx_{m(k)}, Tx_{n(k)-1})} \leq \beta \left( d(Tx_{m(k)}, Tx_{n(k)-1}) \right) < 1$$

Therefore  $\psi(\epsilon) \leq \lim_{k \rightarrow \infty} \beta \left( \psi(m(Tx_{m(k)}, Tx_{n(k)-1})) \right) \cdot \psi(\epsilon)$

$$1 = \frac{\psi(\epsilon)}{\psi(\epsilon)} \leq \lim_{k \rightarrow \infty} \beta \left( \psi(m(Tx_{m(k)}, Tx_{n(k)-1})) \right) \leq 1.$$

So that  $\beta \left( \psi(m(Tx_{m(k)}, Tx_{n(k)-1})) \right) \rightarrow 1$  as  $k \rightarrow \infty$ ,

Since  $\beta \in S$ ,  $\psi(m(Tx_{m(k)}, Tx_{n(k)-1})) \rightarrow 0$  as  $k \rightarrow \infty$ .

Since  $\psi$  is continuous, Hence it follows that  $\epsilon = 0$ , a contradiction.

Therefore  $\{Tx_n\}$  is a Cauchy sequence in  $X$ , and since  $(TX, d)$  is a complete Hausdorff rectangular metric space, there exists  $u \in Tx$  such that  $Tx_{n+1} = Sx_n \rightarrow u$  as  $n \rightarrow \infty$ .

Now let  $y \in X$  such that  $Ty = u$ .

Suppose that if  $Sy \neq Ty$  i.e.,  $d(Sy, Ty) > 0$ .

Since  $\{Tx_n\}$  is a decreasing sequence of non-negative reals and  $\{Tx_n\}$  converges to  $Tu$  for some  $u \in X$ .

From the inequality (2.4.1), with  $x = x_n$ , we get

$$\psi(d(Sx_n, Sy)) \leq \beta \left( \psi(m(Tx_n, Ty)) \right) \psi(m(Tx_n, Ty)) \tag{2.4.6}$$

where

$$m(Tx_n, Ty) = \max\{ (Tx_n, Ty), (Tx_n, Sx_n), (Ty, Sy), 1/2[(Tx_n, Sy) + (Ty, Sx_n)] \}$$

$$= \max\{ (Tx_n, Ty), (Tx_n, Tx_{n+1}), (Ty, Sy), 1/2[(Tx_n, Sy) + (Ty, Tx_{n+1})] \}$$

On letting  $n \rightarrow \infty$

$$= \max\{ (u, Ty), (u, u), (Ty, Sy), 1/2[(Ty, Sy) + (Ty, u)] \}$$

$$= \max\{ (u, u), (u, u), (Ty, Sy), 1/2[(Ty, Sy) + (u, u)] \}$$

$$= d(Ty, Sy)$$

Taking limit as  $n \rightarrow \infty$  in (2.4.6)

$$\lim_{n \rightarrow \infty} \psi(d(Sx_n, Sy)) \leq \lim_{n \rightarrow \infty} \beta \left( \psi(m(Tx_n, Ty)) \right) \psi(m(Tx_n, Ty))$$

$$\lim_{n \rightarrow \infty} \psi(d(Sx_n, Sy)) \leq \lim_{n \rightarrow \infty} \beta \left( \psi(m(Tx_n, Ty)) \right) \psi(d(Ty, Sy))$$

$$\lim_{n \rightarrow \infty} d(Sx_n, Sy) = 0$$

$$d(Ty, Sy) = 0$$

Therefore  $Ty = Sy$  and hence  $y$  is a coincidence point of  $T$  and  $S$ .

Since  $T(X)$  is complete, there exists  $u \in T(X)$  such that  $\lim_{n \rightarrow \infty} Tx_{n+1} = \lim_{n \rightarrow \infty} Sx_n = Tu = u$  for some  $y \in X$ .

Suppose that if  $Sy \neq Ty$  i.e.,  $d(Sy, Ty) > 0$

We get Since  $S$  and are weakly commutative

$$\psi(d(T(Sx_n), S(Tx_n))) \leq \psi(d(Tx_n), Sx_n)$$

Which implies  $\psi(d(T(u), S(u))) \leq \psi(d(u, u))$

Therefore  $T(u) = S(u)$  and hence  $u$  is a coincidence point of  $S$  and  $T$ .

So that  $T(S(u)) = S(T(u)) = S(S(u))$

Suppose that  $\psi(d(S(u), S(S(u)))) > 0$

$$\psi(d(S(u), S(S(u)))) \leq \beta \left( \psi(m(T(u), T(S(u)))) \psi(m(T(u), T(S(u)))) \right)$$

Since  $\beta$  in  $S$ , it follows that

$$\psi(d(S(u), S(S(u)))) < \psi(d(T(u), T(S(u))))$$

$$\psi(d(S(u), S(S(u)))) < \psi(d(S(u), S(S(u))))$$
, which is contradiction

so that  $\psi(d(S(u), S(S(u)))) = 0$

Since

$\psi$  is continuous, it follows that  $d(S(u), S(S(u))) = 0$  and hence  $S(u) = S(S(u))$  so that  $S(u)$  is a fixed point of  $S$ .  $S(u)$  is common fixed point of  $S$  and  $T$ .

**Theorem 2.5:** Let  $(X, d)$  be a Hausdorff rectangular metric space and let  $S, T: X \rightarrow X$  be selfmaps such that  $S(X) \subset T(X)$ . Assume that  $(T(X), d)$  be a complete rectangular metric space. Suppose that  $(T, S)$  is a pair of  $\psi$ -Generalized Geraghty contraction maps. Then  $S$  and  $T$  have a unique coincidence point in  $X$ . Moreover, if  $S$  and  $T$  commute, then  $S$  and  $T$  have a unique common fixed point.

**Proof:** As in the proof of the theorem  $\{T(x_n)\}$  is a Cauchy sequence in  $X$ , and since  $(X, d)$  is a Hausdorff rectangular metric space,  $\{S(x_n)\}$  converges in  $(X, d)$ . Since  $T(X)$  is complete, there exists  $u \in T(X)$  such that

$$\lim_{n \rightarrow \infty} \{T(x_{n+1})\} = \lim_{n \rightarrow \infty} \{S(x_n)\} = Tu = u \text{ for some } y \in X. \text{ Suppose that if } Sy \neq Ty$$

i.e.,  $d(Sy, Ty) > 0$

Since  $S$  and  $T$  commutes so that  $S(T(x_n)) = T(S(x_n))$  for all  $n$ .

Thus  $S(u) = T(u)$ , and consequently by commutativity,  $T(T(u)) = T(S(u)) = S(S(u))$ .

So that  $T(S(u)) = S(T(u)) = S(S(u))$ .

Suppose that  $d(S(u), S(S(u))) > 0$

$$d(S(u), S(S(u))) \leq \beta(d(T(u), T(S(u)))d(T(u), T(S(u))))$$

Since  $\beta$  in  $S$ , it follows that

$$\psi(d(S(u), S(S(u)))) < \psi(d(T(u), T(S(u))))$$

$$\psi(d(S(u), S(S(u)))) < \psi(d(S(u), S(S(u))))$$
, which is contradiction

so that  $\psi(d(S(u), S(S(u)))) = 0$  and hence  $S(u) = S(S(u))$  so that  $S(u)$  is a fixed point of  $S$ .  $S(u)$  is common fixed point of  $S$  and  $T$ .

The following is the Example in support of Theorem 2.4.

**Example 2.5:** Let  $X = A \cup B$ , where  $A = \{\frac{1}{2}, \frac{2}{3}, \frac{3}{4}\}$  and  $B = [2, 3]$ . We define the generalized metric  $d$  on  $X$  such that

$$d\left(\frac{1}{2}, \frac{2}{3}\right) = 0.2, \quad d\left(\frac{1}{2}, \frac{3}{4}\right) = 0.6, \quad d\left(\frac{2}{3}, \frac{3}{4}\right) = 0.3,$$

$$d\left(\frac{1}{2}, \frac{1}{2}\right) = d\left(\frac{2}{3}, \frac{2}{3}\right) = d\left(\frac{3}{4}, \frac{3}{4}\right) = 0.$$

$$d(x, y) = |x - y|, \text{ if } x, y \in B \text{ or } x \in A.$$

We define  $S: X \rightarrow X$  by  $\begin{cases} \frac{2}{3} & \text{if } x \in [2, 3] \\ \frac{1}{2} & \text{if } x \in \{\frac{1}{2}, \frac{2}{3}, \frac{3}{4}\} \end{cases}$  and  $T: X \rightarrow X$  by  $\begin{cases} \frac{3}{4} & \text{if } x \in [2, 3] \\ \frac{1}{2} & \text{if } x = \frac{1}{2} \\ \frac{2}{3} & \text{if } x \in \{\frac{2}{3}, \frac{3}{4}\} \end{cases}$

We define  $\beta: [0, \infty) \rightarrow [0, 1)$  by  $\beta(t) = \begin{cases} 0 & \text{if } t = 0 \\ \frac{1}{1+t} & \text{if } t > 0 \end{cases}$

And  $\psi(t) = t^2$  for all  $t \geq 0$ .

Now we verify the inequality (2.3.10) in the following cases:

**Case-(i):**  $x \in [2, 3]$  and  $y = \frac{1}{2}$

$$\psi(d(S(x), S(y))) = \psi\left(d\left(\frac{2}{3}, \frac{1}{2}\right)\right) = \psi(0.2) = 0.04$$

$$m(T(x), T(y)) = \max\left\{d\left(\frac{3}{4}, \frac{1}{2}\right), d\left(\frac{3}{4}, \frac{2}{3}\right), d\left(\frac{1}{2}, \frac{1}{2}\right), \left[\frac{d\left(\frac{2}{3}, \frac{1}{2}\right) + d\left(\frac{3}{4}, \frac{1}{2}\right)}{2}\right]\right\}$$

$$= \max\{0.6, 0.3, 0, (0.2+0.6)/2\} = 0.6$$

$$\beta(\psi(0.6))\psi(0.6) = \beta(0.36)(0.36) = \frac{0.36}{1.36} = 0.264$$

Therefore,  $0.04 = \psi(d(S(x), S(y))) \leq 0.264 = \beta(\psi(m(T(x), T(y))))\psi(m(T(x), T(y)))$

**Case-(ii):**  $x \in [2, 3]$  and  $y \in \{\frac{2}{3}, \frac{3}{4}\}$

$$\psi(d(S(x), S(y))) = \psi\left(d\left(\frac{2}{3}, \frac{1}{2}\right)\right) = \psi(0.2) = 0.04$$

$$m(T(x), T(y)) = \max\left\{d\left(\frac{3}{4}, \frac{2}{3}\right), d\left(\frac{3}{4}, \frac{2}{3}\right), d\left(\frac{2}{3}, \frac{1}{2}\right), \left[\frac{d\left(\frac{2}{3}, \frac{2}{3}\right) + d\left(\frac{3}{4}, \frac{1}{2}\right)}{2}\right]\right\}$$

$$= \max\{0.3, 0.3, 0.2, (0+0.6)/2\} = 0.3$$

$$(\psi(0.3)\psi(0.3)) = \beta(0.09)(0.09) = \frac{0.09}{1.09} = 0.0825$$

$0.04 = \psi(d(S(x), S(y))) \leq 0.0825 = \beta(\psi(m(T(x), T(y))))\psi(m(T(x), T(y)))$

And the remaining cases also the inequality (2.3.1) holds.

Therefore  $S$  and  $T$  have the unique common fixed point  $\frac{1}{2}$ .

## REFERENCES

1. S. Banach, Sur les operations dans les ensembles abstraitsetleurs applications aux equations integrales, Fund.Math., 3 (1922), 133-181.
2. Arslan Hojat Ansari, Haseen Aydi, P Sumati Kumari and Isa Yildirim, New fixed point results via c-class functions in b- rectangular metric spaces. Communications in Mathematics and Applications, Vol. 9, No.2, pp 109-126, 2018.
3. D. Baleanu, Sh. Rezapour and M. Mohammadi, Some existence results on nonlinear fractional differential equations, Philos. Trans. R. Soc. A, Math. Phys. Eng. Sci. 371(1990), Article ID 20120144 (2013).
4. A. Branciari, A fixed point theorem of Banach-Caccioppoli type on a class of generalized metric spaces, Publicationes Mathematicae Debrecen, 57(1) (2000)31-37.
5. E. Karapinar, Discussion on  $\alpha$ - $\psi$  contractions on generalized metric spaces, Abstract and Applied Analysis, vol. 2014, Article ID 962784, 7 pages.
6. I. M. Erhan, E. Karapinar, and T Sekulic, "Fixed point of  $(\psi-\Phi)$  – contractions on rectangular metric spaces," Fixed Point Theory and Applications, Vol. 22012, article 138, 2012.
7. G. V. R. Babu, P. D. Sailaja. A Fixed point theorem of generalized weak contractive maps in orbitally complete metric spaces. Thai Journal of Mathematics, 2011, 9(1): 1-10.
8. Babu, G. V. R., Sarma, K. K. M. and Krishna, P. H., Fixed points of  $\psi$ -weak Geraghty contractions in partially ordered metric spaces, J. of Adv. Res. in Pure Math., 6(4) 2014, 9 - 23.
9. Babu, G. V. R., Sarma, K. K. M. and Krishna, P. H., Necessary and sufficient conditions for the existence of fixed points of weak generalized Geraghty contractions, Int. J. Mathematics and Scientific computing, 5(1) 2015, 31 - 38.
10. M. A. Geraghty. On contractive maps. proc. of Amer. Math. soc., 1973, 40: 604-608.
11. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, 2006 Theory and applications of fractional differeneential equations. North-Holland Mathematics Studies, vol. 204. Amsterdam, The Netherlands: Elsevier.
12. Jungck G, Commuting mappings and fixed point, Amer. Math, Monthly 1976, (83), 261-263.
13. Mahammad Arshad, Jamshaid Ahamad, and Erdal Karpinar: Some common fixed point results in rectangular metric spaces. Hindawi Publishing Corporation, International Journal of Analysis, Vol. 2013, Article ID 307234, 7 pages.
14. M. S. Khan, M. Swaleh, S. Sessa. (1984). Fixed point theorems by altering distances between the points. Bull, Aust, Math. Soc., 1, 30: pp. 205-212.
15. P. H. Krishna, V Prasad, P Mahesh and P Hemalatha, Fixed Points of  $\psi$  Geraghty contractions in rectangular metric spaces, International Journal of Management, Technology And Engineering Volume 8, Issue XII, December-2018 p.p 3048-3054.
16. Geraghty Contraction principle for digital Images: P. H. Krishna and D. A. Tatajee International Research Journal of Pure Algebra-9(9), 2019, 1-6.

**Source of Support: Nil, Conflict of interest: None Declared**

***[Copy right © 2020, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]***