



COMPUTATION OF SOME NEW STATUS NEIGHBORHOOD INDICES OF GRAPHS

V. R. KULLI*

Department of Mathematics,
Gulbarga University, Kalaburgi (Gulbarga) - 585 106, India.

(Received On: 11-05-2020; Revised & Accepted On: 04-06-2020)

ABSTRACT

In this paper, we propose the total status neighborhood index, modified vertex status neighborhood index, status neighborhood inverse degree, status neighborhood zeroth order index, F-status neighborhood index, F_1 -status neighborhood index, general vertex status neighborhood index of a graph. Also we introduce the total status neighborhood polynomial, third status neighborhood polynomial, F-status neighborhood polynomial, F_1 -status neighborhood polynomial of a graph. We compute exact formulas for complete graphs, complete bipartite graphs, wheel graphs and friendship graphs.

Keywords: status, distance, status neighborhood index, F-status neighborhood index, graph.

Mathematics Subject Classification: 05C05, 05C07, 05C12, 05C35.

1. INTRODUCTION

Throughout the paper, we consider only finite, undirected simple, connected graphs. Let $V(G)$ be the vertex set and $E(G)$ be the edge set of a graph G . The edge between the vertices u and v is denoted by uv . The degree of a vertex u is the number of vertices adjacent to u and is denoted by $d_G(u)$. The distance $d(u, v)$ between any two vertices u and v is the length of shortest path connecting u and v . The status $\sigma(u)$ of a vertex u in G is the sum of its distance from every other vertex of G . Let $N(v) = N_G(v) = \{u: uv \in E(G)\}$. Let $\sigma_n(v) = \sum_{u \in N(v)} \sigma(u)$ be the status sum of neighbor vertices.

For graph theoretic terminology, we refer the book [1].

Many distance based indices of a graph such as Wiener index [4] have been appeared in the literature. In this paper, we introduce some new status neighborhood indices of graphs.

The third or vertex status neighborhood index was introduced by Kulli in [5] and it is defined as

$$SN_3(G) = \sum_{u \in V(G)} \sigma_n(u)^2$$

Recently some variants of status neighborhood indices were studied in [6].

We introduce the following status neighborhood indices:

The modified the third or vertex status neighborhood index of a graph G is defined as

$${}^m SN_3(G) = \sum_{u \in V(G)} \frac{1}{\sigma_n(u)^2}$$

The F -status neighborhood index of a graph G is defined as

$$FSN(G) = \sum_{u \in V(G)} \sigma_n(u)^3$$

Corresponding Author: V. R. Kulli*

Department of Mathematics, Gulbarga University, Kalaburgi (Gulbarga) - 585 106, India.

The total status neighborhood index of a graph G is defined as

$$T_{sn}(G) = \sum_{u \in V(G)} \sigma_n(u).$$

The status neighborhood inverse degree of a graph G is defined as

$$SNI(G) = \sum_{u \in V(G)} \frac{1}{\sigma_n(u)}.$$

The status neighborhood zeroth order index of a graph G is defined as

$$SNZ(G) = \sum_{u \in V(G)} \frac{1}{\sqrt{\sigma_n(u)}}.$$

We continue this generalization and introduce the general third or vertex status neighborhood index of a graph G , and it is defined as

$$SN_3^a(G) = \sum_{u \in V(G)} \sigma_n(u)^a,$$

where a is a real number.

Also we introduce the F_1 -status neighborhood index of a graph G and it is defined as

$$F_1SN(G) = \sum_{uv \in E(G)} [\sigma_n(u)^2 + \sigma_n(v)^2].$$

Recently, some variants of status indices were studied, for example, in [7, 8, 9, 10, 11, 12, 13, 14, 15].

The third or vertex status neighborhood polynomial was defined by Kulli in [5], defined as

$$SN_3(G, x) = \sum_{u \in V(G)} x^{\sigma_n(u)^2}.$$

We now introduce the total status neighborhood polynomial, F -status neighborhood polynomial, F_1 -status neighborhood polynomial of a graph G , and they are defined as

$$T_{sn}(G, x) = \sum_{u \in V(G)} x^{\sigma_n(u)}.$$

$$FSN(G, x) = \sum_{u \in V(G)} x^{\sigma_n(u)^3}.$$

$$F_1SN(G, x) = \sum_{uv \in E(G)} x^{\sigma_n(u)^2 + \sigma_n(v)^2}.$$

Recently some different polynomials were studied in [16, 17, 18, 19, 20, 21].

In this paper, the modified vertex status neighborhood index, status neighborhood zeroth order index, F -status neighborhood index, F_1 -status neighborhood index, general vertex status neighborhood index of some standard graphs and friendship graphs are determined. Also the total status neighborhood polynomial, vertex status neighborhood polynomial, F_1 -status neighborhood polynomial of some standard graphs and friendship graphs are computed.

2. RESULTS FOR COMPLETE GRAPHS

Let K_n be a complete graph with n vertices and $\frac{n(n-1)}{2}$ edges.

Theorem 1: The general third or vertex status neighborhood index of a complete graph K_n is

$$SN_3^a(K_n) = n(n-1)^{2a}. \tag{1}$$

Proof: Let K_n be a complete graph with n vertices. For any vertex u of K_n , $\sigma(u) = n - 1$. Thus $\sigma_n(u) = (n - 1)^2$ for any vertex of K_n . Thus

$$SN_3^a(K_n) = \sum_{u \in V(K_n)} \sigma_n(u)^a = n(n-1)^{2a}.$$

We obtain the following results by using Theorem 1.

Corollary 1.1: Let K_n be a complete graph with K_n with n vertices. Then

$$\begin{aligned}
 \text{(i)} \quad SN_3(K_n) &= n(n-1)^4 & \text{(ii)} \quad {}^m SN_3(K_n) &= \frac{n}{(n-1)^4}. \\
 \text{(iii)} \quad FSN(K_n) &= n(n-1)^6 & \text{(iv)} \quad T_{sn}(K_n) &= n(n-1)^2. \\
 \text{(v)} \quad SNI(K_n) &= \frac{n}{(n-1)^2} & \text{(vi)} \quad SNZ(K_n) &= \frac{n}{n-1}.
 \end{aligned}$$

Proof: Put $a = 2, -2, 3, 1, -1, -1/2$ in equation (1), we obtain the desired results.

Theorem 2: The general second status neighborhood index of a complete graph K_n is

$$\text{(i)} \quad F_1SN(K_n) = n(n-1)^5. \quad \text{(ii)} \quad F_1SN(K_n, x) = \frac{n(n-1)}{2} x^{2(n-1)^4}.$$

Proof: Let K_n be a complete graph with n vertices and $\frac{n(n-1)}{2}$ edges. For any vertex u of K_n , $\sigma_n(u) = (n-1)^2$.

Therefore

$$\begin{aligned}
 \text{(i)} \quad F_1SN(K_n) &= \sum_{uv \in E(K_n)} [\sigma_n(u)^2 + \sigma_n(v)^2] = [(n-1)^4 + (n-1)^4] \frac{n(n-1)}{2} \\
 &= n(n-1)^5. \\
 \text{(ii)} \quad F_1SN(K_n, x) &= \sum_{uv \in E(K_n)} x^{\sigma_n(u)^2 + \sigma_n(v)^2} = x^{(n-1)^4 + (n-1)^4} \times \frac{n(n-1)}{2} \\
 &= \frac{n(n-1)}{2} x^{2(n-1)^4}.
 \end{aligned}$$

Theorem 3: The total status neighborhood polynomial and F -status neighborhood polynomial of a complete graph K_p are given by

$$\text{(i)} \quad T_{sn}(K_n, x) = nx^{(n-1)^2}. \quad \text{(ii)} \quad FSN(K_n, x) = nx^{(n-1)^6}.$$

Proof: Let K_n be a complete graph with n vertices. Then $\sigma_n(u) = (n-1)^2$ for any vertex u of K_n . Thus

$$\begin{aligned}
 \text{(i)} \quad T_{sn}(K_n, x) &= \sum_{u \in V(G)} x^{\sigma_n(u)} = nx^{(n-1)^2}. \\
 \text{(ii)} \quad FSN(K_n, x) &= \sum_{u \in V(G)} x^{\sigma_n(u)^3} = nx^{(n-1)^6}.
 \end{aligned}$$

3. RESULTS FOR COMPLETE BIPARTITE GRAPHS

Let $K_{p,q}$ be a complete bipartite graph with $p+q$ vertices and pq edges. For vertex set of $K_{p,q}$ can be partitioned into two independent sets V_1 and V_2 such that $u \in V_1$ and $v \in V_2$ for every edge uv in $K_{p,q}$. Therefore $d_K(u)=q$, $d_K(v)=p$, where $K=K_{p,q}$. Then $\sigma(u)=q+2p-2$ and $\sigma(v)=p+2q-2$. By calculation, we obtain $\sigma_n(u)=p(q+2p-2)$ and $\sigma_n(v)=q(p+2q-2)$. Therefore

$\sigma_n(u) \setminus u \in V(G)$	$q(p+2q-2)$	$p(q+2p-2)$
Number of edges	p	q

Table-1: Status neighborhood vertex partition of $K_{p,q}$

Theorem 4: The general vertex status neighborhood index of a complete bipartite graph $K_{p,q}$ is

$$SN_v^a(K_{p,q}) = p[q(p+2q-2)]^a + q[p(q+2p-2)]^a. \tag{2}$$

Proof: By definition and by using Table 1, we deduce

$$SN^a(K_{p,q}) = \sum_{u \in V(G)} \sigma_n(u)^a = p[q(p+2q-2)]^a + q[p(q+2p-2)]^a.$$

From Theorem 4, we establish the following results.

Corollary 4.1: Let $K_{p,q}$ be a complete bipartite graph. Then

- (i) $SN(K_{p,q}) = pq^2(p+2q-2)^2 + p^2q(q+2p-2)^2$.
- (ii) ${}^m SN(K_{p,q}) = \frac{p}{q^2(p+2q-2)^2} + \frac{q}{p^2(q+2p-2)^2}$.
- (iii) $FSN(K_{p,q}) = pq^3(p+2q-2)^3 + p^3q(q+2p-2)^3$.
- (iv) $T_{sn}(K_{p,q}) = pq(3pq-4)$.
- (v) $SNI(K_{p,q}) = \frac{p}{q(p+2q-2)} + \frac{q}{p(q+2p-2)}$.
- (vi) $SNZ(K_{p,q}) = \frac{p}{\sqrt{q(p+2q-2)}} + \frac{q}{\sqrt{p(q+2p-2)}}$.

Proof: Put $a = 2, -2, 3, 1, -1, -\sqrt{2}$ in equation (2), we obtain the desired results.

Theorem 5: Let $K_{p,q}$ be a complete bipartite graph with $p+q$ vertices and pq edges. Then

- (i) $F_1SN(K_{p,q}) = pq[q^2(p+2q-2)^2 + p^2(q+2p-2)^2]$.
- (ii) $F_1SN(K_{p,q}, x) = pqx^{q^2(p+2q-2)^2 + p^2(q+2p-2)^2}$.

Proof: We have

- (i) $F_1SN(K_{p,q}) = \sum_{uv \in E(G)} [\sigma_n(u)^2 + \sigma_n(v)^2] = pq[q^2(p+2q-2)^2 + p^2(q+2p-2)^2]$.
- (ii) $F_1SN(K_{p,q}, x) = \sum_{uv \in E(G)} x^{\sigma_n(u)^2 + \sigma_n(v)^2} = pqx^{q^2(p+2q-2)^2 + p^2(q+2p-2)^2}$.

Theorem 6: The total status neighborhood polynomial and F -status neighborhood polynomial of a complete bipartite graph $K_{p,q}$ is

- (i) $T_{sn}(K_{p,q}, x) = px^{q(p+2q-2)} + qx^{p(q+2p-2)}$
- (ii) $FSN(K_{p,q}, x) = px^{q^3(p+2q-2)^3} + qx^{p^3(q+2p-2)^3}$.

Proof: We have

- (i) $T_{sn}(K_{p,q}, x) = \sum_{u \in V(G)} x^{\sigma_n(u)} = px^{q(p+2q-2)} + qx^{p(q+2p-2)}$.
- (ii) $FSN(K_{p,q}, x) = \sum_{u \in V(G)} x^{\sigma_n(u)^3} = px^{q^3(p+2q-2)^3} + qx^{p^3(q+2p-2)^3}$.

4. RESULTS FOR WHEEL GRAPHS

A wheel graph W_n is the join of K_1 and C_n . A graph W_4 is shown in Figure 1.




Figure-1: Wheel graph W_4

A graph W_n has $n + 1$ vertices and $2n$ edges. In this graph, there are two types of status vertices as follows:

$$V_1 = \{u \in V(W_n) \mid \sigma(u) = n\}, \quad |V_1| = 1.$$

$$V_2 = \{u \in V(W_n) \mid \sigma(u) = 2n - 3\}, \quad |V_2| = n.$$

By calculation, we find that there are two types of status neighborhood vertices as given in Table 2.

$\sigma_n(u) \setminus u \in V(W_n)$	$n(2n - 3)$	$5n - 6$
Number of vertices	1	n

Table-2: Status neighborhood vertex partition of W_n

In W_n , we obtain that there are types of status edges as follows:

$$E_1 = \{uv \in E(W_n) \mid \sigma(u) = \sigma(v) = 2n - 3\}, \quad |E_1| = n.$$

$$E_2 = \{uv \in E(W_n) \mid \sigma(u) = n, \sigma(v) = 2n - 3\}, \quad |E_2| = n.$$

By calculation, in W_n , there are two types of status neighborhood edges as given in Table 3.

$\sigma_n(u), \sigma_n(v) \setminus uv \in E(W_n)$	$(5n - 6, 5n - 6)$	$(5n - 6, n(2n - 3))$
Number of edges	n	n

Table-3: Status neighborhood edge partition of W_n

Theorem 7: The general vertex status neighborhood index of a wheel graph W_n is given by

$$SN^a(W_n) = [n(2n - 3)]^a + n(5n - 6)^a. \tag{3}$$

Proof: From definition and by using Table 2, we deduce

$$SN^a(W_n) = \sum_{u \in V(W_n)} \sigma_n(u)^a = [n(2n - 3)]^a + n(5n - 6)^a.$$

We obtain the following results from Theorem 7.

Corollary 7.1: Let W_n be a wheel graph with $n + 1$ vertices and $2n$ edges. Then

- (i) $SN(W_n) = 4n^4 + 13n^3 - 51n^2 + 36n.$
- (ii) ${}^m SN(W_n) = \frac{1}{n^2(2n - 3)^2} + \frac{n}{(5n - 6)^2}.$
- (iii) $FSN(W_n) = n^3(2n - 3)^3 + n(5n - 6)^3.$
- (iv) $T_{sn}(W_n) = 7n^2 - 9n.$
- (v) $SNI(W_n) = \frac{1}{n(2n - 3)} + \frac{n}{5n - 6}.$
- (vi) $SNZ(W_n) = \frac{1}{\sqrt{n(2n - 3)}} + \frac{n}{\sqrt{5n - 6}}.$

Proof: Put $a = 2, -2, 3, 1, -1, -\frac{1}{2}$ in equation (3), we obtain the desired results.

Theorem 8: The F_1 -status neighborhood index and F_1 -status neighborhood polynomial of a wheel graph W_n are given by

- (i) $F_1SN(W_n) = 4n^5 - 12n^4 + 84n^3 - 180n^2 + 108n.$
- (ii) $F_1SN(W_n, x) = nx^{50n^2 - 120n + 72} + nx^{4n^4 - 12n^3 + 34n^2 - 60n + 36}.$

Proof:

(i) By definition and by using Table 3, we derive

$$\begin{aligned} F_1SN(W_n) &= \sum_{uv \in E(W_n)} [\sigma_n(u)^2 + \sigma_n(v)^2] \\ &= n[(5n - 6)^2 + (5n - 6)^2] + n[(5n - 6)^2 + (2n^2 - 3n)^2] \\ &= 4n^5 - 12n^4 + 84n^3 - 180n^2 + 108n. \end{aligned}$$

(ii) From definition and by using Table 3, we have

$$\begin{aligned} F_1SN(W_n, x) &= \sum_{uv \in E(W_n)} x^{\sigma_n(u)^2 + \sigma_n(v)^2} \\ &= nx^{(5n-6)^2 + (5n-6)^2} + nx^{(5n-6)^2 + (2n^2-3n)^2} \\ &= nx^{50n^2-120n+72} + nx^{4n^4-12n^3+34n^2-60n+36} \end{aligned}$$

Theorem 9: The total status neighborhood polynomial and F -status neighborhood polynomial of a wheel graph W_n are given by

- (i) $T_{sn}(W_n, x) = x^{n(2n-3)} + nx^{5n-6}$.
- (ii) $FSN(W_n, x) = x^{n^3(2n-3)^3} + nx^{(5n-6)^3}$.

Proof:

(i) By definition and by using Table 2, we obtain

$$T_{sn}(W_n, x) = \sum_{u \in V(W_n)} x^{\sigma_n(u)} = x^{n(2n-3)} + nx^{5n-6}$$

(ii) From definition and by using Table 2, we have

$$FSN(W_n, x) = \sum_{u \in V(W_n)} x^{\sigma_n(u)^3} = x^{n^3(2n-3)^3} + nx^{(5n-6)^3}$$

5. RESULTS FOR FRIENDSHIP GRAPHS

A friendship graph F_n is the graph obtained by taking $n \geq 2$ copies of C_3 with vertex in common. A graph F_4 is shown in Figure 2.




Figure-2: Friendship graph F_4 .

A friendship graph F_n has $2n+1$ vertices and $3n$ edges. In F_n , we obtain two types of status vertices as follows:

$$\begin{aligned} V_1 &= \{u \in V(F_n) \mid \sigma(u) = 2n\}, & |V_1| &= 1. \\ V_2 &= \{u \in V(F_n) \mid \sigma(u) = 4n - 2\}, & |V_2| &= 2n. \end{aligned}$$

By calculation, there are two types of status neighborhood vertices in F_n as given in Table 4.

$\sigma_n(u) \setminus u \in V(F_n)$	$2n(4n - 2)$	$6n - 2$
Number of vertices	1	$2n$

Table-4: Status neighborhood vertex partition of F_n

In a graph F_n there are two types of status edges as follows:

$$\begin{aligned} E_1 &= \{uv \in E(F_n) \mid \sigma(u) = \sigma(v) = 4n - 2\}, & |E_1| &= n. \\ E_2 &= \{uv \in E(F_n) \mid \sigma(u) = 2n, \sigma(v) = 4n - 2\}, & |E_2| &= 2n. \end{aligned}$$

By calculation, we have two types of status neighborhood edges in F_n as given in Table 5.

$\sigma_n(u), \sigma_n(v) \setminus uv \in E(F_n)$	$(6n - 2, 6n - 2)$	$(6n - 2, 2n(4n - 2))$
Number of edges	n	$2n$

Table-5: Status neighborhood edge partition of F_n

Theorem 10: The general vertex status neighborhood index of a friendship graph F_n is given by

$$SN^a(F_n) = [2n(4n-2)]^a + 2n(6n-2)^a. \tag{4}$$

Proof: From definition and by using Table 4, we deduce

$$SN^a(F_n) = \sum_{u \in V(F_n)} \sigma_n(u)^a = [2n(4n-2)]^a + 2n(4n-2)^a.$$

We establish the following results by using Theorem 10.

Corollary 10.1: Let F_n be a friendship graph with $2n+1$ vertices and $3n$ edges. Then

- (i) $S(F_n) = 64n^4 + 8n^3 - 32n^2 + 8n.$
- (ii) ${}^m SN(F_n) = \frac{1}{4n^2(4n-2)^2} + \frac{n}{2(3n-1)^2}.$
- (iii) $FNS(F_n) = 8n^3(4n-2)^3 + 2n(6n-2)^3.$
- (iv) $T_{sn}(F_n) = 20n^2 - 8n.$
- (v) $SNI(F_n) = \frac{1}{2n(4n-2)} + \frac{n}{3n-1}.$
- (vi) $SNZ(F_n) = \frac{1}{2\sqrt{n(n-1)}} + \frac{2n}{\sqrt{6n-2}}.$

Proof: Put $a = 2, -2, 3, 1, -1, -1/2$ in equation (4), we obtain the desired results.

Theorem 11: The F_1 -status neighborhood index and F_1 -status neighborhood polynomial of a friendship graph F_n are given by

- (i) $F_1SN(F_n) = 128n^5 - 128n^4 + 176n^3 - 96n^2 + 16n.$
- (ii) $F_1SN(F_n, x) = nx^{2(6n-2)^2} + 2nx^{(6n-2)^2 + (8n^2-4n)^2}.$

Proof:

(i) By definition and by using Table 5, we deduce

$$\begin{aligned} F_1SN(F_n) &= \sum_{uv \in E(F_n)} [\sigma_n(u)^2 + \sigma_n(v)^2] \\ &= n[(6n-2)^2 + (6n-2)^2] + 2n[(6n-2)^2 + (8n^2-4n)^2] \\ &= 128n^5 - 128n^4 + 176n^3 - 96n^2 + 16n. \end{aligned}$$

(ii) By using definition and Table 5, we derive

$$\begin{aligned} F_1SN(F_n, x) &= \sum_{uv \in E(F_n)} x^{\sigma_n(u)^2 + \sigma_n(v)^2} \\ &= nx^{(6n-2)^2 + (6n-2)^2} + 2nx^{(6n-2)^2 + (8n^2-4n)^2} \\ &= nx^{2(6n-2)^2} + 2nx^{(6n-2)^2 + (8n^2-4n)^2} \end{aligned}$$

Theorem 12: The total status neighborhood polynomial and F -status neighborhood polynomial of a friendship graph F_n are given by

- (i) $T_{sn}(F_n, x) = x^{2n(4n-2)} + 2nx^{6n-2}.$
- (ii) $FNS(F_n, x) = x^{8n^3(4n-2)^3} + 2nx^{(6n-2)^3}.$

Proof:

(i) By using definition and Table 4, we obtain

$$T_{sn}(F_n, x) = \sum_{u \in V(F_n)} x^{\sigma_n(u)} = x^{2n(4n-2)} + 2nx^{6n-2}.$$

(ii) From definition and by using Table 4, we deduce

$$FNS(F_n, x) = \sum_{u \in V(F_n)} x^{\sigma_n(u)^3} = x^{8n^3(4n-2)^3} + 2nx^{(6n-2)^3}.$$

REFERENCES

1. V.R.Kulli, *Collegiate Graph Theory*, Vishwa International Publications, Gulbarga, India (2012).
2. H. Wiener, Structural determination of paraffin boiling points, *J. Amer. Chem. Soc.* 69(1947) 17-20.
3. V.R.Kulli, Computation of status indices of graphs, *International Journal of Mathematics Trends and Technology*, 65(12) (2019) 54-61.
4. D.Plavsic, S. Nikolić and N. Trinajstić, On the Harary index for the characterization of chemical graphs, *J. Math. Chem.*, 12 (1993) 235-250.
5. V.R.Kulli, Computation of status neighborhood indices of graphs, *International Journal of Recent Scientific Research*, 11 (4) (2020) 38079-38085..
6. V.R.Kulli, Distance based connectivity status neighborhood indices of certain graphs, *International Journal of Mathematical Archive*, 11(6) (2020) 17-23.
7. V.R.Kulli, Some new status indices of graphs, *International Journal of Mathematics Trends and Technology*, 65(10) (2019) 70-76.
8. V.R.Kulli, Computation of multiplicative (a, b)-status index of certain graphs, *Journal of Mathematics and Informatics* 18 (2020) 45-50.
9. V.R.Kulli, Some new multiplicative status indices of graphs, *International Journal of Recent Scientific Research*, 10, 10(F) (2019) 35568-35573.
10. V.R.Kulli, Status Gourava indices of graphs, *International Journal of Recent Scientific Research*, 11, 1(A) (2020) 36770-36773.
11. V.R.Kulli, Multiplicative ABC, GA, AG, augmented and harmonic status indices of graphs, *International Journal of Mathematical Archive*, 11(1) (2020) 32-40.
12. V.R.Kulli, Computation of ABC, AG and augmented status indices of graphs, *International Journal of Mathematics Trends and Technology*, 66(1) (2020) 1-7.
13. V.R.Kulli, Computation of multiplicative status indices of graphs, *International Journal of Mathematical Archive*, 11(4) (2020) 1-6.
14. H.S.Ramane, B. Basavanagoud and A.S. Yalnaik, Harmonic status index of graphs, *Bulletin of Mathematical Sciences and Applications*, 17(2016) 24-32.
15. K.P. Narayankar and D.Selvan, Geometric-arithmetic index of graphs, *International Journal of Mathematical Archive*, 8(7) (2017) 230-233.
16. V.R.Kulli, Square reverse index and its polynomial of certain networks, *International Journal of Mathematical Archive*, 9(10) (2018) 27-33.
17. V.R.Kulli, Computing square Revan index and its polynomial of certain benzenoid systems, *International Journal of Mathematics Archive*, 9(12) (2018) 41-49.
18. V.R.Kulli, On KV indices and their polynomials of two families of dendrimers, *International Journal of Current Research in Life Sciences*, 7(9) (2018) 2739-2744.
19. V.R. Kulli, Computing F-reverse index and F-reverse polynomial of certain networks, *International Journal of Mathematical Archive*, 9(8) (2018) 27-33.
20. V.R.Kulli, On augmented leap index and its polynomial of some wheel type graphs, *International Research Journal of Pure Algebra*, 9(4) (2019) 1-7.
21. V.R. Kulli, Minus F and square F-indices and their polynomials of certain dendrimers, *Earthline Journal of Mathematical Sciences*, 1(2) (2019) 171-185.

Source of Support: Nil, Conflict of interest: None Declared

[Copy right © 2020, RJPA. All Rights Reserved. This is an Open Access article distributed under the terms of the International Research Journal of Pure Algebra (IRJPA), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.]